

Reference List

Excluded: Not about screening program

Aalto-Setälä T, Marttunen M, Tuulio-Henriksson A, Poikolainen K, Lonnqvist J. One-month prevalence of depression and other DSM-IV disorders among young adults. *Psychol Med*. 2001 Jul;31(5):791-801.

Aben I, Verhey F, Strik J, Lousberg R, Lodder J, Honig A. A comparative study into the one year cumulative incidence of depression after stroke and myocardial infarction. *J Neurol Neurosurg Ps*. 2003;74(5):581-5.

Abolfotouh MA, Daffallah AA, Khan MY, Khattab MS, Abdulmoneim I. Psychosocial assessment of geriatric subjects in Abha City, Saudi Arabia. *East Mediterr Health J*. 2001 May;7(3):481-91.

Abou-Saleh MT, Ghubash R, Karim L, Krymski M, Bhai I. Hormonal aspects of postpartum depression. *Psychoneuroendocrinol*. 1998 Jul;23(5):465-75.

Abou-Saleh MT, Ghubash R, Karim L, Krymski M, Anderson DN. The role of pterins and related factors in the biology of early postpartum depression. *Eur Neuropsychopharmacol*. 1999 Jun;9(4):295-300.

Abraham IL, Wofford AB, Lichtenberg PA, Holroyd S. Factor structure of the geriatric depression scale in a cohort of depressed nursing home residents. *Int J Geriatr Psychiatry*. 1994;9(8):611-7.

Adams KB. Specific effects of caring for a spouse with dementia: Differences in depressive symptoms between caregiver and non-caregiver spouses. *Int Psychogeriatr*. 2008;20(3):508-20.

Adewuya AO, Eegunranti AB, Lawal AM. Prevalence of postnatal depression in Western Nigerian women: A controlled study. *Int J Psychiatry Clin Pract*. 2005;9(1):60-4.

Adewuya AO, Fatoye FO, Ola BA, Ijaodola OR, Ibigbami SM. Sociodemographic and obstetric risk factors for postpartum depressive symptoms in Nigerian women. *J Psychiatr Pract*. 2005 Sep;11(5):353-8.

Adewuya AO, Afolabi OT. The course of anxiety and depressive symptoms in Nigerian postpartum women. *Arch Womens Ment Health*. 2005 Nov;8(4):257-9.

Agoub M, Moussaoui D, Battas O. Prevalence of postpartum depression in a Moroccan sample. *Arch Womens Ment Health*. 2005 May;8(1):37-43.

Al-Otaibi B, Al-Weqayyan A, Taher H, Sarkhou E, Gloom A, Aseeri F, et al. Depressive symptoms among Kuwaiti population attending primary healthcare setting: Prevalence and influence of sociodemographic factors. *Med Princ Pract*. 2007;16(5):384-8.

Al-Shammari SA, Al-Subaie A. Prevalence and correlates of depression among Saudi elderly. *Int J Geriatr Psychiatry*. 1999 Sep;14(9):739-47.

Al-Windi A. Depression in general practice. *Nordic Journal of Psychiatry*. 2005;59(4):272-7.

Al Ansari A, Al Alaiwat B, Hamadeh R, Ali M, Mukarim B, Othman R. Treatment and prevalence of generalized anxiety disorder and depression among primary care attendees. *Bahrain Med Bull.* 2010;32(1).

Alderete E, Juarbe TC, Kaplan CP, Pasick R, Perez-Stable EJ. Depressive symptoms among women with an abnormal mammogram. *Psycho Oncol.* 2006 Jan;15(1):66-78.

Alexandre PK, French MT. Labor supply of poor residents in metropolitan Miami, Florida: The role of depression and the co-morbid effects of substance use. *J Ment Health Pol Econ.* 2001 Dec;4(4):161-73.

Ali NS, Hussein AA. Depression in elderly patients attending primary health care clinics in Baghdad City. *Arab J Psychiatr.* 2005 Nov;16(2):107-17.

Alim TN, Graves E, Mellman TA, Aigbogun N, Gray E, Lawson W, et al. Trauma exposure, posttraumatic stress disorder and depression in an African-American primary care population. *J Natl Med Assoc.* 2006;98(10):1630-6.

Allen RL, Walker Z, Shergill SS, D'Ath P, Katona CLE. Attitudes to depression in hospital inpatients: A comparison between older and younger subjects. *Aging Ment Health.* 1998;2(1):36-9.

Almeida-Filho N, Mari JdJ, Coutinho E, Franca JF, Fernandes J, Andreoli SB, et al. Brazilian multicentric study of psychiatric morbidity: Methodological features and prevalence estimates. *Br J Psychiatry.* 1997;171:524-9.

Alvarenga ME, Caniato RN, Mauritz A, Braun A, Aljeesh Y, Baune BT. Health service utilization in patients with major depression and co-morbid pain. *Psychiatry Clin Neurosci.* 2009 Feb;63(1):101-6.

Ames D, Tuckwell V. Psychiatric disorders among elderly patients in a general hospital. *Med J Aust.* 1994;160(11):671-5.

Amoran O, Lawoyin T, Lasebikan V. Prevalence of depression among adults in Oyo State, Nigeria: A comparative study of rural and urban communities. *Aust J Rural Health.* 2007 Jun;15(3):211-5.

Andersson L, Sundstrom-Poromaa I, Bixo M, Wulff M, Bondestam K, Astrom M. Point prevalence of psychiatric disorders during the second trimester of pregnancy: A population-based study. *Am J Obstet Gynecol.* 2003 Jul;189(1):148-54.

Andersson L, Poromaa I, Wulff M, Astrom M, Bixo M. Depression and anxiety during pregnancy and six months postpartum: A follow-up study. *Acta Obstet Gynecol Scand.* 2006;85(8):937-44.

Angst J, Gamma A, Benazzi F, Silverstein B, Ajdacic-Gross V, Eich D, et al. Atypical depressive syndromes in varying definitions. *Eur Arch Psychiatry Clin Neurosci.* 2006 Feb;256(1):44-54.

Ani C, Bazargan M, Hindman D, Bell D, Farooq MA, Akhanjee L, et al. Depression symptomatology and diagnosis: Discordance between patients and physicians in primary care settings. *BMC Fam Pract.* 2008;9:1.

Ansseau M, Dierick M, Buntinx F, Cnockaert P, De Smedt J, Van Den Haute M, et al. High prevalence of mental disorders in primary care. *J Affect Disord*. 2004 Jan;78(1):49-55.

Ansseau M, Fischler B, Dierick M, Mignon A, Leyman S. Prevalence and impact of generalized anxiety disorder and major depression in primary care in Belgium and Luxemburg: The GADIS study. *Eur Psychiatry*. 2005;20(3):229-35.

Ansseau M, Fischler B, Dierick M, Albert A, Leyman S, Mignon A. Socioeconomic correlates of generalized anxiety disorder and major depression in primary care: The GADIS II study (Generalized Anxiety and Depression Impact Survey II). *Depress Anxiety*. 2008;25(6):506-13.

Aragones E, Pinol JL, Labad A, Masdeu RM, Pino M, Cervera J. Prevalence and determinants of depressive disorders in primary care practice in Spain. *Int J Psychiatr Med*. 2004;34(1):21-35.

Aragones E, Pinol JL, Labad A. The overdiagnosis of depression in non-depressed patients in primary care. *Fam Pract*. 2006 Jun;23(3):363-8.

Aragones E, Labad A, Pinol JL, Lucena C, Alonso Y. Somatized depression in primary care attenders. *J Psychosom Res*. 2005 Feb;58(2):145-51.

Aranda MP, Lee PJ, Wilson S. Correlates of depression in older Latinos. *Home Health Care Serv Q*. 2001;20(1):1-20.

Arch JJ, Craske MG, Stein MB, Sherbourne CD, Roy-Byrne PP. Correlates of alcohol use among anxious and depressed primary care patients. *Gen Hosp Psychiatry*. 2006;28(1):37-42.

Arean PA, Alvidrez J. The prevalence of psychiatric disorders and subsyndromal mental illness in low-income, medically ill elderly. *Int J Psychiatr Med*. 2001;31(1):9-24.

Argyriadou S, Melissopoulou H, Krania E, Karagiannidou A, Vlachonicolis I, Lionis C. Dementia and depression: Two frequent disorders of the aged in primary health care in Greece. *Fam Pract*. 2001 Feb;18(1):87-91.

Aro AR, de Koning HJ, Absetz P, Schreck M. Two distinct groups of non-attenders in an organized mammography screening program. *Breast Canc Res Treat*. 2001;70(2):145-53.

Artinian NT, Washington OG, Flack JM, Hockman EM, Jen KL. Depression, stress, and blood pressure in urban African-American women. *Prog Cardiovasc Nurs*. 2006;21(2):68-75.

Asal A-R, Abdel-Fattah MM. Prevalence, symptomatology, and risk factors for depression among high school students in Saudi Arabia. *Neurosciences*. 2007;12(1):8-16.

Asarnow JR, Jaycox LH, Duan N, LaBorde AP, Rea MM, Tang L, et al. Depression and role impairment among adolescents in primary care clinics. *J Adolesc Health*. 2005 Dec;37(6):477-83.

Assadi SM, Noroozian M, Pakravannejad M, Yahyazadeh O, Aghayan S, Shariat SV, et al. Psychiatric morbidity among sentenced prisoners: Prevalence study in Iran. *Br J Psychiatry*. 2006 Feb;188(2):159-64.

Augustin A, Sahel JA, Bandello F, Dardennes R, Maurel F, Negrini C, et al. Anxiety and depression prevalence rates in age-related macular degeneration. *Investig Ophthalmol Vis Sci*. 2007;48(4):1498-503.

Augusto A, Kumar R, Calheiros JM, Matos E, Figueiredo E. Post-natal depression in an urban area of Portugal: Comparison of childbearing women and matched controls. *Psychol Med*. 1996 Jan;26(1):135-41.

Avila-Funes JA, Gray-Donald K, Payette H. Association of nutritional risk and depressive symptoms with physical performance in the elderly: The Quebec longitudinal study of nutrition as a determinant of successful aging (NuAge). *J Am Coll Nutr*. 2008 Aug;27(4):492-8.

Axon RN, Zhao Y, Egede LE. Association of depressive symptoms with all-cause and ischemic heart disease mortality in adults with self-reported hypertension. *Am J Hypertens*. 2010 Jan;23(1):30-7.

Azar AR, Murrell SA, Mast BT. Race and vascular depression risk in community-dwelling older adults. *Am J Geriatr Psychiatry*. 2005 Apr;13(4):329-32.

Baillio JN, Lyddon WJ. The self-confrontation method and the assessment of depression: A between groups comparison study. *Construct Human Sci*. 2000;5(2):89-96.

Baker FM, Parker DA, Wiley C, Velli SA, Johnson JT. Depressive symptoms in African American medical patients. *Int J Geriatr Psychiatry*. 1995;10(1):9-14.

Baker L, Cross S, Greaver L, Wei G, Lewis R, Healthy Start CORP. Prevalence of postpartum depression in a Native American population. *Matern Child Health J*. 2005 Mar;9(1):21-5.

Balestrieri M, Bisoffi G, Tansella M, Martucci M, Goldberg DP. Identification of depression by medical and surgical general hospital physicians. *Gen Hosp Psychiatry*. 2002 Jan;24(1):4-11.

Barcelos-Ferreira R, Pinto JAJ, Nakano EY, Steffens DC, Litvoc J, Bottino CMC. Clinically significant depressive symptoms and associated factors in community elderly subjects from Sao Paulo, Brazil. *Am J Geriatr Psychiatry*. 2009 Jul;17(7):582-90.

Barkin JL. The development and testing of the Barkin Index of maternal functioning. Jennifer Lynn: U Pittsburgh, US: Barkin; 2010.

Barkow K, Heun R, Bedirhan UT, Maier W. Identification of items which predict later development of depression in primary health care. *Eur Arch Psychiatry Clin Neurosci*. 2001;251(SUPPL. 2):21-6.

Barnes DM. Mental health screening in a refugee population: A program report. *J Immigrant Health*. 2001 Jul;3(3):141-9.

Barnes J, Senior R, MacPherson K. The utility of volunteer home-visiting support to prevent maternal depression in the first year of life. *Child*. 2009 Nov;35(6):807-16.

Barreto do Carmo MB, Neves Santos D, Alves Ferreira Amorim LD, Fiaccone RL, Souza da Cunha S, Cunha Rodrigues L, et al. Minor psychiatric disorders in mothers and asthma in children. *Soc Psychiatry Psychiatr Epidemiol*. 2009 May;44(5):416-20.

Barry KL, Fleming MF, Manwell LB, Copeland LA, Appel S. Prevalence of and factors associated with current and lifetime depression in older adult primary care patients. *Fam Med*. 1998;30(5):366-71.

Bartels SJ, Miles KM, Oxman TE, Zimmerman S, Cori LA, Pomerantz AS, et al. Correlates of co-occurring depressive symptoms and alcohol use in an older primary care clinic population. *J Dual Diagn*. 2006;2(3):57-72.

Bass DS, Attix DK, Phillips-Bute B, Monk TG. An efficient screening tool for preoperative depression: The Geriatric Depression Scale-Short Form. *Anesth Analg*. 2008;106(3):805-9.

Bauer HM, Rodriguez MA, Perez-Stable EJ. Prevalence and determinants of intimate partner abuse among public hospital primary care patients. *J Gen Intern Med*. 2000;15(11):811-7.

Baumeister H, Morar V. The impact of clinical significance criteria on subthreshold depression prevalence rates. *Acta Psychiatrica Scandinavica*. 2008 Dec;118(6):443-50.

Bazargan M, Ani CO, Hindman DW, Bazargan-Hejazi S, Baker RS, Bell D, et al. Correlates of complementary and alternative medicine utilization in depressed, underserved African American and Hispanic patients in primary care settings. *J Alt Complement Med*. 2008 Jun;14(5):537-44.

Beard JR, Heathcote K, Brooks R, Earnest A, Kelly B. Predictors of mental disorders and their outcome in a community based cohort. *Soc Psychiatry Psychiatr Epidemiol*. 2007;42(8):623-30.

Beck CT. Maternal depression and child behaviour problems: A meta-analysis. *J Adv Nurs*. 1999 Mar;29(3):623-9.

Beck CT, Froman RD, Bernal H. Acculturation level and postpartum depression in Hispanic mothers. *Am J Matern Child Nurs*. 2005 Sep;30(5):299-304.

Beck CT, Indman P. The many faces of postpartum depression. *JOGNN*. 2005 Sep;34(5):569-76.

Beekman AT, Deeg DJ, Braam AW, Smit JH, van Tilburg W. Consequences of major and minor depression in later life: A study of disability, well-being and service utilization. *Psychol Med*. 1997 Nov;27(6):1397-409.

Beekman AT, Penninx BW, Deeg DJ, Ormel J, Braam AW, van Tilburg W. Depression and physical health in later life: Results from the Longitudinal Aging Study Amsterdam. *J Affect Disord*. 1997 Dec;46(3):219-31.

Beekman AT, Deeg DJ, Smit JH, Comijs HC, Braam AW, de Beurs E, et al. Dysthymia in later life: A study in the community. *J Affect Disord*. 2004 Sep;81(3):191-9.

Beekman ATF, de Beurs E, van Balkom AJLM, Deeg DJH, van Dyck R, van Tilburg W. Anxiety and depression in later life: Co-occurrence and communality of risk factors. *Am J Psychiatry*. 2000 Jan;157(1):89-95.

Beghi E, Allais G, Cortelli P, D'Amico D, Simone R, d'Onofrio F, et al. Headache and anxiety-depressive disorder comorbidity: The HADAS study. *Neurol Sci*. 2007;28(Suppl 2):S217-S219.

Bell B, Chalklin L, Mills M, Browne G, Steiner M, Roberts J, et al. Burden of dysthymia and comorbid illness in adults in a Canadian primary care setting: High rates of psychiatric illness in the offspring. *J Affect Disord*. 2004 Jan;78(1):73-80.

Benazzi F. Late-life chronic depression: A 399-case study in private practice. *Int J Geriatr Psychiatry*. 2000;15(1):1-6.

Benedict RHB, Cox D, Thompson LL, Foley F, Weinstock-Guttman B, Munschauer F. Reliable screening for neuropsychological impairment in multiple sclerosis. *Mult Scler*. 2004;10(6):675-8.

Berardi D, Berti Ceroni G, Leggieri G, Rucci P, Ustun B, Ferrari G. Mental, physical and functional status in primary care attenders. *Int J Psychiatr Med*. 1999;29(2):133-48.

Berardi D, Leggieri G, Ceroni GB, Rucci P, Pezzoli A, Paltrinieri E, et al. Depression in primary care. A nationwide epidemiological survey. *Fam Pract*. 2002 Aug;19(4):397-400.

Bergdahl E, Gustavsson JM, Kallin K, von Heideken W, Lundman B, Bucht G, et al. Depression among the oldest old: The Umea 85+ study. *Int Psychogeriatr*. 2005 Dec;17(4):557-75.

Bergdahl E, Allard P, Lundman B, Gustafson Y. Depression in the oldest old in urban and rural municipalities. *Aging Ment Health*. 2007 Sep;11(5):570-8.

Bergdahl E, Allard P, Alex L, Lundman B, Gustafson Y. Gender differences in depression among the very old. *Int Psychogeriatr*. 2007 Dec;19(6):1125-40.

Bermejo I, Frey C, Kriston L, Schneider F, Gaebel W, Hegerl U, et al. Stability of the effects of guideline training in primary care on the identification of depressive disorders. *Prim Care Community Psychiatr*. 2007;12(3-4):99-107.

Bern-Klug M, Kramer KW, Sharr P. Depression screening in nursing homes: Involvement of social services departments. *Am J Geriatr Psychiatr*. 2010 Mar;18(3):266-75.

Bernstein IH, Rush AJ, Yonkers K, Carmody TJ, Woo A, McConnell K, et al. Symptom features of postpartum depression: Are they distinct? *Depress Anxiety*. 2008;25(1):20-6.

Bhagwanjee A, Parekh A, Paruk Z, Petersen I, Subedar H. Prevalence of minor psychiatric disorders in an adult African rural community in South Africa. *Psychol Med*. 1998 Sep;28(5):1137-47.

Bhui K, Bhugra D, Goldberg D, Sauer J, Tylee A. Assessing the prevalence of depression in Punjabi and English primary care attenders: The role of culture, physical illness and somatic symptoms. *Transcult Psychiatr*. 2004;41(3):307-22.

Biderman A, Cwikel J, Fried AV, Galinsky D. Depression and falls among community dwelling elderly people: A search for common risk factors. *J Epidemiol Community Health*. 2002 Aug;56(8):631-6.

Bisserbe J-C, Weiller E, Boyer P, Lepine J-P, Lecrubier Y. Social phobia in primary care: Level of recognition and drug use. *Int Clin Psychopharmacol*. 1996;11(Suppl 3):Jun-28.

Black SA, Markides KS, Miller TQ. Correlates of depressive symptomatology among older community-dwelling Mexican Americans: The Hispanic EPESE. *J Gerontol B Psychol Sci Soc Sci*. 1998 Jul;53(4):S198-S208.

Blacker CV, Thomas JM, Thompson C. Seasonality prevalence and incidence of depressive disorder in a general practice sample: Identifying differences in timing by caseness. *J Affect Disord*. 1997 Mar;43(1):41-52.

Bodlund O, Andersson SO, Mallon L. Effects of consulting psychiatrist in primary care. 1-year follow-up of diagnosing and treating anxiety and depression. *Scand J Prim Health Care*. 1999 Sep;17(3):153-7.

Bogner HR, Ford DE, Gallo JJ. The role of cardiovascular disease in the identification and management of depression by primary care physicians. *Am J Geriatr Psychiatr*. 2006 Jan;14(1):71-8.

Bolkhir A, Loisel MM, Evon DM, Hayashi PH. Depression in primary caregivers of patients listed for liver or kidney transplantation. *Prog Transplant*. 2007;17(3):193-8.

Bosch JA, Engeland CG, Cacioppo JT, Marucha PT. Depressive symptoms predict mucosal wound healing. *Psychosom Med*. 2007;69(7):597-605.

Bossuyt PMM, Raaymakers TWM, Bonsel GJ, Rinkel GJE. Screening families for intracranial aneurysms: Anxiety, perceived risk, and informed choice. *Prev Med*. 2005;41(3-4):795-9.

Boudreaux ED, Cagande C, Kilgannon H, Kumar A, Camargo J. A prospective study of depression among adult patients in an urban emergency department. *Prim Care Comp J Clin Psychiatr*. 2006;8(2):66-70.

Boyd RC, Zayas LH, McKee MD. Mother-infant interaction, life events and prenatal and postpartum depressive symptoms among urban minority women in primary care. *Matern Child Health J*. 2006 Mar;10(2):139-48.

Brandon AR, Trivedi MH, Hynan LS, Miltenberger PD, Labat DB, Rifkin JB, et al. Prenatal depression in women hospitalized for obstetric risk. *J Clin Psychiatr*. 2008;69(4):635-43.

Breen SJ, Baravelli CM, Schofield PE, Jefford M, Yates PM, Aranda SK. Is symptom burden a predictor of anxiety and depression in patients with cancer about to commence chemotherapy? *Med J Aust*. 2009;190(7 SUPPL.):S99-S104.

Brilman EI, Ormel J. Life events, difficulties and onset of depressive episodes in later life. *Psychol Med*. 2001 Jul;31(5):859-69.

Brindle LA, Oliver SE, Dedman D, Donovan JL, Neal DE, Hamdy FC, et al. Measuring the psychosocial impact of population-based prostate-specific antigen testing for prostate cancer in the UK. *BJU Int*. 2006;98(4):777-82.

Britton JR. Maternal anxiety: Course and antecedents during the early postpartum period. *Depress Anxiety*. 2008 Sep;25(9):793-800.

Brodaty H, Kemp NM, Low L-F. Characteristics of the GPCOG, a screening tool for cognitive impairment. *Int J Geriatr Psychiatry*. 2004;19(9):870-4.

Brody DS, Dietrich AJ, deGruy F, III, Kroenke K. The depression in primary care tool kit. *Int J Psychiatr Med*. 2000;30(2):99-110.

Brook OH, van Hout H, Stalman W, Nieuwenhuysen H, Bakker B, Heerdink E, et al. A pharmacy-based coaching program to improve adherence to antidepressant treatment among primary care patients. *Psychiatr Serv*. 2005 Apr;56(4):487-9.

Bruce DG, Casey GP, Grange V, Clarnette RC, Almeida OP, Foster JK, et al. Cognitive impairment, physical disability and depressive symptoms in older diabetic patients: The Fremantle Cognition in Diabetes Study. *Diabetes Res Clin Pract*. 2003;61(1):59-67.

Bruce ML, Seeman TE, Merrill SS, Blazer DG. The impact of depressive symptomatology on physical disability: MacArthur Studies of Successful Aging. *Am J Public Health*. 1994 Nov;84(11):1796-9.

Brugha TS, Wheatley S, Taub NA, Culverwell A, Friedman T, Kirwan P, et al. Pragmatic randomized trial of antenatal intervention to prevent post-natal depression by reducing psychosocial risk factors. *Psychol Med*. 2000 Nov;30(6):1273-81.

Buist AE, Austin MP, Hayes BA, Speelman C, Bilszta JL, Gemmill AW, et al. Postnatal mental health of women giving birth in Australia 2002-2004: Findings from the beyondblue National Postnatal Depression Program. *Aust NZ J Psychiatry*. 2008 Jan;42(1):66-73.

Bulloch AG, Patten SB. Non-remission of depression in the general population as assessed by the HAMD-7 scale. *Depress Anxiety*. 2008;25(5):393-7.

Bunevicius R, Kusminskas L, Bunevicius A, Nadisauskiene RJ, Jureniene K, Pop VJ. Psychosocial risk factors for depression during pregnancy. *Acta Obstet Gynecol Scand*. 2009;88(5):599-605.

Burman ME, McCabe S, Pepper CM. Treatment practices and barriers for depression and anxiety by primary care advanced practice nurses in Wyoming. *J Am Acad Nurse Pract*. 2005 Sep;17(9):370-80.

Cairney J, Corna LM, Veldhuizen S, Herrmann N, Streiner DL. Comorbid depression and anxiety in later life: Patterns of association, subjective well-being, and impairment. *Am J Geriatr Psychiatry*. 2008 Mar;16(3):201-8.

Callahan CM, Hendrie HC, Dittus RS, Brater DC, Hui SL, Tierney WM. Depression in late life: The use of clinical characteristics to focus screening efforts. *J Gerontol*. 1994 Jan;49(1):M9-14.

Callahan CM, Wolinsky FD, Stump TE, Nienaber NA, Hui SL, Tierney WM. Mortality, symptoms, and functional impairment in late-life depression. *J Gen Intern Med*. 1998 Nov;13(11):746-52.

Callahan EJ, Bertakis KD, Azari R, Robbins J, Helms LJ, Miller J. The influence of depression on physician-patient interaction in primary care. *Fam Med*. 1996 May;28(5):346-51.

Callen BL, Wells TJ. Screening for nutritional risk in community-dwelling old-old. *Public Health Nurs*. 2005;22(2):138-46.

Campbell A, Hayes B, Buckby B. Aboriginal and Torres Strait Islander women's experience when interacting with the Edinburgh Postnatal Depression Scale: A brief note. *Aust J Rural Health*. 2008 Jun;16(3):124-31.

Campbell DG, Felker BL, Liu CF, Yano EM, Kirchner JE, Chan D, et al. Prevalence of depression-PTSD comorbidity: Implications for clinical practice guidelines and primary care-based interventions. *J Gen Intern Med*. 2007 Jun;22(6):711-8.

Cannon-Spoor HE, Levy JA, Zubenko GS, Zubenko WW, Cohen RM, Mirza N, et al. Effects of previous major depressive illness on cognition in Alzheimer disease patients. *Am J Geriatr Psychiatry*. 2005;13(4):312-8.

Cantilino A, Zambaldi CF, De Albuquerque TLC, Paes JA, Montenegro ACP, Sougey EB. Postpartum depression in Recife - Brazil: Prevalence and association with bio-socio-demographic factors. *Jornal Brasileiro de Psiquiatria*. 2010;59(1):1-9.

Cantwell R, Cox JL. Psychiatric disorders in pregnancy and the puerperium. *Curr Obstet Gynaecol*. 2003;13(1):7-13.

Caplan S, Alvidrez J, Paris M, Escobar JI, Dixon JK, Desai MM, et al. Subjective versus objective: An exploratory analysis of latino primary care patients with self-perceived depression who do not fulfill primary care evaluation of mental disorders patient health questionnaire criteria for depression. *Prim Care Comp J Clin Psychiatry*. 2010;12(5):e1-e12.

Capoccia KL, Boudreau DM, Blough DK, Ellsworth AJ, Clark DR, Stevens NG, et al. Randomized trial of pharmacist interventions to improve depression care and outcomes in primary care. *Am J Health*. 2004 Feb 15;61(4):364-72.

Carmin CN, Klocek JW. To screen or not to screen: Symptoms identifying primary care medical patients in need of screening for depression. *Int J Psychiatr Med*. 1998;28(3):293-302.

Carta MG, Loviselli A, Hardoy MC, Massa S, Cadeddu M, Sardu C, et al. The link between thyroid autoimmunity (antithyroid peroxidase autoantibodies) with anxiety and mood disorders in the community: A field of interest for public health in the future. *BMC Psychiatry*. 2004;4 , 2004. Date of Publication: 18 Aug 2004.

Casey P, Goolsby S, Berkowitz C, Frank D, Cook J, Cutts D, et al. Maternal depression, changing public assistance, food security, and child health status. *Pediatrics*. 2004 Feb;113(2):298-304.

Casey P, Birbeck G, McDonagh C, Horgan A, Dowrick C, Dalgard O, et al. Personality disorder, depression and functioning: Results from the ODIN study. *J Affect Disord*. 2004;82(2):277-83.

Casey P, Maracy M, Kelly BD, Lehtinen V, Ayuso-Mateos J-L, Dalgard OS, et al. Can adjustment disorder and depressive episode be distinguished? Results from ODIN. *J Affect Disord*. 2006;92(2-3):291-7.

Casey P, Dunn G, Kelly BD, Lehtinen V, Dalgard OS, Dowrick C, et al. The prevalence of suicidal ideation in the general population: Results from the Outcome of Depression International Network (ODIN) study. *Soc Psychiatry Psychiatr Epidemiol*. 2008 Apr;43(4):299-304.

Castilla-Puentes RC, Secin R, Grau A, Galeno R, De Mello MF, Pena N, et al. A multicenter study of major depressive disorder among emergency department patients in Latin-American countries. *Depress Anxiety*. 2008;25(12):E199-E204.

Castro-Costa E, Dewey M, Stewart R, Banerjee S, Huppert F, Mendonca-Lima C, et al. Prevalence of depressive symptoms and syndromes in later life in ten European countries: The SHARE study. *Br J Psychiatry*. 2007;191(NOV.):393-401.

Chahine LM, Bijlsma A, Hospers APN, Chemali Z. Dementia and depression among nursing home residents in Lebanon: A pilot study. *Int J Geriatr Psychiatry*. 2007 Apr;22(4):283-5.

Chakrabhand MLS, Visanuyothin T, Bhugra D. The impact of tsunami on psychological well-being in Thailand. *Int J Cult Ment Health*. 2008 Dec;1(2):85-92.

Chan S, Levy V. Postnatal depression: A qualitative study of the experiences of a group of Hong Kong Chinese women. *J Clin Nurs*. 2004 Jan;13(1):120-3.

Chan SSM, Lam LCW, Tam CWC, Lui VWC, Chan WC, Wong S, et al. Prevalence of clinically significant depressive symptoms in an epidemiologic sample of community-dwelling elders with milder forms of cognitive impairment in Hong Kong SAR. *Int J Geriatr Psychiatry*. 2008 Jun;23(6):611-7.

Chaturvedi SK, Kalyanasundaram S, Jagadish A, Prabhu V, Narasimha V. Detection of stress, anxiety and depression in IT/ITES professionals in the Silicon Valley of India: A preliminary study. *Prim Care Community Psychiatr*. 2008;12(2):75-80.

Chaudron LH, Klein MH, Remington P, Palta M, Allen C, Essex MJ. Predictors, prodromes and incidence of postpartum depression. *J Psychosom Obstet Gynaecol*. 2001 Jun;22(2):103-12.

Chaudron LH, Kitzman HJ, Szilagyi PG, Sidora-Arcoleo K, Anson E. Changes in maternal depressive symptoms across the postpartum year at well child care visits. *Ambul Pediatr*. 2006;6(4):221-4.

Cheasty M, Clare AW, Collins C. Child sexual abuse - A predictor of persistent depression in adult rape and sexual assault victims. *J Ment Health*. 2002;11(1):79-84.

Chee CY, Lee DT, Chong YS, Tan LK, Ng TP, Fones CS. Confinement and other psychosocial factors in perinatal depression: A transcultural study in Singapore. *J Affect Disord*. 2005 Dec;89(1-3):157-66.

Chee CYI, Chong Y-S, Ng TP, Lee DTS, Tan LK, Fones CSL. The association between maternal depression and frequent non-routine visits to the infant's doctor - A cohort study. *J Affect Disord*. 2008;107(1-3):247-53.

Chen CH, Wang SY, Chung UL, Tseng YF, Chou FH. Being reborn: The recovery process of postpartum depression in Taiwanese women. *J Adv Nurs*. 2006 May;54(4):450-6.

Chen HC, Chou FH, Chen MC, Su SF, Wang SY, Feng WW, et al. A survey of quality of life and depression for police officers in Kaohsiung, Taiwan. *Qual Life Res*. 2006 Jun;15(5):925-32.

Chiriboga DA, Jang Y, Banks S, Kim G. Acculturation and its effect on depressive symptom structure in a sample of Mexican American elders. *Hispanic J Behav Sci*. 2007 Feb;29(1):83-100.

Chisholm D, Diehr P, Knapp M, Patrick D, Treglia M, Simon G, et al. Depression status, medical comorbidity and resource costs. Evidence from an international study of major depression in primary care (LIDO). *Br J Psychiatry*. 2003 Aug;183:121-31.

Chong J, Reinschmidt KM, Moreno FA. Symptoms of depression in a Hispanic primary care population with and without chronic medical illnesses. *Prim Care Comp J Clin Psychiatr*. 2010;12(3).

Chou FH-C, Wu HC, Chou P, Su CY, Tsai KY, Chao SS, et al. Epidemiologic psychiatric studies on post-disaster impact among Chi-Chi earthquake survivors in Yu-Chi, Taiwan. *Psychiatry Clin Neurosci*. 2007 Aug;61(4):370-8.

Chung KF, Tso KC, Cheung E, Wong M. Validation of the Chinese version of the Mood Disorder Questionnaire in a psychiatric population in Hong Kong. *Psychiatry Clin Neurosci*. 2008 Aug;62(4):464-71.

Cianciarullo Minett TSC, Da Silva RV, Ortiz KZ, Bertolucci PHF. Subjective memory complaints in an elderly sample: A cross-sectional study. *Int J Geriatr Psychiatry*. 2008;23(1):49-54.

Clarke PJ. Validation of two postpartum depression screening scales with a sample of First Nations and Metis women. *Can J Nurs Res*. 2008 Mar;40(1):113-25.

Clayer JR, McFarlane AC, Bookless CL, Air T, Wright G, Czechowicz AS. Prevalence of psychiatric disorders in rural South Australia. *Med J Aust*. 128 Sep;163(3):124-5.

Cleland JA, Lee AJ, Hall S. Associations of depression and anxiety with gender, age, health-related quality of life and symptoms in primary care COPD patients. *Fam Pract*. 2007 Jun;24(3):217-23.

Codori A-M, Slavney PR, Rosenblatt A, Brandt J. Prevalence of major depression one year after predictive testing for Huntington's disease. *Genet Testing*. 2004;8(2):114-9.

Coffman MJ, Norton CK. Demands of immigration, health literacy, and depression in recent Latino immigrants. *Home Health Care Manag Pract*. 2010 Feb;22(2):116-22.

Cohen MM, Schei B, Ansara D, Gallop R, Stuckless N, Stewart DE. A history of personal violence and postpartum depression: Is there a link? *Arch Womens Ment Health*. 2002;4(3):83-92.

Colenda CC, Legault C, Rapp SR, Debon MW, Hogan P, Wallace R, et al. Psychiatric disorders and cognitive dysfunction among older, postmenopausal women: Results from the Women's Health Initiative Memory Study. *Am J Geriatr Psychiatr*. 2010 Feb;18(2):177-86.

Comino EJ, Silove D, Manicavasagar V, Harris E, Harris MF. Agreement in symptoms of anxiety and depression between patients and GPs: The influence of ethnicity. *Fam Pract*. 2001 Feb;18(1):71-7.

Concin H, Ulmer H, Hefler L. Mental well-being in 5000 women participating in the 'Women-Plus' preventive medicine program. *Maturitas*. 2002 Apr 15;41 Suppl 1:S9-12.

Cook CAL, Flick LH, Homan SM, Campbell C, McSweeney M, Gallagher ME. Psychiatric disorders and treatment in low-income pregnant women. *J Womens Health*. 2010 Jul;19(7):1251-62.

Cooler CY, Finkel SI, Moran MB, Richter EM. Recruitment and enrollment of elderly persons with depression into a clinical drug trial. *Int J Geriatr Psychopharmacol*. 1999;2(1):23-7.

Coryell W, Akiskal HS, Leon AC, Winokur G, Maser JD, Mueller TI, et al. The time course of nonchronic major depressive disorder. Uniformity across episodes and samples. National Institute of Mental Health Collaborative Program on the Psychobiology of Depression--Clinical Studies. *Arch Gen Psychiatry*. 1994 May;51(5):405-10.

Costa E, Barreto SM, Uchoa E, Firmo JO, Lima-Costa MF, Prince M. Prevalence of International Classification of Diseases, 10th Revision common mental disorders in the elderly in a Brazilian community: The Bambui Health Ageing Study. *Am J Geriatr Psychiatr*. 2007 Jan;15(1):17-27.

Coyne JC, Fechner-Bates S, Schwenk TL. Prevalence, nature, and comorbidity of depressive disorders in primary care. *Gen Hosp Psychiatry*. 1994 Jul;16(4):267-76.

Coyne JC, Thompson R, Racioppo MW. Validity and efficiency of screening for history of depression by self-report. *Psychol Assess*. 2001;13(2):163-70.

Coyne JC, Brown G, Datto C, Bruce ML, Schulberg HC, Katz I. The benefits of a broader perspective in case-finding for disease management of depression: Early lessons from the PROSPECT study. *Int J Geriatr Psychiatry*. 2001 Jun;16(6):570-6.

Creed F, Winterbottom M, Tomenson B, Britt R, Anand IS, Wander GS, et al. Preliminary study of non-psychotic disorders in people from the Indian subcontinent living in the UK and India. *Acta Psychiatrica Scandinavica*. 1999 Apr;99(4):257-60.

Da Canhota CM, Piterman L. Depressive disorders in elderly Chinese patients in Macau: A comparison of general practitioners' consultations with a depression screening scale. *Aust NZ J Psychiatry*. 2001 Jun;35(3):336-44.

da Silva RA, Jansen K, Souza LDDM, Moraes IGDS, Tomasi E, Da Silva GDG, et al. Depression during pregnancy in the Brazilian public health care system. *Rev Bras Psiquiatr*. 2010;32(2):139-44.

Dahlberg KM, Waern M, Runeson B. Mental health literacy and attitudes in a Swedish community sample - Investigating the role of personal experience of mental health care. *BMC Public Health*. 2008;8, 2008. Article Number: 8. Date of Publication: 2008.

Dal FG, Palermo MT, Donohue JE, Karagiozis H, Zonderman AB, Kawas CH. Depressive symptoms, sex, and risk for Alzheimer's disease. *Ann Neurol*. 2005;57(3):381-7.

Daly RC, Danaceau MA, Rubinow DR, Schmidt PJ. Concordant restoration of ovarian function and mood in perimenopausal depression. *Am J Psychiatry*. 2003;160(10):1842-6.

Dao TK, Youssef NA, Gopaldas RR, Chu D, Bakaeen F, Wear E, et al. Autonomic cardiovascular dysregulation as a potential mechanism underlying depression and coronary artery bypass grafting surgery outcomes. *J Cardiothorac Surg*. 2010;5:36.

Dave S, Nazareth I, Sherr L, Senior R. The association of paternal mood and infant temperament: A pilot study. *Br J Dev Psychol*. 2005 Nov;23(4):609-21.

Davila M, McFall SL, Cheng D. Acculturation and depressive symptoms among pregnant and postpartum Latinas. *Matern Child Health J*. 2009 May;13(3):318-25.

Davis TM, Bush KR, Kivlahan DR, Dobie DJ, Bradley KA. Screening for substance abuse and psychiatric disorders among women patients in a VA health care system. *Psychiatr Serv*. 2003;54(2):214-8.

Deave T, Heron J, Evans J, Emond A. The impact of maternal depression in pregnancy on early child development. *BJOG*. 2008 Jul;115(8):1043-51.

Deave T, Heron J, Evans J, Emond A. The impact of maternal depression in pregnancy on early child development. *Obstetrical and Gynecological Survey*. 2008;63(10):626-8.

Deb S, Thomas M, Bright C. Mental disorder in adults with intellectual disability. I: Prevalence of functional psychiatric illness among a community-based population aged between 16 and 64 years. *J Intellect Disabil Res*. 2001 Dec;45(6):495-505.

Delatte R, Cao H, Meltzer-Brody S, Menard MK. Universal screening for postpartum depression: An inquiry into provider attitudes and practice. *Am J Obstet Gynecol*. 2009 May;200(5):e63-e64.

Dennis C-L, Ross LE. The clinical utility of maternal self-reported personal and familial psychiatric history in identifying women at risk for postpartum depression. *Acta Obstet Gynecol Scand*. 2006;85(10):1179-85.

Dennis CL. Influence of depressive symptomatology on maternal health service utilization and general health. *Arch Womens Ment Health*. 2004 Jul;7(3):183-91.

Dickens C, McGowan L, Percival C, Tomenson B, Cotter L, Heagerty A, et al. Negative illness perceptions are associated with new-onset depression following myocardial infarction. *Gen Hosp Psychiatry*. 2008 Sep;30(5):414-20.

Dickens CM, McGowan L, Percival C, Douglas J, Tomenson B, Cotter L, et al. Lack of a close confidant, but not depression, predicts further cardiac events after myocardial infarction. *Heart*. 2004;90(5):518-22.

Dindar I, Erdogan S. Screening of Turkish women for postpartum depression within the first postpartum year: The risk profile of a community sample. *Public Health Nurs*. 2007 Mar;24(2):176-83.

Dogar IA, Azeem MW, Kiran M, Hussain I, Mehmood K, Hina I. Depression and anxiety in cancer patients in outpatient department of a tertiary care hospital in Pakistan. *Pakistan J Med Sci*. 2009;25(5):734-7.

Douglas KM, Taylor AJ, O'Malley PG. Relationship between depression and C-reactive protein in a screening population. *Psychosom Med*. 2004;66(5):679-83.

Dowdy DW, Dinglas V, Mendez-Tellez PA, Bienvenu OJ, Sevransky J, Dennison CR, et al. Intensive care unit hypoglycemia predicts depression during early recovery from acute lung injury. *Crit Care Med*. 2008 Oct;36(10):2726-33.

Downey G, Coyne JC. Children of depressed parents: An integrative review. *Psychol Bull*. 1990 Jul;108(1):50-76.

Driscoll KA, Johnson SB, Barker D, Quittner AL, Deeb LC, Geller DE, et al. Risk factors associated with depressive symptoms in caregivers of children with type 1 diabetes or cystic fibrosis. *J Pediatr Psychol*. 2010 Sep;35(8):814-22.

Dudley M, Roy K, Kelk N, Bernard D. Psychological correlates of depression in fathers and mothers in the first postnatal year. *J Reprod Infant Psychol*. 2001;19(3):187-202.

Duffy SA, Terrell JE, Valenstein M, Ronis DL, Copeland LA, Connors M. Effect of smoking, alcohol, and depression on the quality of life of head and neck cancer patients. *Gen Hosp Psychiatry*. 2002;24(3):140-7.

Dunham NC, Sager MA. Functional status, symptoms of depression, and the outcomes of hospitalization in community-dwelling elderly patients. *Arch Fam Med*. 1996;3(8):676-80.

Dunn LL, Hundley MC, Shelton MM. Spiritual well-being, anxiety, and depression in antepartal women on bedrest. *Issues Ment Health Nurs*. 2007;28(11):1235-46.

Duran B, Sanders M, Skipper B, Waitzkin H, Malcoe LH, Paine S, et al. Prevalence and correlates of mental disorders among Native American women in primary care. *Am J Public Health*. 2004 Jan;94(1):71-7.

Durieux-Paillard S, Whitaker-Clinch B, Bovier PA, Eytan A. Screening for major depression and posttraumatic stress disorder among asylum seekers: Adapting a standardized instrument to the social and cultural context. [References]. *Can J Psychiatry*. 2006 Aug;51(9):587-97.

Ebeigbe PN, Akhigbe KO. Incidence and associated risk factors of postpartum depression in a tertiary hospital in Nigeria. *Nigerian Postgraduate Medical Journal*. 2008 Mar;15(1):15-8.

Edge D, Baker D, Rogers A. Perinatal depression among black Caribbean women. *Health Soc Care Community*. 2004 Sep;12(5):430-8.

Edge D. Ethnicity, psychosocial risk, and perinatal depression--A comparative study among inner-city women in the United Kingdom. *J Psychosom Res*. 2007 Sep;63(3):291-5.

Eisenberg D, Golberstein E, Gollust SE. Help-seeking and access to mental health care in a university student population. *Medical Care*. 2007 Jul;45(7):594-601.

Ekeberg O, Skjauff H, Karesen R. Screening for breast cancer is associated with a low degree of psychological distress. *Breast*. 2001;10(1):20-4.

El-Rufaie OE, Al-Sabosy MA, Bener A, Abuzeid MS. Somatized mental disorder among primary care Arab patients: I. Prevalence and clinical and sociodemographic characteristics. *J Psychosom Res*. 1999 Jun;46(6):549-55.

Ell K, Sanchez K, Vourlekis B, Lee P-J, Dwight-Johnson M, Lagomasino I, et al. Depression, correlates of depression, and receipt of depression care among low-income women with breast or gynecologic cancer. *J Clin Oncol*. 2005;23(13):3052-60.

Ell K, Katon W, Cabassa LJ, Xie B, Lee PJ, Kapetanovic S, et al. Depression and diabetes among low-income Hispanics: Design elements of a socio-culturally adapted collaborative care model randomized controlled trial. *Int J Psychiatr Med*. 2009;39(2):113-32.

Ercolani M, Trombini G, Chattat R, Cervini C, Piergiacomi G, Salaffi F, et al. Fibromyalgic syndrome: Depression and abnormal illness behavior. Multicenter investigation. *Psychother Psychosom.* 1994;61(3-4):178-86.

Ericsson M, Poston WS, Linder J, Taylor JE, Haddock CK, Foreyt JP. Depression predicts disability in long-term chronic pain patients. *Disabil Rehabil.* 2002 Apr 15;24(6):334-40.

Esler DM, Johnston F, Thomas D. The acceptability of a depression screening tool in an urban, Aboriginal community-controlled health service. *Aust NZ J Public Health.* 2007 Jun;31(3):259-63.

Evans JR, Fletcher AE, Wormald RP. Depression and anxiety in visually impaired older people. *Ophthalmology.* 2007 Feb;114(2):283-8.

Evans LM, Myers MM, Monk C. Pregnant women's cortisol is elevated with anxiety and depression - But only when comorbid. *Arch Womens Ment Health.* 2008;11(3):239-48.

Evren C, Can S, Evren B, Saatcioglu O, Cakmak D. Lifetime posttraumatic stress disorder in Turkish alcohol-dependent inpatients: Relationship with depression, anxiety and erectile dysfunction. *Psychiatry Clin Neurosci.* 2006;60(1):77-84.

Fahrenkopf AM, Sectish TC, Barger LK, Sharek PJ, Lewin D, Chiang VW, et al. Rates of medication errors among depressed and burnt out residents: Prospective cohort study. *BMJ.* 2008 Mar 1;336(7642):488-91.

Falck RS, Wang J, Carlson RG, Eddy M, Siegal HA. The prevalence and correlates of depressive symptomatology among a community sample of crack-cocaine smokers. *J Psychoactive Drugs.* 2002;34(3):281-8.

Fan F, Zou Y, Ma A, Yue Y, Mao W, Ma X. Hormonal changes and somatopsychologic manifestations in the first trimester of pregnancy and post partum. *Int J Gynaecol Obstet.* 2009 Apr;105(1):46-9.

Feijo RB, Saueressig M, Salazar C, Chaves MLF. Mental health screening by self-report questionnaire among community adolescents in southern Brazil. *J Adolesc Health.* 1997 Mar;20(3):232-7.

Felice E, Saliba J, Grech V, Cox J. Prevalence rates and psychosocial characteristics associated with depression in pregnancy and postpartum in Maltese women. *J Affect Disord.* 2004 Oct 15;82(2):297-301.

Felker B, Katon W, Hedrick SC, Rasmussen J, McKnight K, McDonnell MB, et al. The association between depressive symptoms and health status in patients with chronic pulmonary disease. *Gen Hosp Psychiatry.* 2001;23(2):56-61.

Ferketich AK, Schwartzbaum JA, Frid DJ, Moeschberger ML. Depression as an antecedent to heart disease among women and men in the NHANES I study. National Health and Nutrition Examination Survey. *Arch Intern Med.* 2000 May 8;160(9):1261-8.

Ferraro FR, Chelminski I. Preliminary normative data on the Geriatric Depression Scale-Short Form (GDS-SF) in a young adult sample. *J Clin Psychol.* 1996;52(4):443-7.

Ferreira RB, Steffens DC, De Campos Bottino CM. Clinically significant depressive symptoms and associated factors in a community sample of elderly subjects. *Alzheimers Dement.* 2009;Conference: Alzheimer's Association International Conference on Alzheimer's Disease Vienna Austria. Conference Start: 20090711 Conference End: 20090716. Conference: Alzheimer's Association International Conference on Alzheimer's Disease Vienna Austria. Conference Start: 20090711 Conference End: 20090716. Conference Publication:(var.pagings):294-5.

Ferrie JE, Shipley MJ, Davey SG, Stansfeld SA, Marmot MG. Change in health inequalities among British civil servants: The Whitehall II study. *J Epidemiol Community Health.* 2002;56(12):922-6.

Ferro T, Verdelli H, Pierre F, Weissman MM. Screening for depression in mothers bringing their offspring for evaluation or treatment of depression. *Am J Psychiatry.* 2000 Mar;157(3):375-9.

Field T, Diego M, Hernandez-Reif M, Figueiredo B, Deeds O, Contogeorgos J, et al. Prenatal paternal depression. *Infant Behav Devel.* 2006;29(4):579-83.

Fifer SK, Buesching DP, Henke CJ, Potter LP, Mathias SD, Schonfeld WH, et al. Functional status and somatization as predictors of medical offset in anxious and depressed patients. *Value Health.* 2003 Jan;6(1):40-50.

Finley PR, Rens HR, Pont JT, Gess SL, Louie C, Bull SA, et al. Impact of a collaborative care model on depression in a primary care setting: A randomized controlled trial. *Pharmacotherapy.* 2003 Sep;23(9):1175-85.

Fisher L, Skaff MM, Mullan JT, Arean P, Glasgow R, Masharani U. A longitudinal study of affective and anxiety disorders, depressive affect and diabetes distress in adults with Type 2 diabetes. *Diabetic Med.* 2008 Sep;25(9):1096-101.

Flynn HA, Davis M, Marcus SM, Cunningham R, Blow FC. Rates of maternal depression in pediatric emergency department and relationship to child service utilization. *Gen Hosp Psychiatry.* 2004 Jul;26(4):316-22.

Forkmann T, Norra C, Wirtz M, Gauggel S, Boecker M. Psychometric evaluation of the Rasch-based depression screening in patients with neurologic disorders. *Arch Phys Med Rehabil.* 2010 Aug;91(8):1188-93.

Foss GF, Chantal AW, Hendrickson S. Maternal depression and anxiety and infant development: A comparison of foreign-born and native-born mothers. *Public Health Nurs.* 2004;21(3):237-46.

Francia-Martinez M, Roca de Torres I, Alvarado CS, Martinez-Taboas A, Sayers S. Dissociation, depression and trauma in psychiatric inpatients in Puerto Rico. *J Trauma Dissociat.* 2003;4(4):47-61.

Freudenstein U, Arthur AJ, Matthews RJ, Jagger C. Community surveys of late-life depression: Who are the non-responders? *Age Ageing*. 2001 Nov;30(6):517-21.

Freyenhagen R, Baron R, Tolle T, Stemmler E, Gockel U, Stevens M, et al. Screening of neuropathic pain components in patients with chronic back pain associated with nerve root compression: A prospective observational pilot study (MIPOPT). *Curr Med Res Opin*. 2006;22(3):529-37.

Friedman B, Conwell Y, Delavan RR, Wamsley BR, Eggert GM. Depression and suicidal behaviors in Medicare primary care patients under age 65. *J Gen Intern Med*. 2005 May;20(5):397-403.

Frojd K, Hakansson A, Karlsson I, Molarius A. Deceased, disabled or depressed--A population-based 6-year follow-up study of elderly people with depression. *Soc Psychiatry Psychiatr Epidemiol*. 2003 Oct;38(10):557-62.

Fu C-W, Tan AW, Sheng F, Luan R-S, Zhan S-Y, Chen W-Q, et al. The prevalence of anxiety symptoms and depressive symptoms in patients with somatic disorders in urban China: A multi-center cross-sectional study. *Int J Psychiatr Med*. 2007;37(2):185-99.

Fuller CD, Schillerstrom JE, Jones WE, III, Boersma M, Royall DR, Fuss M. Prospective evaluation of pretreatment executive cognitive impairment and depression in patients referred for radiotherapy. *Int J Radiat Oncol Biol Phys*. 2008 Oct 1;72(2):529-33.

Gahm GA, Lucenko BA, Retzlaff P, Fukuda S. Relative impact of adverse events and screened symptoms of posttraumatic stress disorder and depression among active duty soldiers seeking mental health care. *J Clin Psychol*. 2007 Mar;63(3):199-211.

Gahm GA, Lucenko BA. Screening soldiers in outpatient care for mental health concerns. *Mil Med*. 2008 Jan;173(1):17-24.

Galeazzi GM, Ferrari S, Giaroli G, Mackinnon A, Merelli E, Motti L, et al. Psychiatric disorders and depression in multiple sclerosis outpatients: Impact of disability and interferon beta therapy. *Neurol Sci*. 2005;26(4):255-62.

Gallo JJ, Bogner HR, Morales KH, Post EP, Ten Have T, Bruce ML. Depression, cardiovascular disease, diabetes, and two-year mortality among older, primary-care patients. *Am J Geriatr Psychiatr*. 2005 Sep;13(9):748-55.

Ganatra HA, Zafar SN, Qidwai W, Rozi S. Prevalence and predictors of depression among an elderly population of Pakistan. *Aging Ment Health*. 2008 May;12(3):349-56.

Ganguli M, Gilby J, Seaberg E, Belle S. Depressive symptoms and associated factors in a rural elderly population: The MoVIES Project. *Am J Geriatr Psychiatr*. 1995;3(2):144-60.

Gao W, Paterson J, Abbott M, Carter S, Iusitini L. Maternal mental health and child behaviour problems at 2 years: Findings from the Pacific Islands Families Study. *Aust NZ J Psychiatry*. 2007 Nov;41(11):885-95.

Gaugue-Finot J, Devouche E, Wendland J, Varescon I. Prenatal depression in a French sample: Associations with psychological distress, anxiety and perceived social support. *Neuropsychiatrie de l'Enfance et de l'Adolescence*. 2010;58(8):441-7.

Gawronski I, Privette G. Empathy and reactive depression. *Psychol Rep*. 1997;80(3, Pt 1):Jun-1049.

Gaynes BN, Rush AJ, Trivedi MH, Wisniewski SR, Balasubramani GK, Spencer DC, et al. Major depression symptoms in primary care and psychiatric care settings: A cross-sectional analysis. *Ann Fam Med*. 2007 Mar;5(2):126-34.

George L, Elliott SA. Searching for antenatal predictors of postnatal depressive symptomatology: Unexpected findings from a study of obsessive-compulsive personality traits. *J Reprod Infant Psychol*. 2004;22(1):25-31.

Georgiopoulos AM, Bryan TL, Yawn BP, Houston MS, Rummans TA, Therneau TM. Population-based screening for postpartum depression. *Obstetrics and Gynecology*. 1999 May;93(5:Pt 1):653-7.

Gerrity MS, Corson K, Dobscha SK. Screening for posttraumatic stress disorder in VA primary care patients with depression symptoms. *J Gen Intern Med*. 2007;22(9):1321-4.

Ghubash R, El-Rufaie O, Zoubeidi T, Al-Shboul QM, Sabri SM. Profile of mental disorders among the elderly United Arab Emirates population: Sociodemographic correlates. *Int J Geriatr Psychiatry*. 2004 Apr;19(4):344-51.

Giakoumaki O, Vasilaki K, Lili L, Skouroliakou M, Liosis G. The role of maternal anxiety in the early postpartum period: Screening for anxiety and depressive symptomatology in Greece. *J Psychosom Obstet Gynaecol*. 2009 Mar;30(1):21-8.

Girling DM, Barkley C, Paykel ES, Gehlhaar E, Brayne C, Gill C, et al. The prevalence of depression in a cohort of the very elderly. *J Affect Disord*. 1995 Aug 18;34(4):319-29.

Glasser M, Vogels L, Gravdal J. Geriatric depression assessment by rural primary care physicians. *Rural & Remote Health*.9(4):1180-Dec.

Glasser M, Stearns JA, de Kemp E, van Hout J, Hott D. Dementia and depression symptomatology as assessed through screening tests of older patients in an outpatient clinic. *Fam Pract Res J*. 1994 Sep;14(3):261-72.

Glasser M, Gravdal JA. Assessment and treatment of geriatric depression in primary care settings. *Arch Fam Med*. 1997 Sep;6(5):433-8.

Glasser S, Barell V, Shoham A, Ziv A, Boyko V, Lusky A, et al. Prospective study of postpartum depression in an Israeli cohort: Prevalence, incidence and demographic risk factors. *J Psychosom Obstet Gynaecol*. 1998 Sep;19(3):155-64.

Glasser S, Barell V, Boyko V, Ziv A, Lusky A, Shoham A, et al. Postpartum depression in an Israeli cohort: Demographic, psychosocial and medical risk factors. *J Psychosom Obstet Gynaecol*. 2000 Jun;21(2):99-108.

Glavin K, Smith L, Sorum R. Prevalence of postpartum depression in two municipalities in Norway. *Scand J Caring Sci*. 2009 Dec;23(4):705-10.

Godard C, Chevalier A, Siret B, Giorla JF, Hergueta T, Lecrubier Y, et al. A strategy for therapy of depressive disorders and anxiety disorders by a health education intervention in medical consultations: The results of the APRAND program. *Revue d'epidemiologie et de sante publique*. 2007;55(2):113-21.

Goebert D, Morland L, Frattarelli L, Onoye J, Matsu C. Mental health during pregnancy: A study comparing Asian, Caucasian and Native Hawaiian women. *Matern Child Health J*. 2007 May;11(3):249-55.

Goel V, Glazier R, Summers A, Holzapfel S. Psychological outcomes following maternal serum screening: A cohort study. *Can Med Assoc J*. 1998 Sep 22;159(6):651-6.

Goldberg D, Privett M, Ustun B, Simon G, Linden M. The effects of detection and treatment on the outcome of major depression in primary care: A naturalistic study in 15 cities. *Br J Gen Pract*. 1998 Dec;48(437):1840-4.

Goldberg HI, Wagner EH, Fihn SD, Martin DP, Horowitz CR, Christensen DB, et al. A randomized controlled trial of CQI teams and academic detailing: Can they alter compliance with guidelines? *Joint Comm J Qual Improv*. 1998 Mar;24(3):130-42.

Gonzalez JS, Safren SA, Cagliero E, Wexler DJ, Delahanty L, Wittenberg E, et al. Depression, self-care, and medication adherence in type 2 diabetes: Relationships across the full range of symptom severity. *Diabetes Care*. 2007;30(9):2222-7.

Gonzalez JS, Safren SA, Delahanty LM, Cagliero E, Wexler DJ, Meigs JB, et al. Symptoms of depression prospectively predict poorer self-care in patients with Type 2 diabetes. *Diabetic Med*. 2008 Sep;25(9):1102-7.

Goodman JH, Tyer-Viola L. Detection, treatment, and referral of perinatal depression and anxiety by obstetrical providers. *J Womens Health*. 2010;19(3):477-90.

Goodwin RD, Olfson M, Shea S, Lantigua RA, Carrasquillo O, Gameroff MJ, et al. Asthma and mental disorders in primary care. *Gen Hosp Psychiatry*. 2003 Nov;25(6):479-83.

Goodwin RD, Kroenke K, Hoven CW, Spitzer RL. Major depression, physical illness, and suicidal ideation in primary care. *Psychosom Med*. 2003 Jul;65(4):501-5.

Gordon PH, Wang Y, Doorish C, Lewis M, Battista V, Mitsumoto H, et al. A screening assessment of cognitive impairment in patients with ALS. *Amyotrophic Lat Scler*. 2007;8(6):362-5.

Gough K, Hudson P. Psychometric properties of the Hospital Anxiety and Depression Scale in family caregivers of palliative care patients. *J Pain Symptom Manag.* 2009 May;37(5):797-806.

Goyal D, Murphy SO, Cohen J. Immigrant Asian Indian women and postpartum depression. *JOGNN.* 2006 Jan;35(1):98-104.

Grant BF, Stinson FS, Dawson DA, Chou SP, Dufour MC, Compton W, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry.* 2004 Aug;61(8):807-16.

Green JM. Postnatal depression or perinatal dysphoria? Findings from a longitudinal community-based study using the Edinburgh Postnatal Depression Scale. *J Reprod Infant Psychol.* 1998 May;16(2-3):143-55.

Greenberg PE, Kessler RC, Birnbaum HG, Leong SA, Lowe SW, Berglund PA, et al. The economic burden of depression in the United States: How did it change between 1990 and 2000? *J Clin Psychiatr.* 2003 Dec;64(12):1465-75.

Gresenz CR, Sturm R, Tang L. Income and mental health: Unraveling community and individual level relationships. *J Ment Health Pol Econ.* 2001 Dec;4(4):197-203.

Grupp-Phelan J, Whitaker RC, Naish AB. Depression in mothers of children presenting for emergency and primary care: Impact on mothers' perceptions of caring for their children. *Ambul Pediatr.* 2003 May;3(3):142-6.

Grussu P, Quatraro RM. Prevalence and risk factors for a high level of postnatal depression symptomatology in Italian women: A sample drawn from ante-natal classes. *Eur Psychiatry.* 2009 Jun;24(5):327-33.

Gunnell D, Rasul F, Stansfeld SA, Hart CL, Davey SG. Gender differences in self-reported minor mental disorder and its association with suicide. A 20-year follow-up of the Renfrew and Paisley cohort. *Soc Psychiatry Psychiatr Epidemiol.* 2002;37(10):457-9.

Gureje O, Uwakwe R, Oladeji B, Makanjuola VO, Esan O. Depression in adult Nigerians: Results from the Nigerian Survey of Mental Health and Well-being. *J Affect Disord.* 2010 Jan;120(1-3):158-64.

Hadi N, Jamali E, Rahimi F. The prevalence of anxiety and depression in adult hospitalized patients in internal and surgical wards of Shiraz Hospitals-1387(2008). *Shiraz E Med J.* 2010;11(3):137-47.

Haggarty J, Cernovsky Z, Kermeen P, Merskey H. Psychiatric disorders in an Arctic community. *Can J Psychiatry.* 2000;45(4):357-62.

Hajibeigi B, Azarkeyvan A, Alavian SM, Lankarani MM, Assari S. Anxiety and depression affects life and sleep quality in adults with beta-thalassemia. *Indian J Hematol Blood Transfus.* 2009;25(2):59-65.

Hamid F, Asif A, Haider II. Study of anxiety and depression during pregnancy. *Pakistan J Med Sci.* 2008;24(6):861-4.

Hamirani MM, Sultana A, Ibrahim Z, Iqbal H, Sultana S. Frequency of prenatal depression in second and third trimesters of pregnancy in Karachi: A hospital based study. *J Liaquat Uni Med Health Sci.* 2006;5(3):106-9.

Hammond AJ, Yu S, Esa K, Jabbour J, Wakefield L, Ryan P, et al. Factors associated with persistent risk of depression in older people following discharge from an acute cardiac unit. *Int Psychogeriatr.* 2008 Aug;20(4):738-51.

Hankin CS, Spiro A, III, Miller DR, Kazis L. Mental disorders and mental health treatment among U.S. Department of Veterans Affairs outpatients: The Veterans Health Study. *Am J Psychiatry.* 1999 Dec;156(12):1924-30.

Hankin CS, Skinner KM, Sullivan LM, Miller DR, Frayne S, Tripp TJ. Prevalence of depressive and alcohol abuse symptoms among women VA outpatients who report experiencing sexual assault while in the military. *J Traum Stress.* 1999 Oct;12(4):601-12.

Hankin CS, Spiro A, Mansell D, Miller DR, Kazis LE. Mental disorders and medical care utilization of VA ambulatory care patients: The Veterans Health Study. *J Ambul Care Manag.* 2006;29(1):51-60.

Hansen MS, Fink P, Frydenberg M, Oxhøj M-L, Sondergaard L, Munk-Jørgensen P. Mental disorders among internal medical inpatients: Prevalence, detection, and treatment status. *J Psychosom Res.* 2001;50(4):199-204.

Harman JS, Veazie PJ, Lyness JM. Primary care physician office visits for depression by older Americans. *J Gen Intern Med.* 2006 Sep;21(9):926-30.

Harris Y, Cooper JK. Depressive symptoms in older people predict nursing home admission. *J Am Geriatr Soc.* 2006 Apr;54(4):593-7.

Harrison DL, Miller MJ, Schmitt MR, Touchet BK. Variations in the probability of depression screening at community-based physician practice visits. *Prim Care Comp J Clin Psychiatr.* 2010;12(5):e1-e8.

Harrison KS, Rogers R. Axis I screens and suicide risk in jails: A comparative analysis. *Assessment.* 2007;14(2):171-80.

Harvey ST, Pun PK. Analysis of positive Edinburgh depression scale referrals to a consultation liaison psychiatry service in a two-year period. *Int J Ment Health Nurs.* 2007 Jun;16(3):161-7.

Harwood DG, Barker WW, Ownby RL, Mullan MJ, Duara R. Family history of dementia and current depression in nondemented community-dwelling older adults. *J Geriatr Psychiatr Neurol.* 2000;13(2):65-71.

Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: Results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Arch Gen Psychiatry*. 2005 Oct;62(10):1097-106.

Hassett AL, Radvanski DC, Buyske S, Savage SV, Sigal LH. Psychiatric comorbidity and other psychological factors in patients with "chronic Lyme disease". *Am J Med*. 2009;122(9):843-50.

Hatzenbuehler ML, Nolen-Hoeksema S, Erickson SJ. Minority stress predictors of HIV risk behavior, substance use, and depressive symptoms: Results from a prospective study of bereaved gay men. *Health Psychol*. 2008 Jul;27(4):455-62.

Hauser W, Janke K-H, Klump B, Gregor M, Hinz A. Anxiety and depression in adult patients with celiac disease on a gluten-free diet. *World J Gastroenterol*. 2010;16(22):2780-7.

Havranek EP, Ware MG, Lowes BD. Prevalence of depression in congestive heart failure. *Am J Cardiol*. 1999;84(3):348-50.

Havranek EP, Spertus JA, Masoudi FA, Jones PG, Rumsfeld JS. Predictors of the onset of depressive symptoms in patients with heart failure. *J Am Coll Cardiol*. 2004;44(12):2333-8.

Hedayati SS, Minhajuddin AT, Toto RD, Morris DW, Rush AJ. Prevalence of major depressive episode in CKD. *Am J Kidney Dis*. 2009;54(3):424-32.

Heh SS, Fu YY. Effectiveness of informational support in reducing the severity of postnatal depression in Taiwan. *J Adv Nurs*. 2003 Apr;42(1):30-6.

Heikkinen R-L, Kauppinen M. Depressive symptoms in late life: A 10-year follow-up. *Arch Gerontol Geriatr*. 2004;38(3):239-50.

Heisel MJ, Duberstein PR, Lyness JM, Feldman MD. Screening for suicide ideation among older primary care patients. *J Am Board Fam Med*. 2010 Mar;23(2):260-9.

Helvik A-S, Skancke RH, Selbaek G. Screening for depression in elderly medical inpatients from rural area of Norway: Prevalence and associated factors. *Int J Geriatr Psychiatry*. 2010;25(2):150-9.

Hendrie HC, Callahan CM, Levitt EE, Hui SL. Prevalence rates of major depressive disorders: The effects of varying the diagnostic criteria in an older primary care population. *Am J Geriatr Psychiatry*. 1995;3(2):119-31.

Heneghan AM, Silver EJ, Bauman LJ, Stein RE. Do pediatricians recognize mothers with depressive symptoms? *Pediatrics*. 2000 Dec;106(6):1367-73.

Henk HJ, Katzelnick DJ, Kobak KA, Greist JH, Jefferson JW. Medical costs attributed to depression among patients with a history of high medical expenses in a health maintenance organization. *Arch Gen Psychiatry*. 1996 Oct;53(10):899-904.

Henshaw C, Foreman D, Cox J. Postnatal blues: A risk factor for postnatal depression. *J Psychosom Obstet Gynaecol*. 2004 Sep;25(3-4):267-72.

Hensley ML, Robson ME, Kauff ND, Korytowsky B, Castiel M, Ostroff J, et al. Pre- and postmenopausal high-risk women undergoing screening for ovarian cancer: Anxiety, risk perceptions, and quality of life. *Gynecol Oncol*. 2003 Jun;89(3):440-6.

Herrman H, Patrick DL, Diehr P, Martin ML, Fleck M, Simon GE, et al. Longitudinal investigation of depression outcomes in primary care in six countries: The LIDO study. Functional status, health service use and treatment of people with depressive symptoms. *Psychol Med*. 2002 Jul;32(5):889-902.

Herrmann-Lingen C, Klemme H, Meyer T. Depressed mood, physician-rated prognosis, and comorbidity as independent predictors of 1-year mortality in consecutive medical inpatients. *J Psychosom Res*. 2001 Jun;50(6):295-301.

Herrmann C, Brand-Driehorst S, Kaminsky B, Leibing E, Staats H, Ruger U. Diagnostic groups and depressed mood as predictors of 22-month mortality in medical inpatients. *Psychosom Med*. 1998 Sep;60(5):570-7.

Herzig L, Muhlemann N, Bischoff T. Mental disorders in primary care. *Revue Medicale Suisse*. 2010;6(249):1000-5.

Hibino Y, Takaki J, Kambayashi Y, Hitomi Y, Sakai A, Sekizuka N, et al. Relationship between the Noto-Peninsula earthquake and maternal postnatal depression and child-rearing. *Environ Health Prevent Med*. 2009;14(5):255-60.

Hillemacher T, Graassel E, Tigges S, Bleich S, Neundorfer B, Kornhuber J, et al. Depression and bulbar involvement in amyotrophic lateral sclerosis. *Amyotrophic Lat Scler*. 2004;5(4):245-9.

Hillemeier MM, Weisman CS, Chase GA, Dyer AM. Mental health status among rural women of reproductive age: Findings from the Central Pennsylvania Women's Health Study. *Am J Public Health*. 2008 Jul;98(7):1271-9.

Hinton L, Jenkins CN, McPhee S, Wong C, Lai KQ, Le A, et al. A survey of depressive symptoms among Vietnamese-American men in three locales: Prevalence and correlates. *J Nervous Ment Dis*. 1998 Nov;186(11):677-83.

Hinton WL, Tiet Q, Tran CG, Chesney M. Predictors of depression among refugees from Vietnam: A longitudinal study of new arrivals. *J Nervous Ment Dis*. 1997 Jan;185(1):39-45.

Hirschman KB, Shea JA, Xie SX, Karlawish JHT. The development of a rapid screen for caregiver burden. *J Am Geriatr Soc*. 2004;52(10):1724-9.

Ho PM, Masoudi FA, Spertus JA, Peterson PN, Shroyer AL, McCarthy J, et al. Depression predicts mortality following cardiac valve surgery. *Ann Thorac Surg*. 2005;79(4):1255-9.

Holt RI, Phillips DI, Jameson KA, Cooper C, Dennison EM, Peveler RC, et al. The relationship between depression and diabetes mellitus: Findings from the Hertfordshire Cohort Study. *Diabetic Med.* 2009 Jun;26(6):641-8.

Holzman C, Eyster J, Tiedje LB, Roman LA, Seagull E, Rahbar MH. A life course perspective on depressive symptoms in mid-pregnancy. *Matern Child Health J.* 2006 Mar;10(2):127-38.

Homish GG, Cornelius JR, Richardson GA, Day NL. Antenatal risk factors associated with postpartum comorbid alcohol use and depressive symptomatology. *Alcohol Clin Exp Res.* 2004;28(8):1242-8.

Hood KK, Johnson SB, Carmichael SK, Laffel LMB, She J-X, Schatz DA. Depressive symptoms in mothers of infants identified as genetically at risk for type 1 diabetes. *Diabetes Care.* 2005;28(8):1898-903.

Hopwood P, Stephens RJ. Depression in patients with lung cancer: Prevalence and risk factors derived from quality-of-life data. *J Clin Oncol.* 2000;18(4):893-903.

Horan-Smith J, Gullone E. Screening an Australian community sample for risk of postnatal depression. *Aust Psychol.* 1998 Jul;33(2):138-42.

Horowitz A, Reinhardt JP, Kennedy GJ. Major and subthreshold depression among older adults seeking vision rehabilitation services. *Am J Geriatr Psychiatr.* 2005;13(3):180-7.

Hsu GLK, Wan YM, Adler D, Rand W, Choi E, Tsang BYP. Detection of major depressive disorder in Chinese Americans in primary care. *Hong Kong J Psychiatr.* 2005;15(3):71-6.

Huang FY, Chung H, Kroenke K, Delucchi KL, Spitzer RL. Using the Patient Health Questionnaire-9 to measure depression among racially and ethnically diverse primary care patients. *J Gen Intern Med.* 2006;21(6):547-52.

Huang YC, Mathers NJ. Postnatal depression and the experience of South Asian marriage migrant women in Taiwan: Survey and semi-structured interview study. *Int J Nurs Stud.* 2008 Jun;45(6):924-31.

Hunkeler EM, Meresman JF, Hargreaves WA, Fireman B, Berman WH, Kirsch AJ, et al. Efficacy of nurse telehealth care and peer support in augmenting treatment of depression in primary care. *Arch Fam Med.* 2000 Aug;9(8):700-8.

Hunt M, Forand NR. Cognitive vulnerability to depression in never depressed subjects. *Cognit Emot.* 2005;19(5):763-70.

Husain N, Creed F, Tomenson B. Adverse social circumstances and depression in people of Pakistani origin in the UK. *Br J Psychiatry.* 1997 Nov;171:434-8.

Husain N, Creed F, Tomenson B. Depression and social stress in Pakistan. *Psychol Med.* 2000 Mar;30(2):395-402.

Husaini BA, Sherkat DE, Bragg R, Levine R, Emerson JS, Menten CM, et al. Predictors of breast cancer screening in a panel study of African American women. *Women and Health.* 2001;34(3):35-51.

Hybels CF, Pieper CF, Blazer DG. The complex relationship between depressive symptoms and functional limitations in community-dwelling older adults: the impact of subthreshold depression. *Psychol Med*. 2009 Oct;39(10):1677-88.

Iliffe S, Kharicha K, Harari D, Swift C, Gillmann G, Stuck AE. Health risk appraisal in older people 2: The implications for clinicians and commissioners of social isolation risk in older people. *Br J Gen Pract*. 2007 Apr;57(537):277-82.

Imran A, Azidah AK, Asrenee AR, Rosediani M. Prevalence of depression and its associated factors among elderly patients in outpatient clinic of Universiti Sains Malaysia Hospital. *Med J Malaysia*. 2009 Jun;64(2):134-9.

Imran N, Haider II. Screening of antenatal depression in Pakistan: Risk factors and effects on obstetric and neonatal outcomes. *Asia Pacific Psychiatr*. 2010;2(1):26-32.

Ingles J, Lind JM, Phongsavan P, Semsarian C. Psychosocial impact of specialized cardiac genetic clinics for hypertrophic cardiomyopathy. *Genet Med*. 2008;10(2):117-20.

Ingram JC, Greenwood RJ, Woolridge MW. Hormonal predictors of postnatal depression at 6 months in breastfeeding women. *J Reprod Infant Psychol*. 2003;21(1):61-8.

Ip WY, Martin C. The factor structure of the Chinese version of the Edinburgh Postnatal Depression Scale (EPDS). *J Reprod Infant Psychol*. 2007;25(1):83-6.

Isaksson AK, Gunnarsson LG, Ahlstrom G. The presence and meaning of chronic sorrow in patients with multiple sclerosis. *J Clin Nurs*. 2007 Nov;16(11C):315-24.

Jackson-Triche ME, Greer SJ, Wells KB, Rogers W, Camp P, Mazel R. Depression and health-related quality of life in ethnic minorities seeking care in general medical settings. *J Affect Disord*. 2000;58(2):89-97.

Jackson JL, Kroenke K. Prevalence, impact, and prognosis of multisomatoform disorder in primary care: A 5-year follow-up study. *Psychosom Med*. 2008 May;70(4):430-4.

James BO, Omoaregba JO, Eze G, Morakinyo O. Depression among patients with diabetes mellitus in a Nigerian teaching hospital. *S Afr J Psychiatr*. 2010;16(2):61-4.

Jang STC. Clinical depression and severe marital maladjustment in an unemployed sample. *Hong Kong Pract*. 2001;23(1):4-9.

Jaser SS, Whittemore R, Ambrosino JM, Lindemann E, Grey M. Coping and psychosocial adjustment in mothers of young children with type 1 diabetes. *Child Health Care*. 2009 Apr;38(2):91-106.

Jelsma J, Mielke J, Powell G, De Weerd W, De Cock P. Disability in an urban black community in Zimbabwe. *Disabil Rehabil*. 2002 Nov 10;24(16):851-9.

Jenkins R, Lewis G, Bebbington P, Brugha T. The National Psychiatric Morbidity Surveys of Great Britain: Initial findings from the Household Survey. *Psychol Med*. 1997 Jul;27(4):775-89.

Jesse DE, Walcott-McQuigg J, Mariella A, Swanson MS. Risks and protective factors associated with symptoms of depression in low-income African American and Caucasian women during pregnancy. *J Midwifery Womens Health*. 2005 Sep;50(5):405-10.

Jesse DE, Swanson MS. Risks and resources associated with antepartum risk for depression among rural southern women. *Nursing Research*. 2007 Nov;56(6):378-86.

Johansson P, Dahlstrom U, Alehagen U. Depressive symptoms and six-year cardiovascular mortality in elderly patients with and without heart failure. *Scand Cardiovasc J*. 2007 Oct;41(5):299-307.

Johnstone A, Goldberg D. Psychiatric screening in general practice. A controlled trial. *Lancet*. 1976 Mar 20;1(7960):605-8.

Jones GY, Payne S. Searching for safety signals: The experience of medical surveillance amongst men with testicular teratomas. *Psycho Oncol*. 2000;9(5):385-94.

Jones J, Barr W, Robinson J, Carlisle C. Depression in patients with chronic venous ulceration. *Br J Nurs*. 2006;15(11):S17-S23.

Jones LE, Doebbeling CC. Depression screening disparities among veterans with diabetes compared with the general veteran population. *Diabetes Care*. 2007 Sep;30(9):2216-21.

Josefsson A, Berg G, Nordin C, Sydsjo G. Prevalence of depressive symptoms in late pregnancy and postpartum. *Acta Obstet Gynecol Scand*. 2001 Mar;80(3):251-5.

Josefsson A. Postpartum depression--Epidemiological and biological aspects. *Nordic Journal of Psychiatry*. 2003;57(5):395-6.

Josefsson A, Larsson C, Sydsjo G, Nylander PO. Temperament and character in women with postpartum depression. *Arch Womens Ment Health*. 2007;10(1):3-7.

Judd LL, Akiskal HS, Zeller PJ, Paulus M, Leon AC, Maser JD, et al. Psychosocial disability during the long-term course of unipolar major depressive disorder. *Arch Gen Psychiatry*. 2000 Apr;57(4):375-80.

Kadir AA, Nordin R, Ismail SB, Yaacob MJ, Mustapha WMRW. Postnatal depression in mothers attending primary care clinics in Kelantan, Malaysia. *Int Med J*. 2005;12(2):105-9.

Kadir AA, Daud MNM, Yaacob MJ, Hussain NHN. Relationship between obstetric risk factors and postnatal depression in Malaysian women. *Int Med J*. 2009;16(2):101-6.

Kane P, Garber J. The relations among depression in fathers, children's psychopathology, and father-child conflict: A meta-analysis. *Clin Psychol Rev*. 2004 Jul;24(3):339-60.

Kang H, Smith M, Buckwalter KC, Ellingrod V, Schultz SK. Anxiety, depression, and cognitive impairment in dementia-specific and traditional assisted living. *J Gerontol Nurs*. 1992 Mar;36(1):18-30.

Kaplan G, Glasser S, Murad H, Atamna A, Alpert G, Goldbourt U, et al. Depression among Arabs and Jews in Israel: A population-based study. *Soc Psychiatry Psychiatr Epidemiol*. 2010;45(10):931-9.

Karlsson H, Joukamaa M, Lehtinen V. Differences between patients with identified and not identified psychiatric disorders in primary care. *Acta Psychiatrica Scandinavica*. 2000;102(5):354-8.

Karmaliani R, Asad N, Bann CM, Moss N, McClure EM, Pasha O, et al. Prevalence of anxiety, depression and associated factors among pregnant women of Hyderabad, Pakistan. *Int J Soc Psychiatr*. 2009 Sep;55(5):414-24.

Kartha A, Anthony D, Manasseh CS, Greenwald JL, Chetty VK, Burgess J, et al. Depression is a risk factor for rehospitalization in medical inpatients. *Prim Care Comp J Clin Psychiatr*. 2007;9(4):256-62.

Kato K, Sullivan PF, Evengard B, Pedersen NL. Chronic widespread pain and its comorbidities: A population-based study. *Arch Intern Med*. 2006 Aug 14;166(15):1649-54.

Kato K, Sullivan PF, Evengard B, Pedersen NL. A population-based twin study of functional somatic syndromes. *Psychol Med*. 2009;39(3):497-505.

Katon W, Von Korff M, Lin E, Simon G, Walker E, Unutzer J, et al. Stepped collaborative care for primary care patients with persistent symptoms of depression: A randomized trial. *Arch Gen Psychiatry*. 1999 Dec;56(12):1109-15.

Katon W, Russo J, Frank E, Barrett J, Williams JW, Jr., Oxman T, et al. Predictors of nonresponse to treatment in primary care patients with dysthymia. *Gen Hosp Psychiatry*. 2002 Jan;24(1):20-7.

Katon WJ, Lin E, Russo J, Unutzer J. Increased medical costs of a population-based sample of depressed elderly patients. *Arch Gen Psychiatry*. 2003;60(9):897-903.

Katz R, Stephen J, Shaw BF, Matthew A, Newman F, Rosenbluth M. The East York Health Needs Study. I: Prevalence of DSM-III-R psychiatric disorder in a sample of Canadian women. *Br J Psychiatry*. 1995;166(JAN.):100-6.

Kelly RH, Russo J, Katon W. Somatic complaints among pregnant women cared for in obstetrics: Normal pregnancy or depressive and anxiety symptom amplification revisited? *Gen Hosp Psychiatry*. 2001 May;23(3):107-13.

Kemp AH, Hopkinson PJ, Stephan BCM, Clark CR, Gordon E, Bryant RA, et al. Predicting severity of non-clinical depression: Preliminary findings using an integrated approach. *J Integra Neurosci*. 2006;5(1):89-110.

Kemp B, Krause JS, Adkins R. Depression among African Americans, Latinos, and Caucasians with spinal cord injury: An exploratory study. *Rehabil Psychol*. 1999;44(3):235-47.

Keqing L, Ze C, Lijun C, Qinpu J, Guang S, Haoran W, et al. Epidemiological survey of mental disorders in the people aged 18 and older in Hebei Province. *Asian J Psychiatr*. 2008;1(2):51-5.

Kerruish NJ, Campbell-Stokes PL, Gray A, Merriman TR, Robertson SP, Taylor BJ. Maternal psychological reaction to newborn genetic screening for type 1 diabetes. *Pediatrics*. 2007;120(2):e324-e335.

Kessler D, Bennewith O, Lewis G, Sharp D. Detection of depression and anxiety in primary care: Follow up study. *Br Med J*. 2002 Nov 2;325(7371):1016-7.

Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005 Jun;62(6):617-27.

Khademi A, Alleyassin A, Agha-Hosseini M, Ramezanzadeh F, Abhari AA. Pretreatment Beck Depression Inventory score is an important predictor for post-treatment score in infertile patients: A before-after study. *BMC Psychiatry*. 2005;5, 2005. Article Number: 25. Date of Publication: 24 May 2005.

Khatti JB, Nepal MK. Study of depression among geriatric population in Nepal. *NMCJ*. 2006 Dec;8(4):220-3.

Khatwa SA, Abdou MH. Adult depression in Alexandria, Egypt, 1998. *J Egypt Publ Health Assoc*. 1999;74(3-4):333-52.

Kim HG, Mandell M, Crandall C, Kuskowski MA, Dieperink B, Buchberger RL. Antenatal psychiatric illness and adequacy of prenatal care in an ethnically diverse inner-city obstetric population. *Arch Womens Ment Health*. 2006 Mar;9(2):103-7.

Kim M, Han H-R, Phillips L. Metric equivalence assessment in cross-cultural research: Using an example of the center for epidemiological studies - Depression scale. *J Nurs Meas*. 2003;11(1):5-18.

Kirby M, Bruce I, Coakley D, Lawlor BA. Dysthymia among the community-dwelling elderly. *Int J Geriatr Psychiatry*. 1999;14(6):440-5.

Kitamura T, Hirano H, Chen Z, Hirata M. Factor structure of the Zung Self-rating Depression Scale in first-year university students in Japan. *Psychiatry Res*. 2004 Oct 30;128(3):281-7.

Klinkman MS, Coyne JC, Gallo S, Schwenk TL. False positives, false negatives, and the validity of the diagnosis of major depression in primary care. *Arch Fam Med*. 1998 Sep;7(5):451-61.

Kodl MM, Lee JW, Matthews AK, Cummings SA, Olopade OI. Correlates of depressive symptoms among women seeking cancer genetic counseling and risk assessment at a high-risk cancer clinic. *J Genet Couns*. 2006 Aug;15(4):267-76.

Kodl MM, Powell AA, Noorbaloochi S, Grill JP, Bangerter AK, Partin MR. Mental health, frequency of healthcare visits, and colorectal cancer screening. *Medical Care*. 2010 Oct;48(10):934-9.

Koenig HG, Vandermeer J, Chambers A, Burr-Crutchfield L, Johnson JL. Comparison of major and minor depression in older medical inpatients with chronic heart and pulmonary disease. *Psychosomatics*. 2006;47(4):296-303.

Koenig HG. Comparison of older depressed hospitalized patients with and without heart failure/pulmonary disease. *Aging Ment Health*. 2006;10(4):335-42.

Koenig HG. Depression outcome in inpatients with congestive heart failure. *Arch Intern Med*. 2006 May 8;166(9):991-6.

Koenig HG, Vandermeer J, Chambers A, Burr-Crutchfield L, Johnson JL. Minor depression and physical outcome trajectories in heart failure and pulmonary disease. *J Nervous Ment Dis*. 2006;194(3):209-17.

Kok RM, Heeren TJ, Hooijer C, Dinkgreve MA, Rooijmans HG. The prevalence of depression in elderly medical inpatients. *J Affect Disord*. 1995 Feb 21;33(2):77-82.

Kolchakova PY, Akabaliev VH. A study of the effect of age on depressivity in Bulgarian urban population. *Folia Medica*. 2003;45(4):11-5.

Kondo N, Kazama M, Suzuki K, Yamagata Z. Impact of mental health on daily living activities of Japanese elderly. *Prev Med*. 2008 May;46(5):457-62.

Korszun A, Moskvina V, Brewster S, Craddock N, Ferrero F, Gill M, et al. Familiality of symptom dimensions in depression. *Arch Gen Psychiatry*. 2004;61(5):468-74.

Kose N, Ekici G, Karakaya MG, Suvalci S, Demir B, Otman AS. The assessment of the pressure pain threshold and its correlation with depression and anxiety in geriatric nursing home residents with cognitive impairment. *Pain Clin*. 2004;16(2):201-6.

Kowacs F, Socal MP, Ziolkowski SC, Borges-Neto VF, Toniolo DP, Francesconi CR, et al. Symptoms of depression and anxiety, and screening for mental disorders in migrainous patients. *Cephalalgia*. 2003 Mar;23(2):79-89.

Kramer D, Allgaier AK, Fejtikova S, Mergl R, Hegerl U. Depression in nursing homes: Prevalence, recognition, and treatment. *Int J Psychiatr Med*. 2009;39(4):345-58.

Krastel B, Margraf J. Depressive and anxious symptomatology after September 2001: Impact of global and local disasters? *Swiss J Psychol*. 2009 Dec;68(4):221-5.

Kumar A, Clark S, Boudreaux ED, Camargo J. A multicenter study of depression among emergency department patients. *Acad Emerg Med*. 2004;11(12):1284-9.

Kvaal K, McDougall FA, Brayne C, Matthews FE, Dewey ME, MRC CFAS. Co-occurrence of anxiety and depressive disorders in a community sample of older people: Results from the MRC CFAS (Medical Research Council Cognitive Function and Ageing Study). *Int J Geriatr Psychiatry*. 2008 Mar;23(3):229-37.

Lachiewicz A, Dawson D, Spiridigliozzi G, Cuccaro M, Lachiewicz M, McConkie-Rosell A. Indicators of anxiety and depression in women with the fragile X premutation: Assessment of a clinical sample. *J Intellect Disabil Res.* 2010 Jul;54(7):597-610.

LaCoursiere DY, Barrett-Connor E, O'Hara MW, Hutton A, Varner MW. The association between prepregnancy obesity and screening positive for postpartum depression. *BJOG.* 2010 Jul;117(8):1011-8.

Lai DW. Impact of culture on depressive symptoms of elderly Chinese immigrants. *Can J Psychiatry.* 2004 Dec;49(12):820-7.

Lambert MT, Terrell JE, Copeland LA, Ronis DL, Duffy SA. Cigarettes, alcohol, and depression: Characterizing head and neck cancer survivors in two systems of care. *Nicotine and Tobacco Research.* 2005;7(2):233-41.

Lamberty GJ, Bieliauskas LA, Holt CS. Depressive symptom covariation in geriatric clinic patients. *Clin Gerontol.* 1994;15(1):15-27.

Lampic C, Thurfjell E, Bergh J, Sjoden P-O. Short- and long-term anxiety and depression in women recalled after breast cancer screening. *Eur J Cancer.* 2001;37(4):463-9.

Lanzi RG, Pascoe JM, Keltner B, Ramey SL. Correlates of maternal depressive symptoms in a national head start program sample. *Arch Pediatr Adolesc Med.* 1999;153(8):801-7.

Larijani B, Bayat MKS, Gorgani MK, Bandarian F, Akhondzadeh S, Sadjadi SA. Association between depression and diabetes. *Ger J Psychiatr.* 2004;7(4):62-5.

LaRosa AC, Glascoe FP, Macias MM. Parental depressive symptoms: Relationship to child development, parenting, health, and results on parent-reported screening tools. *J Pediatr.* 2009 Jul;155(1):124-8.

Lavretsky H, Mistry R, Bastani R, Gould R, Gokhman I, Huang D, et al. Symptoms of depression and anxiety predict mortality in elderly veterans enrolled in the UPBEAT program. *Int J Geriatr Psychiatry.* 2003;18(2):183-4.

Lawrence JM, Standiford DA, Loots B, Klingensmith GJ, Williams DE, Ruggiero A, et al. Prevalence and correlates of depressed mood among youth with diabetes: The SEARCH for Diabetes in Youth study. *Pediatrics.* 2006;117(4):1348-58.

Le HN, Munoz RF, Soto JA, Delucchi KL, Ippen CG. Identifying risk for onset of major depressive episodes in low-income Latinas during pregnancy and postpartum. *Hispanic J Behav Sci.* 2004 Nov;26(4):463-82.

Lee ATH, Tong MCF, Yuen KCP, Tang PSO, Van Hasselt CA. Hearing impairment and depressive symptoms in an older chinese population. *J Otolaryngol Head Neck Surg.* 2010;39(5):498-503.

Lee D, Yip A, Chiu H, Leung T, Chung T. A psychiatric epidemiological study of postpartum Chinese women. *Am J Psychiatry.* 2001 Feb;158(2):220-6.

Lee DT, Chan SS, Sahota DS, Yip AS, Tsui M, Chung TK. A prevalence study of antenatal depression among Chinese women. *J Affect Disord*. 2004 Oct 1;82(1):93-9.

Lee EJ, Kim JB, Shin IH, Lim KH, Lee SH, Cho GA, et al. Current use of depression rating scales in mental health setting. *Psychiatry Invest*. 2010;7(3):170-6.

Lee IFK, Lee DTF, MacKenzie AE. Correlates of functional limitations in older Chinese patients with chronic obstructive pulmonary disease in Hong Kong. *Heart Lung*. 2006;35(5):324-33.

Lehtinen V, Michalak E, Wilkinson C, Dowrick C, Ayuso-Mateos JL, Dalgard OS, et al. Urban-rural differences in the occurrence of female depressive disorder in Europe--Evidence from the ODIN study. *Soc Psychiatry Psychiatr Epidemiol*. 2003 Jun;38(6):283-9.

Lehtinen V, Sohlman B, Nummelin T, Salomaa M, Ayuso-Mateos JL, Dowrick C. The estimated incidence of depressive disorder and its determinants in the Finnish ODIN sample. *Soc Psychiatry Psychiatr Epidemiol*. 2005 Oct;40(10):778-84.

Lehto S, Koukkunen H, Hintikka J, Viinamaki H, Laakso M, Pyorala K. Depression after coronary heart disease events. *Scand Cardiovasc J*. 2000;34(6):580-3.

Leiferman J. The effect of maternal depressive symptomatology on maternal behaviors associated with child health. *Health Educ Behav*. 2002 Oct;29(5):596-607.

LePage JP, Mogge NL, Garcia-Rea EA. Detecting random responding using the Assessment of Depression Inventory: A brief screening measure of depression. *Depress Anxiety*. 2009;26(6):592-5.

Lepine JP, Gastpar M, Mendlewicz J, Tylee A. Depression in the community: The first pan-European study DEPRES (Depression Research in European Society). *Int Clin Psychopharmacol*. 1997 Jan;12(1):19-29.

Leung WC, Kung F, Lam J, Leung TW, Ho PC. Domestic violence and postnatal depression in a Chinese community. *Int J Gynaecol Obstet*. 2002;79(2):159-66.

Lexis MAS, Jansen NWH, Stevens FCJ, van Amelsvoort LGPM, Kant IJ. Experience of health complaints and help seeking behavior in employees screened for depressive complaints and risk of future sickness absence. *J Occup Rehabil*. 2010 Dec;20(4):537-46.

Li Y-S, Meyer JS, Thornby J. Depressive symptoms among cognitively normal versus cognitively impaired elderly subjects. *Int J Geriatr Psychiatry*. 2001;16(5):455-61.

Licht-Strunk E, van der Kooij KG, van Schaik DJ, van Marwijk HW, van Hout HP, de Haan M, et al. Prevalence of depression in older patients consulting their general practitioner in The Netherlands. *Int J Geriatr Psychiatry*. 2005 Nov;20(11):1013-9.

Lin PC, Wang HH, Huang HT. Depressive symptoms among older residents at nursing homes in Taiwan. *J Clin Nurs*. 2007 Sep;16(9):1719-25.

Linfante AH, Allan R, Smith J, Mosca L. Psychosocial factors predict coronary heart disease, but what predicts psychosocial risk in women. *J Am Med Womens Assoc.* 2003;58(4):248-53.

Lipsky S, Caetano R, Field CA, Bazargan S. The role of alcohol use and depression in intimate partner violence among black and Hispanic patients in an urban emergency department. *Am J Drug Alcohol Abuse.* 2005;31(2):225-42.

Lish JD, Zimmerman M, Farber NJ, Lush D, Kuzma MA, Plescia G. Psychiatric screening in geriatric primary care: Should it be for depression alone?. *J Geriatr Psychiatr Neurol.* 1995 Jul;8(3):141-53.

Liu CY, Wang SJ, Teng EL, Fuh JL, Lin CC, Lin KN, et al. Depressive disorders among older residents in a Chinese rural community. *Psychol Med.* 1997 Jul;27(4):943-9.

Liu SI, Prince M, Blizard B, Mann A. The prevalence of psychiatric morbidity and its associated factors in general health care in Taiwan. *Psychol Med.* 2002 May;32(4):629-37.

Livingston G, Watkin V, Milne B, Manela MV, Katona C. Who becomes depressed? The Islington community study of older people. *J Affect Disord.* 2000 May;58(2):125-33.

Llorca P-M, Courtet P, Martin P, Abbar M, Gay C, Meynard J-A, et al. Formalized recommendations of experts (FRE). Screening and management of bipolar disorder: Methodology. [French]. *Encephale.* 2010;36(SUPPL. 4):S79-S85.

Lobo A, Saz P, Marcos G, Dia JL, De-la-Camara C. The prevalence of dementia and depression in the elderly community in a southern European population. The Zaragoza study. *Arch Gen Psychiatry.* 1995 Jun;52(6):497-506.

Lobo A, Campos R, Marcos G, Garcia-Campayo J, Campayo A, Lopez-Anton R, et al. Somatic and psychiatric co-morbidity in primary care patients in Spain. *Eur J Psychiatry.* 2007 Jan;21(1):71-8.

Lobo A, Saz P, Marcos G, Campos R, Garcia-Campayo J, Orozco F, et al. "Depressive co-morbidity in medical in-patients at the time of hospital discharge and outcome in a primary care follow-up. I. rational and design of the project." *Eur J Psychiatry.* 2005 Jul;19(3):172-92.

Lobo A, Saz P, Marcos G, Dia JL, De la Camara C, Ventura T, et al. The ZARADEMP Project on the incidence, prevalence and risk factors of dementia (and depression) in the elderly community: II. Methods and first results. *Eur J Psychiatry.* 2005 Jan;19(1):40-54.

Logsdon MC, Cross R, Williams B, Simpson T. Prediction of postpartum social support and symptoms of depression in pregnant adolescents: A pilot study. *J Sch Nurs.* 2004 Feb;20(1):36-42.

Logsdon MC, Hutti MH. Readability: An important issue impacting healthcare for women with postpartum depression. *Am J Matern Child Nurs.* 2006 Nov;31(6):350-5.

Lotufo-Neto F. The prevalence of mental disorders among clergy in Sao Paulo, Brazil. *J Psychol Theol.* 1996;24(4):313-22.

Lowe GA, Lipps GE, Young R. Factors associated with depression in students at The University of the West Indies, Mona, Jamaica. *W Indian Med J*. 2009 Jan;58(1):21-7.

Lu CH, Liu CY, Yu S. Depressive disorders among the Chinese elderly in a suburban community. *Public Health Nurs*. 1998 Jun;15(3):196-200.

Lucas A, Pizarro E, Granada ML, Salinas I, Sanmarti A. Postpartum thyroid dysfunction and postpartum depression: Are they two linked disorders? *Clin Endocrinol*. 2001 Dec;55(6):809-14.

Luijendijk HJ, van den Berg JF, Dekker MJ, van Tuijl HR, Otte W, Smit F, et al. Incidence and recurrence of late-life depression. *Arch Gen Psychiatry*. 2008 Dec;65(12):1394-401.

Luke S, Salihu HM, Alio AP, Mbah AK, Jeffers D, Berry EL, et al. Risk factors for major antenatal depression among low-income African American women. *J Womens Health*. 2009 Nov;18(11):1841-6.

Luoma I, Koivisto A-M, Tamminen T. Fathers' and mothers' perceptions of their child and maternal depressive symptoms. *Nordic Journal of Psychiatry*. 2004;58(3):205-11.

Luoma JB, Martin CE, Pearson JL. Contact with mental health and primary care providers before suicide: A review of the evidence. *Am J Psychiatry*. 2002 Jun;159(6):909-16.

Lupsakko T, Mantyjarvi M, Kautiainen H, Sulkava R. Combined hearing and visual impairment and depression in a population aged 75 years and older. *Int J Geriatr Psychiatry*. 2002;17(9):808-13.

Lykeridou K, Daskalakis G, Papadatou D, Vaslamatzis G, Dafni O, Antsaklis A. Dysthymic reactions of women undergoing chorionic villus sampling for prenatal diagnosis of hemoglobinopathies or karyotyping. *Fetal Diagn Ther*. 2004;19(2):149-54.

Lyness JM, Caine ED, King DA, Cox C, Yoediono Z. Psychiatric disorders in older primary care patients. *J Gen Intern Med*. 1999 Apr;14(4):249-54.

Lyness JM, Caine ED, King DA, Conwell Y, Duberstein PR, Cox C. Depressive disorders and symptoms in older primary care patients: One-year outcomes. *Am J Geriatr Psychiatry*. 2002 May;10(3):275-82.

Lyness JM, Yu Q, Tang W, Tu X, Conwell Y. Risks for depression onset in primary care elderly patients: Potential targets for preventive interventions. *Am J Psychiatry*. 2009 Dec;166(12):1375-83.

Maartens LW, Knottnerus JA, Pop VJ. Menopausal transition and increased depressive symptomatology: A community based prospective study. *Maturitas*. 2002 Jul 25;42(3):195-200.

Mackin RS, Ayalon L, Feliciano L, Arean PA. The sensitivity and specificity of cognitive screening instruments to detect cognitive impairment in older adults with severe psychiatric illness. *J Geriatr Psychiatr Neurol*. 2010 Jun;23(2):94-9.

Mallik S, Spertus JA, Reid KJ, Krumholz HM, Rumsfeld JS, Weintraub WS, et al. Depressive symptoms after acute myocardial infarction: Evidence for highest rates in younger women. *Arch Intern Med*. 2006;166(8):876-83.

Maloney DM. Postnatal depression: A study of mothers in the metropolitan area of Perth, Western Australia. *J Aust Coll Midwives*. 1998 Jun;11(2):18-23.

Manning VC, Strathdee G, Best D, Keaney F, McGillivray L, Witton J. Dual diagnosis screening: Preliminary findings on the comparison of 50 clients attending community mental health services and 50 clients attending community substance misuse services. *J Subst Use*. 2002;7(4):221-8.

Manwell LB, Czabala JC, Ignaczak M, Mundt MP. Correlates of depression among heavy drinkers in Polish primary care clinics. *Int J Psychiatr Med*. 2004;34(2):165-78.

Manzoli P, Nunes MA, Schmidt MI, Pinheiro AP, Soares RM, Giacomello A, et al. Violence and depressive symptoms during pregnancy: A primary care study in Brazil. *Soc Psychiatry Psychiatr Epidemiol*. 2010;45(10):983-8.

Marcus DA. Obesity and the impact of chronic pain. *Clin J Pain*. 2004;20(3):186-91.

Marcus SM, Flynn HA, Blow F, Barry K. A screening study of antidepressant treatment rates and mood symptoms in pregnancy. *Arch Womens Ment Health*. 2005 May;8(1):25-7.

Mariam KA, Srinivasan K. Antenatal psychological distress and postnatal depression: A prospective study from an urban clinic. *Asian J Psychiatr*. 2009;2(2):71-3.

Marlow RA, Kegowicz CL, Starkey KN. Prevalence of depression symptoms in outpatients with a complaint of headache. *J Am Board Fam Med*. 2009 Nov;22(6):633-7.

Marshall RD, Olfson M, Hellman F, Blanco C, Guardino M, Struening EL. Comorbidity, impairment, and suicidality in subthreshold PTSD. *Am J Psychiatry*. 2001 Sep;158(9):1467-73.

Martens EJ, Smith OR, Winter J, Denollet J, Pedersen SS. Cardiac history, prior depression and personality predict course of depressive symptoms after myocardial infarction. *Psychol Med*. 2008 Feb;38(2):257-64.

Martucci M, Balestrieri M, Bisoffi G, Bonizzato P, Covre MG, Cunico L, et al. Evaluating psychiatric morbidity in a general hospital: A two-phase epidemiological survey. *Psychol Med*. 1999 Jul;29(4):823-32.

Masmoudi J, Trabelsi S, Charfeddine F, Ben Ayed B, Guerhazi M, Jaoua A, et al. Evaluation of affective temperaments in the postpartum depressive symptomatology. *Encephale*. 2010;36(SUPPL. 2):D14-D21.

Mathers C, Bernard C, Iburg K. Global burden of disease in 2002: Data sources, methods and results. World Health Organization (WHO); 2003. 54.

Mattsson P, Ekselius L. Migraine, major depression, panic disorder, and personality traits in women aged 40-74 years: A population-based study. *Cephalalgia*. 2002;22(7):543-51.

McCabe MP, Davison T, Mellor D, George K, Moore K, Ski C. Depression among older people with cognitive impairment: Prevalence and detection. *Int J Geriatr Psychiatry*. 2006 Jul;21(7):633-44.

McCoy SJ, Beal JM, Payton ME, Stewart AL, DeMers AM, Watson GH. Correlations of visual analog scales with Edinburgh Postnatal Depression Scale. *J Affect Disord*. 2005 Jun;86(2-3):295-7.

McCoy SJ, Beal JM, Payton ME, Stewart AL, DeMers AM, Watson GH. Postpartum thyroid measures and depressive symptomology: A pilot study. *J Am Osteopath Assoc*. 2008 Sep;108(9):503-7.

McCoy SJ, Beal JM, Saunders B, Hill EN, Payton ME, Watson GH. Risk factors for postpartum depression: A retrospective investigation. *J Reprod Med*. 2008 Mar;53(3):166-70.

McCrone S, Cotton S, Jones L, Hawkins TA, Costante J, Nuss M. Depression in a rural, free clinic providing primary care: Prevalence and predictive factors. *Arch Psychiatr Nurs*. 2007 Oct;21(5):291-3.

McCusker J, Karp E, Yaffe MJ, Cole M, Bellavance F. Determining detection of depression in the elderly by primary care physicians: Chart review or questionnaire? *Prim Care Psychiatr*. 1996;2(4):217-21.

McCusker J, Cole M, Dufouil C, Dendukuri N, Latimer E, Windholz S, et al. The prevalence and correlates of major and minor depression in older medical inpatients. *J Am Geriatr Soc*. 2005 Aug;53(8):1344-53.

McCusker J, Cole M, Ciampi A, Latimer E, Windholz S, Belzile E. Major depression in older medical inpatients predicts poor physical and mental health status over 12 months. *Gen Hosp Psychiatry*. 2007 Jul;29(4):340-8.

McDougall FA, Kvaal K, Matthews FE, Paykel E, Jones PB, Dewey ME, et al. Prevalence of depression in older people in England and Wales: The MRC CFA Study. *Psychol Med*. 2007 Dec;37(12):1787-95.

McGee HM, Doyle F, Conroy RM, De La Harpe D, Shelley E. Impact of briefly-assessed depression on secondary prevention outcomes after acute coronary syndrome: A one-year longitudinal survey. *BMC Health Serv Res*. 2006;6:9.

McMillan GP, Timken DS, Lapidus J, C'de Baca J, Lapham SC, McNeal M. Underdiagnosis of comorbid mental illness in repeat DUI offenders mandated to treatment. *J Subst Abuse Treat*. 2008 Apr;34(3):320-5.

McQuade WH, Levy SM, Yanek LR, Davis SW, Liepman MR. Detecting symptoms of alcohol abuse in primary care settings. *Arch Fam Med*. 2000;9(9):814-21.

McSweeney K, O'Connor DW. Depression among newly admitted Australian nursing home residents. *Int Psychogeriatr*. 2008 Aug;20(4):724-37.

Means-Christensen AJ, Sherbourne CD, Roy-Byrne PP, Schulman MC, Wu J, Dugdale DC, et al. In search of mixed anxiety-depressive disorder: A primary care study. *Depress Anxiety*. 2006;23(4):183-9.

Mehnert A, Koch U. Psychological comorbidity and health-related quality of life and its association with awareness, utilization, and need for psychosocial support in a cancer register-based sample of long-term breast cancer survivors. *J Psychosom Res*. 2008 Apr;64(4):383-91.

Melville JL, Delaney K, Newton K, Katon W. Incontinence severity and major depression in incontinent women. *Obstetrics and Gynecology*. 2005 Sep;106(3):585-92.

Menchetti M, Fava C, Berardi D. Disability associated with depressive symptoms in elderly primary care attenders. *Arch Gerontol Geriatr*. 2001;Suppl 7:261-6.

Menchetti M, Cevenini N, De Ronchi D, Quartesan R, Berardi D. Depression and frequent attendance in elderly primary care patients. *Gen Hosp Psychiatry*. 2006 Mar;28(2):119-24.

Menkes DB, Slegg GP, Rowe NF. The frequency of anxiety and depression in purchasers of over-the-counter sleep aids. *Pharmaceut J*. 2006;276(7397):478-9.

Menon AS, Gruber-Baldini AL, Hebel JR, Kaup B, Loreck D, Itkin ZS, et al. Relationship between aggressive behaviors and depression among nursing home residents with dementia. *Int J Geriatr Psychiatry*. 2001;16(2):139-46.

Merckaert I, Libert Y, Messin S, Milani M, Slachmuylder JL, Razavi D. Cancer patients' desire for psychological support: Prevalence and implications for screening patients' psychological needs. *Psycho Oncol*. 2010 Feb;19(2):141-9.

Micali N, Simonoff E, Treasure J. Pregnancy and post-partum depression and anxiety in a longitudinal general population cohort: The effect of eating disorders and past depression. *J Affect Disord*. 2011;131(1-3):150-7.

Michalak EE, Wilkinson C, Hood K, Srinivasan J, Dowrick C, Dunn G, et al. Prevalence and risk factors for depression in a rural setting. Results from the North Wales arm of the ODIN project. *Soc Psychiatry Psychiatr Epidemiol*. 2002 Dec;37(12):567-71.

Michalak EE, Murray G, Wilkinson C, Dowrick C, Lasa L, Lehtinen V, et al. Estimating depression prevalence from the Beck Depression Inventory: Is season of administration a moderator? *Psychiatry Res*. 2004 Nov 30;129(1):99-106.

Mier N, Bocanegra-Alonso A, Zhan D, Wang S, Stoltz SM, Acosta-Gonzalez RI, et al. Clinical depressive symptoms and diabetes in a binational border population. *J Am Board Fam Med*. 2008;21(3):223-33.

Mikolajczyk RT, Maxwell AE, El Ansari W, Naydenova V, Stock C, Ilieva S, et al. Prevalence of depressive symptoms in university students from Germany, Denmark, Poland and Bulgaria. *Soc Psychiatry Psychiatr Epidemiol*. 2008 Feb;43(2):105-12.

Milgrom J, Gemmill AW, Bilszta JL, Hayes B, Barnett B, Brooks J, et al. Antenatal risk factors for postnatal depression: A large prospective study. *J Affect Disord*. 2008 May;108(1-2):147-57.

Miller AM, Sorokin O, Wilbur J, Chandler PJ. Demographic characteristics, menopausal status, and depression in midlife immigrant women. *Womens Health Iss*. 2004 Nov;14(6):227-34.

Miller DK, Malmstrom TK, Joshi S, Andresen EM, Morley JE, Wolinsky FD. Clinically relevant levels of depressive symptoms in community-dwelling middle-aged African Americans. *J Am Geriatr Soc.* 2004 May;52(5):741-8.

Modabernia MJ, Tehrani HS, Fallahi M, Shirazi M, Modabernia AH. Prevalence of depressive disorders in Rasht, Iran: A community based study. *Clin Pract Epidemiol Ment Health.* 2008;4 , 2008. Article Number: 20. Date of Publication: 04 Jul 2008.

Moeller AA, Hell D. Affective disorder and 'psychopathy' in a sample of younger male delinquents. *Acta Psychiatrica Scandinavica.* 2003;107(3):203-7.

Mohan D, Fitzgerald M, Collins C. The relationship between maternal depression (antenatal and pre-school stage) and childhood behavioural problems. *Ir J Psychol Med.* 1998;15(1):10-3.

Molassiotis A, Wilson B, Blair S, Howe T, Cavet J. Unmet supportive care needs, psychological well-being and quality of life in patients living with multiple myeloma and their partners. *Psycho Oncol.* 2011 Jan;20(1):88-97.

Monti F, Agostini F, Marano G, Lupi F. The course of maternal depressive symptomatology during the first 18 months postpartum in an Italian sample. *Arch Womens Ment Health.* 2008 Jul;11(3):231-8.

Mora PA, Bennett IM, Elo IT, Mathew L, Coyne JC, Culhane JF. Distinct trajectories of perinatal depressive symptomatology: Evidence from growth mixture modeling. *Am J Epidemiol.* 2009;169(1):24-32.

Mosca L, Edelman D, Mochari H, Christian AH, Paultre F, Pollin I. Waist circumference predicts cardiometabolic and global Framingham risk among women screened during national woman's heart day. *J Womens Health.* 2006;15(1):24-34.

Mozley CG, Challis D, Sutcliffe C, Bagley H, Burns A, Huxley P, et al. Psychiatric symptomatology in elderly people admitted to nursing and residential homes. *Aging Ment Health.* 2000 May;4(2):136-41.

Muller MA, Bleker OP, Bonsel GJ, Bilardo CM. Nuchal translucency screening and anxiety levels in pregnancy and puerperium. *Ultrasound Obstetr Gynecol.* 2006;27(4):357-61.

Muluk NB, Oguzturk O, Ekici A, Koc C. Emotional effects of nasal packing measured by the hospital anxiety and depression scale in patients following nasal surgery. *J Otolaryngol.* 2005;34(3):172-7.

Mumford DB, Saeed K, Ahmad I, Latif S, Mubbashar MH. Stress and psychiatric disorder in rural Punjab: A community survey. *Br J Psychiatry.* 1997;170:473-8.

Mumford DB, Minhas FA, Akhtar I, Akhter S, Mubbashar MH. Stress and psychiatric disorder in urban Rawalpindi. Community survey. *Br J Psychiatry.* 2000;177:557-62.

Muneer A, Minhas FA, Tamiz-ud-Din NA, Mujeeb F, Usmani AT. Frequency and associated factors for postnatal depression. *J Coll Phys Surg Pakistan.* 2009 Apr;19(4):236-9.

Mykletun A, Overland S, Aaro LE, Liabo HM, Stewart R. Smoking in relation to anxiety and depression: Evidence from a large population survey: The HUNT study. *Eur Psychiatry*. 2008 Mar;23(2):77-84.

Nagy E, Molnar P, Pal A, Orvos H. Prevalence rates and socioeconomic characteristics of post-partum depression in Hungary. *Psychiatry Res*. 2011;185(1-2):113-20.

Naqvi TZ, Rafique AM, Andreas V, Rahban M, Mirocha J, Naqvi SS. Predictors of depressive symptoms post-acute coronary syndrome. *Gender Med*. 2007 Dec;4(4):339-51.

Narrow WE, Rae DS, Robins LN, Regier DA. Revised prevalence estimates of mental disorders in the United States: Using a clinical significance criterion to reconcile 2 surveys' estimates. *Arch Gen Psychiatry*. 2002 Feb;59(2):115-23.

Nasser J, Habib F, Hasan M, Khalil N. Prevalence of depression among people with diabetes attending diabetes clinics at primary health settings. *Bahrain Med Bull*. 2009;31(3).

Navarro P, Esteve L, Ascaso C, Aguado J, Gelabert E, Santos R. Non-psychotic psychiatric disorders after childbirth: Prevalence and comorbidity in a community sample. *J Affect Disord*. 2008 Jul;109(1-2):171-6.

Nazemi H, Larkin AA, Sullivan MD, Katon W. Methodological issues in the recruitment of primary care patients with depression. *Int J Psychiatr Med*. 2001;31(3):277-88.

Nelson KR, Bui H, Samet JH. Screening in special populations: A "case study" of recent Vietnamese immigrants. *Am J Med*. 1997 May;102(5):435-40.

Newmann JP, Klein MH, Jensen JE, Essex MJ. Depressive symptom experiences among older women: A comparison of alternative measurement approaches. *Psychol Aging*. 1996 Mar;11(1):112-26.

Ng A, Earnest A, Yeo D, Kandiah N. Cognitive screening in Asia: Recognizing the role of the patient-family unit. *Dement Geriatr Cognit Disord*. 2010;29(4):351-5.

Niaz S, Izhar N, Bhatti MR. Anxiety and depression in pregnant women presenting in the OPD of a teaching hospital. *Pakistan J Med Sci*. 2004;20(2):117-9.

Nicholson RA, Gramling SE, Ong JC, Buenevar L. Differences in anger expression between individuals with and without headache after controlling for depression and anxiety. *Headache*. 2003;43(6):651-63.

Nidhinandana S, Chinvarun Y, Sithinamsuwan P, Udommongkol C, Suwantamee J, Wongmek W, et al. Prevalence of depression among epileptic patients at Phramongkutklo Hospital. *J Med Assoc Thailand*. 2007 Jan;90(1):32-6.

Noltorp S, Gottfries CG, Norgaard N. Simple steps to diagnosis at primary care centres. *Int Clin Psychopharmacol*. 1998 Sep;13:Suppl-4.

Nordstrom A, Bodlund O. Every third patient in primary care suffers from depression, anxiety or alcohol problems. *Nordic Journal of Psychiatry*. 2008;62(3):250-5.

Norholm V, Bech P. Quality of life in schizophrenic patients: Association with depressive symptoms. *Nordic Journal of Psychiatry*. 2006;60(1):32-7.

Norton J, de Roquefeuil G., David M, Boulenger JP, Ritchie K, Mann A. Prevalence of psychiatric disorders in French general practice using the patient health questionnaire: Comparison with GP case-recognition and psychotropic medication prescription. *Encephale*. 2009 Dec;35(6):560-9.

Norton MC, Skoog I, Toone L, Corcoran C, Tschanz JT, Lisota RD, et al. Three-year incidence of first-onset depressive syndrome in a population sample of older adults: The Cache County study. *Am J Geriatr Psychiatry*. 2006 Mar;14(3):237-45.

Nourissat A, Mille D, Delaroche G, Jacquin JP, Vergnon JM, Fournel P, et al. Estimation of the risk for nutritional state degradation in patients with cancer: Development of a screening tool based on results from a cross-sectional survey. *Ann Oncol*. 2007;18(11):1882-6.

Nugent WR. A psychometric study of the Multi-Problem Screening Inventory depression subscale using item response and generalizability theories. *Res Soc Work Pract*. 2003 Jan;13(1):65-79.

O'Boyle AL, Magann EF, Ricks RE, Jr., Doyle M, Morrison JC. Depression screening in the pregnant soldier wellness program. *Southern Med J*. 2005 Apr;98(4):416-8.

O'Malley PG, Jackson JL, Kroenke K, Yoon K, Hornstein E, Dennis GJ. The value of screening for psychiatric disorders in rheumatology referrals. *Arch Intern Med*. 1998 Nov 23;158(21):2357-62.

O'Rourke N. Research note: Equivalence of French and English language versions of the Center for Epidemiologic Studies-Depression Scale (CES-D) among caregivers of persons with dementia. *Can J Aging*. 2003;22(3):323-9.

Oiji A, Muraoka Y, Nadaoka T, Kashiwakura M, Totsuka S, Ihara K, et al. One-year prognosis of depressive illness in the elderly population in Japan. *Psychiatry Clin Neurosci*. 1998 Aug;52(4):391-5.

Okonkwo OC, Sui X, Ahmed A. Disease-specific depression and outcomes in chronic heart failure: A propensity score analysis. *Compr Ther*. 2007;33(2):65-70.

Okwumabua JO, Baker FM, Wong SP, Pilgram BO. Characteristics of depressive symptoms in elderly urban and rural African Americans. *J Gerontol Biol Med Sci*. 1997 Jul;52(4):M241-M246.

Olivera J, Benabarre S, Lorente T, Rodriguez M, Pelegrin C, Calvo JM, et al. Prevalence of psychiatric symptoms and mental disorders detected in primary care in an elderly Spanish population. The PSICOTARD Study: Preliminary findings. *Int J Geriatr Psychiatry*. 2008;23(9):915-21.

Olson AL, DiBrigida LA. Depressive symptoms and work role satisfaction in mothers of toddlers. *Pediatrics*. 1994 Sep;94(3):363-7.

Onozawa K, Kumar RC, Adams D, Dore C, Glover V. High EPDS scores in women from ethnic minorities living in London. *Arch Womens Ment Health*. 2003 Aug;6 Suppl 2:S51-5.

Osborn DP, Fletcher AE, Smeeth L, Stirling S, Nunes M, Breeze E, et al. Geriatric Depression Scale scores in a representative sample of 14 545 people aged 75 and over in the United Kingdom: Results from the MRC Trial of Assessment and Management of Older People in the Community. *Int J Geriatr Psychiatry*. 2002 Apr;17(4):375-82.

Ostler K, Thompson C, Kinmonth A-LK, Peveler RC, Stevens L, Stevens A. Influence of socio-economic deprivation on the prevalence and outcome of depression in primary care: The Hampshire Depression Project. *Br J Psychiatry*. 2001;178:12-7.

Otsuka K, Yamanaka G, Shinagawa M, Murakami S, Yamanaka T, Shibata K, et al. Chronomic community screening reveals about 31% depression, elevated blood pressure and infradian vascular rhythm alteration. *Biomed Pharmacother*. 2004 Oct;58:Suppl-55.

Owen RR, Cannon D, Thrush CR, Quality ER, I. Mental Health QUERI Initiative: Expert ratings of criteria to assess performance for major depressive disorder and schizophrenia. *Am J Med Qual*. 2003 Jan;18(1):15-20.

Oyane NMF, Bjelland I, Pallesen S, Holsten F, Bjorvatn B. Seasonality is associated with anxiety and depression: The Hordaland health study. *J Affect Disord*. 2008 Jan;105(1-3):147-55.

Pajulo M, Savonlahti E, Sourander A, Helenius H, Piha J. Antenatal depression, substance dependency and social support. *J Affect Disord*. 2001 Jun;65(1):9-17.

Pando Moreno M, Aranda Beltran C, Ponce De Leon Barbosa JL, Salazar Estrada JG, Aguilar Aldrete ME. Mental health in senior citizens in the metropolitan zone of Guadalajara. *Aging Clin Exp Res*. 1999;11(2):96-100.

Panzarine S, Slater E, Sharps P. Coping, social support, and depressive symptoms in adolescent mothers. *J Adolesc Health*. 1995 Aug;17(2):113-9.

Paquay L, De LJ, Schoenmakres B, Stessens J, Bouwen A, van der Burg M, et al. The Qualidem project in Belgium. A two-center study on care needs and provision in dementia care: Inclusion criteria and description of the population. *Arch Public Health*. 2004;62(3-4):125-42.

Patch C, Roderick P, Rosenberg W. Comparison of genotypic and phenotypic strategies for population screening in hemochromatosis: Assessment of anxiety, depression, and perception of health. *Genet Med*. 2005 Oct;7(8):550-6.

Patel V, Prince M. Maternal psychological morbidity and low birth weight in India. *Br J Psychiatry*. 2006;188(MAR.):284-5.

Patten SB. A major depression prognosis calculator based on episode duration. *Clin Pract Epidemiol Ment Health*. 2006;2:13.

Patten SB, Beck CA, Kassam A, Williams JVA, Barbui C, Metz LM. Long-term medical conditions and major depression: Strength of association for specific conditions in the general population. *Can J Psychiatry*. 2005 Mar;50(4):195-202.

Pearson SD, Katzelnick DJ, Simon GE, Manning WG, Helstad CP, Henk HJ. Depression among high utilizers of medical care. *J Gen Intern Med*. 1999 Aug;14(8):461-8.

Pedersen SS, van Domburg RT, Theuns DAMJ, Jordaens L, Erdman RAM. Concerns about the implantable cardioverter defibrillator: A determinant of anxiety and depressive symptoms independent of experienced shocks. *Am Heart J*. 2005;149(4):664-9.

Peindl KS, Wisner KL. Successful recruitment strategies for women in postpartum mental health trials. *J Psychiatr Res*. 2003;37(2):117-25.

Penninx BW, Guralnik JM, Ferrucci L, Simonsick EM, Deeg DJ, Wallace RB. Depressive symptoms and physical decline in community-dwelling older persons. *JAMA*. 1998 Jun 3;279(21):1720-6.

Penninx BW, Leveille S, Ferrucci L, van Eijk JT, Guralnik JM. Exploring the effect of depression on physical disability: Longitudinal evidence from the established populations for epidemiologic studies of the elderly. *Am J Public Health*. 1999 Sep;89(9):1346-52.

Penninx BWJH, Beekman ATF, Johannes HS, Zitman FG, Nolen WA, Spinhoven P, et al. The Netherlands Study of Depression and Anxiety (NESDA): Rationale, objectives and methods. *Int J Meth Psychiatr Res*. 2008;17(3):121-40.

Perez GH, Nicolau JC, Romano BW, Laranjeira R. Depression and acute coronary syndromes: Gender-related differences. *Arquivos Brasileiros de Cardiologia*. 2005;85(5).

Perrino T, Brown SC, Mason CA, Szapocznik J. Depressive symptoms among urban Hispanic older adults in Miami: Prevalence and sociodemographic correlates. *Clin Gerontol*. 2009 Jan;32(1):26-43.

Peterlini M, Tiberio IFLC, Saadeh A, Pereira JCR, Martins MA. Anxiety and depression in the first year of medical residency training. *Med Educ*. 2002;36(1):66-72.

Peters R, Pinto E, Beckett N, Swift C, Potter J, McCormack T, et al. Association of depression with subsequent mortality, cardiovascular morbidity and incident dementia in people aged 80 and over and suffering from hypertension. Data from the Hypertension in the Very Elderly Trial (HYVET). *Age Ageing*. 2010 Jul;39(4):439-45.

Peveler R, George C, Kinmonth AL, Campbell M, Thompson C. Effect of antidepressant drug counselling and information leaflets on adherence to drug treatment in primary care: Randomised controlled trial. *Br Med J*. 1999 Sep 4;319(7210):612-5.

Pfaff JJ, Almeida OP. A cross-sectional analysis of factors that influence the detection of depression in older primary care patients. *Aust NZ J Psychiatry*. 2005 Apr;39(4):262-5.

- Phillips C, Rumsey N. Considerations for the provision of psychosocial services for families following paediatric burn injury-A quantitative study. *Burns*. 2008;34(1):56-62.
- Pincus HA, Tanielian TL, Marcus SC, Olfson M, Zarin DA, Thompson J, et al. Prescribing trends in psychotropic medications: Primary care, psychiatry, and other medical specialties. *JAMA*. 1998 Feb 18;279(7):526-31.
- Pincus T, Griffith J, Pearce S, Isenberg D. Prevalence of self-reported depression in patients with rheumatoid arthritis. *Br J Rheumatol*. 1996 Sep;35(9):879-83.
- Pinheiro RT, da Silva RA, Magalhaes PV, Horta BL, Pinheiro KA. Two studies on suicidality in the postpartum. *Acta Psychiatrica Scandinavica*. 2008 Aug;118(2):160-3.
- Pini S, Perkonig A, Tansella M, Wittchen HU. Prevalence and 12-month outcome of threshold and subthreshold mental disorders in primary care. *J Affect Disord*. 1999 Nov;56(1):37-48.
- Pirraglia PA, Stafford RS, Singer DE. Trends in prescribing of selective serotonin reuptake inhibitors and other newer antidepressant agents in adult primary care. *Prim Care Comp J Clin Psychiatr*. 2003 Aug;5(4):153-7.
- Pitkälä K, Kähönen-Väre M, Valvanne J, Strandberg TE, Tilvis RS. Long-term changes in mood of an aged population: Repeated Zung-tests during a 10-year follow-up. *Arch Gerontol Geriatr*. 2003 Mar;36(2):185-95.
- Podolska MZ, Bidzan M, Majkowicz M, Podolski J, Sipak-Szmigiel O, Ronin-Walknowska E. Personality traits assessed by the NEO Five-Factor Inventory (NEO-FFI) as part of the perinatal depression screening program. *Med Sci Mon*. 2010 Aug 7;16(9):H77-H81.
- Poleshuck EL, Bair MJ, Kroenke K, Watts A, Tu X, Giles DE. Pain and depression in gynecology patients. *Psychosomatics*. 2009 May;50(3):270-6.
- Pollack CE, Lynch J. Health status of people undergoing foreclosure in the Philadelphia region. *Am J Public Health*. 2009 Oct;99(10):1833-9.
- Pompili M, Girardi P, Tatarelli R, Lester D. Subthreshold bipolar traits and suicide risk among undergraduates. *Psychol Rep*. 2006;98(2):417-8.
- Pompili M, Vanacore N, Maccone S, Amore M, Perticoni G, Tonna M, et al. Depression, hopelessness and suicide risk among patients suffering from epilepsy. *Annali Dell'Istituto Superiore di Sanita*. 2007;43(4):425-9.
- Pope H, Watkins KW, Evans AE, Hess P. The perception of depression in long-term-care residents: A qualitative study using residential journaling. *J Appl Gerontol*. 2006;25(2):153-72.
- Post D, Miller K. Use of the health history as a psychiatric screening tool. *J Am Board Fam Pract*. 1998 Nov;11(6):452-8.

Pot AM, Deeg DJH, Twisk JWR, Beekman ATF, Zarit SH. The longitudinal relationship between the use of long-term care and depressive symptoms in older adults. *Gerontologist*. 2005 Jun;45(3):359-69.

Pottinger AM, Trotman-Edwards H, Younger N. Detecting depression during pregnancy and associated lifestyle practices and concerns among women in a hospital-based obstetric clinic in Jamaica. *Gen Hosp Psychiatry*. 2009 May;31(3):254-61.

Poutanen O, Koivisto A-M, Salokangas RKR. Applicability of the DEPS Depression Scale: Assessing format and individual items in subgroups of patients. *Nordic Journal of Psychiatry*. 2010;64(6):384-90.

Powers JR, Young AF, Russell A, Pachana NA. Implications of non-response of older women to a short form of the Center for Epidemiologic Studies Depression Scale. *Int J Aging Human Dev*. 2003;57(1):37-54.

Probst JC, Laditka SB, Moore CG, Harun N, Powell MP, Baxley EG. Rural-urban differences in depression prevalence: Implications for family medicine. *Fam Med*. 2006;38(9):653-60.

Prodromidis M, Abrams S, Field T, Scafidi F. Psychosocial stressors among depressed adolescent mothers. *Adolesc*. 1994;29(114):331-43.

Qiu C, Sanchez SE, Lam N, Garcia P, Williams MA. Associations of depression and depressive symptoms with preeclampsia: Results from a Peruvian case-control study. *BMC Womens Health*. 2007;7:15.

Rajkumar AP, Thangadurai P, Senthilkumar P, Gayathri K, Prince M, Jacob KS. Nature, prevalence and factors associated with depression among the elderly in a rural south Indian community. *Int Psychogeriatr*. 2009 Apr;21(2):372-8.

Ramos-Marcuse F, Oberlander SE, Papas MA, McNary SW, Hurley KM, Black MM. Stability of maternal depressive symptoms among urban, low-income, African American adolescent mothers. *J Affect Disord*. 2010 Apr;122(1-2):68-75.

Rauch SAM, Morales KH, Zubritsky C, Knott K, Oslin D. Posttraumatic stress, depression, and health among older adults in primary care. *Am J Geriatr Psychiatry*. 2006 Apr;14(4):316-24.

Reed BD, Haefner HK, Punch MR, Roth RS, Gorenflo DW, Gillespie BW. Psychosocial and sexual functioning in women with vulvodynia and chronic pelvic pain: A comparative evaluation. *J Reprod Med Obstet Gynecol*. 2000;45(8):624-32.

Reger MA, Gahm GA, Swanson RD, Duma SJ. Association between number of deployments to Iraq and mental health screening outcomes in US Army soldiers. *J Clin Psychiatry*. 2009;70(9):1266-72.

Reighard FT, Evans ML. Use of the Edinburgh Postnatal Depression Scale in a southern, rural population in the United States. *Prog Neuropsychopharmacol Biol Psychiatry*. 1995 Nov;19(7):1219-24.

Rice DP, Miller LS. The economic burden of affective disorders. *Br J Psychiatry*. 1995 Apr;(27):34-42.

Ried LD, Planas LG. Aging, health, and depressive symptoms: Are women and men different? *J Womens Health*. 2002 Nov;11(9):813-24.

Riekert KA, Bartlett SJ, Boyle MP, Krishnan JA, Rand CS. The association between depression, lung function, and health-related quality of life among adults with cystic fibrosis. *Chest*. 2007 Jul;132(1):231-7.

Righetti-Veltema M, Conne-Perreard E, Bousquet A, Manzano J. Construction and multicentric validation of an antepartum screening questionnaire for postpartum depression. *La Psychiatrie de l'Enfant*. 2006;49(2):513-41.

Riolo SA, Nguyen TA, Greden JF, King CA. Prevalence of depression by race/ethnicity: Findings from the National Health and Nutrition Examination Survey III. *Am J Public Health*. 2005 Jun;95(6):998-1000.

Roca-Bennasar M, Gili-Planas M, Ferrer-Perez V, Bernardo-Arroyo M. Mental disorders and medical conditions: A community study in a small island in Spain. *J Psychosom Res*. 2001;50(1):39-44.

Rodriguez CA, Schonfeld L, King-Kallimanis B, Gum AM. Depressive symptoms and alcohol abuse/misuse in older adults: Results from the Florida BRITE Project. *Best Pract Ment Health*. 2010 Jan;6(1):90-102.

Rogers G, Curry M, Oddy J, Pratt N, Beilby J, Wilkinson D. Depressive disorders and unprotected casual anal sex among Australian homosexually active men in primary care. *HIV Med*. 2003 Jul;4(3):271-5.

Ronzio CR, Mitchell SJ. The highs and lows of maternal depression: Cluster analysis of depression symptoms in a sample of African American women. *J Investig Med*. 2010;58(7):887-92.

Rovner BW, Casten RJ, Tasman WS. Effect of depression on vision function in age-related macular degeneration. *Arch Ophthalmol*. 2002 Aug;120(8):1041-4.

Rowe HJ, Fisher JRW, Loh WM. The Edinburgh Postnatal Depression Scale detects but does not distinguish anxiety disorders from depression in mothers of infants. *Arch Womens Ment Health*. 2008 Jun;11(2):103-8.

Roy-Byrne P, Katon W, Broadhead WE, Lepine JP, Richards J, Brantley PJ, et al. Subsyndromal ("mixed") anxiety--Depression in primary care. *J Gen Intern Med*. 1994 Sep;9(9):507-12.

Rubenstein LV, Rayburn NR, Keeler EB, Ford DE, Rost KM, Sherbourne CD. Predicting outcomes of primary care patients with major depression: development of a depression prognosis index. *Psychiatr Serv*. 2007 Aug;58(8):1049-56.

Rubertsson C, Wickberg B, Gustavsson P, Radestad I. Depressive symptoms in early pregnancy, two months and one year postpartum-prevalence and psychosocial risk factors in a national Swedish sample. *Arch Womens Ment Health*. 2005 Jun;8(2):97-104.

Rumbold AR, Crowther CA. Women's experiences of being screened for gestational diabetes mellitus. *Aust NZ J Obstet Gynaecol*. 2002 May;42(2):131-7.

Ruoff GE. Depression and migraine: To what extent is migraine a marker for increased prevalence of selected comorbidities? *Headache Pain*. 2006;17(2):68-72.

Saeed K, Ijaz-Ur-Rehman, Mubbashar MH. Prevalence of psychiatric morbidity among the attendees of a native healer at Rawalpindi. *J Coll Phys Surg Pakistan*. 2000;10(1):7-9.

Sagrestano LM, Rodriguez AC, Carroll D, Bieniarz A, Greenberg A, Castro L, et al. A comparison of standardized measures of psychosocial variables with single-item screening measures used in an urban obstetric clinic. *JOGNN*. 2002;31(2):147-55.

Sakado K, Kuwabara H, Sato T, Uehara T, Sakado M, Someya T. The relationship between personality, dysfunctional parenting in childhood, and lifetime depression in a sample of employed Japanese adults. *J Affect Disord*. 2000;60(1):47-51.

Samuels SC, Katz IR, Parmelee PA, Boyce AA, DiFilippo S. Use of the Hamilton and Montgomery-Asberg depression scales in institutionalized elderly patients. *Am J Geriatr Psychiatry*. 1996;4(3):237-46.

Sandanger I, Nygard JF, Ingebrigtsen G, Sorensen T, Dalgard OS. Prevalence, incidence and age at onset of psychiatric disorders in Norway. *Soc Psychiatry Psychiatr Epidemiol*. 1999 Nov;34(11):570-9.

Sanne B, Mykletun A, Dahl AA, Moen BE, Tell GS. Occupational differences in levels of anxiety and depression: The Hordaland Health Study. *J Occup Environ Med*. 2003;45(6):628-38.

Sanne B, Mykletun A, Moen BE, Dahl AA, Tell GS. Farmers are at risk for anxiety and depression: The Hordaland Health Study. *Occup Med*. 2004;54(2):92-100.

Santos IS, Matijasevich A, Domingues MR, Barros AJD, Barros FCF. Long-lasting maternal depression and child growth at 4 years of age: A cohort study. *J Pediatr*. 2010;157(3):401-6.

Sato Y, Tanda H, Kato S, Onishi S, Nakajima H, Nanbu A, et al. Prevalence of major depressive disorder in self-referred patients in a late onset hypogonadism clinic. *Int J Impot Res*. 2007 Jul;19(4):407-10.

Saus E, Soria V, Escaramis G, Crespo JM, Valero J, Gutierrez-Zotes A, et al. A haplotype of glycogen synthase kinase 3beta is associated with early onset of unipolar major depression. *Genes Brain Behav*. 2010 Oct;9(7):799-807.

Savin-Williams RC, Cohen KM, Joyner K, Rieger G. Depressive symptoms among same-sex oriented young men: Importance of reference group. *Arch Sex Behav*. 2010 Dec;39(6):1213-5.

Saz P, Launer LJ, Dia JL, De la Camara C, Marcos G, Lobo A. Mortality and mental disorders in a Spanish elderly population. *Int J Geriatr Psychiatry*. 1999 Dec;14(12):1031-8.

Scarinci IC, Beech BM, Naumann W, Kovach KW, Pugh L, Fapohunda B. Depression, socioeconomic status, age, and marital status in black women: a national study. *Ethn Dis*. 2002;12(3):421-8.

Schol-Gelok S, Janssens AC, Tiemeier H, Liu F, Lopez-Leon S, Zorkoltseva IV, et al. A genome-wide screen for depression in two independent Dutch populations. *Biol Psychiatry*. 2010 Jul 15;68(2):187-96.

Schroeder V, Borner U, Gutknecht S, Schmid J-P, Saner H, Kohler HP. Relation of depression to various markers of coagulation and fibrinolysis in patients with and without coronary artery disease. *Eur J Cardiovasc Prev Rehabil.* 2007;14(6):782-7.

Schulberg HC, Mulsant B, Schulz R, Rollman BL, Houck PR, Reynolds CFI. Characteristics and course of major depression in older primary care patients. *Int J Psychiatr Med.* 1998;28(4):421-36.

Segre LS, O'Hara MW, Arndt S, Stuart S. The prevalence of postpartum depression: The relative significance of three social status indices. *Soc Psychiatry Psychiatr Epidemiol.* 2007 Apr;42(4):316-21.

Serber ER, Todaro JF, Tilkemeier PL, Niaura R. Prevalence and characteristics of multiple psychiatric disorders in cardiac rehabilitation patients. *J Cardiopulm Rehabil Prev.* 2009;29(3):161-8.

Shah A, Hoxey K. Depression in acutely medically ill elderly inpatients: Prevalence, correlates and longitudinal stability. *Int J Meth Psychiatr Res.* 2001;10(3):147-56.

Shah BM, Gupchup GV, Borrego ME, Raisch DW, Knapp KK. Depressive symptoms in patients with type 2 diabetes in the ambulatory care setting: Opportunities to improve outcomes in the course of routine care. *JAPhA.* 2008;48(6):737-43.

Shah DC, Evans M, King D. Prevalence of mental illness in a rehabilitation unit for older adults. *Postgrad Med J.* 2000 Mar;76(893):153-6.

Shah R, Uren Z, Baker A, Majeed A. Deaths from antidepressants in England and Wales 1993-1997: Analysis of a new national database. *Psychol Med.* 2001 Oct;31(7):1203-10.

Shakespeare J, Blake F, Garcia J. Breast-feeding difficulties experienced by women taking part in a qualitative interview study of postnatal depression. *Midwifery.* 2004;20(3):251-60.

Sharkey JR, Haines PS. Use of telephone-administered survey for identifying nutritional risk indicators among community-living older adults in rural areas. *J Appl Gerontol.* 2002;21(3):385-403.

Sharma P, Avasthi A, Chakrabarti S, Varma S. Depression among hospitalised medically ill patients: A two-stage screening study. *J Affect Disord.* 2002 Jul;70(2):205-9.

Shaw WS, Means-Christensen AJ, Slater MA, Webster JS, Patterson TL, Grant I, et al. Psychiatric disorders and risk of transition to chronicity in men with first onset low back pain. *Pain Med.* 2010 Sep;11(9):1391-400.

Sheppard K, Badger T. The lived experience of depression among culturally deaf adults. *J Psychiatr Ment Health Nurs.* 2010 Nov;17(9):783-9.

Sherina MS, Rampal L, Mustaqim A. The prevalence of depression among the elderly in Sepang, Selangor. *Med J Malaysia.* 2004 Mar;59(1):45-9.

Sherina MS, Rampal L, Hanim MA, Thong PL. The prevalence of depression among elderly warded in a tertiary care centre in Wilayah Persekutuan. *Med J Malaysia.* 2006 Mar;61(1):15-21.

Shiels C, Gabbay M, Exley D. Psychological distress in students registered at a university-based general practice. *Prim Care Community Psychiatr.* 2008;13(1):9-18.

Shoukri MM, Donner A. Bivariate modeling of interobserver agreement coefficients. *Stat Med.* 2009;28(3):430-40.

Shumway M, Sentell T, Unick G, Bamberg W. Cognitive complexity of self-administered depression measures. *J Affect Disord.* 2004;83(2-3):191-8.

Shvartzman P, Weiner Z, Vardy D, Friger M, Sherf M, Biderman A. Health services utilization by depressive patients identified by the MINI questionnaire in a primary care setting. *Scand J Prim Health Care.* 2005 Mar;23(1):18-25.

Siefert K, Heflin CM, Corcoran ME, Williams DR. Food insufficiency and physical and mental health in a longitudinal survey of welfare recipients. *J Health Soc Behav.* 2004;45(2):171-86.

Silove D, Bateman CR, Brooks RT, Fonseca CAZ, Steel Z, Rodger J, et al. Estimating clinically relevant mental disorders in a rural and an urban setting in postconflict Timor Leste. *Arch Gen Psychiatry.* 2008 Oct;65(10):1205-12.

Silverman ME, Loudon H. Antenatal reports of pre-pregnancy abuse is associated with symptoms of depression in the postpartum period. *Arch Womens Ment Health.* 2010;13(5):411-5.

Simon GE, VonKorff M, Ustun TB, Gater R, Gureje O, Sartorius N. Is the lifetime risk of depression actually increasing? *J Clin Epidemiol.* 1995 Sep;48(9):1109-18.

Simon GE, VonKorff M. Recognition, management, and outcomes of depression in primary care. *Arch Fam Med.* 1995 Feb;4(2):99-105.

Simon GE, VonKorff M, Piccinelli M, Fullerton C, Ormel J. An international study of the relation between somatic symptoms and depression. *New Engl J Med.* 1999;341(18):1329-35.

Simon GE, VonKorff M, Rutter C, Wagner E. Randomised trial of monitoring, feedback, and management of care by telephone to improve treatment of depression in primary care. *Br Med J.* 2000 Feb 26;320(7234):550-4.

Simon GE, Goldberg DP, Von Korff M, Ustun TB. Understanding cross-national differences in depression prevalence. *Psychol Med.* 2002 May;32(4):585-94.

Simon G, VonKorff M. Suicide mortality among patients treated for depression in an insured population. *Am J Epidemiol.* 1998;147:155-60.

Sinoff G, Ore L, Zlotogorsky D, Tamir A. Short Anxiety Screening Test--A brief instrument for detecting anxiety in the elderly. *Int J Geriatr Psychiatry.* 1999 Dec;14(12):1062-71.

Sipos K, Bodo M, May Z, Lendvai B, Piros A, Spitzer N, et al. Risk of mental disorders, their changes and somatic consideration in rural Hungary. *Ideggyogyaszati Szemle.* 2008 Mar 30;61(3-4):97-105.

Sirey JA, Bruce ML, Carpenter M, Booker D, Reid MC, Newell KA, et al. Depressive symptoms and suicidal ideation among older adults receiving home delivered meals. *Int J Geriatr Psychiatry*. 2008 Dec;23(12):1306-11.

Skarstein J, Aass N, Fossa SD, Skovlund E, Dahl AA. Anxiety and depression in cancer patients: Relation between the Hospital Anxiety and Depression Scale and the European Organization for Research and Treatment of Cancer Core Quality of Life Questionnaire. *J Psychosom Res*. 2000;49(1):27-34.

Sleath BL, Thomas N, Jackson E, West SL, Gaynes BN. Physician reported communication about depression and psychosocial issues during postpartum visits. *North Carolina Medical Journal*. 2007 May;68(3):151-5.

Slovacek L, Slovackova B, Slanska I, Priester P, Petera J, Kopecky J, et al. Cancer and depression: A prospective study. *Neoplasma*. 2009;56(3):187-93.

Slovacek L, Slanska I, Slovackova B, Petera J, Filip S, Priester P, et al. Screening for depression in survivors of metastatic ovarian cancer in a programme of palliative cancer care. *Bratisl Lek Listy*. 2009;110(10):655-9.

Smith DC, Munroe ML, Foglia LM, Nielsen PE, Deering SH. Effects of deployment on depression screening scores in pregnancy at an army military treatment facility. *Obstetrics and Gynecology*. 2010 Sep;116(3):679-84.

Smith PJ, Attix DK, Weldon BC, Greene NH, Monk TG. Executive function and depression as independent risk factors for postoperative delirium. *Anesthesiol*. 2009;110(4):781-7.

Smolderen KG, Spertus JA, Vriens PW, Kranendonk S, Nooren M, Denollet J. Younger women with symptomatic peripheral arterial disease are at increased risk of depressive symptoms. *J Vasc Surg*. 2010 Sep;52(3):637-44.

Snijders TJ, De LFE, Klumpers UMH, Kappelle LJ, van GJ. Prevalence and predictors of unexplained neurological symptoms in an academic neurology outpatient clinic: An observational study. *J Neurol*. 2004;251(1):66-71.

Sogaard HJ. Choosing screening instrument and cut-point on screening instruments. A comparison of methods. *Scand J Public Health*. 2009;37(8):872-80.

Sogaard HJ, Bech P. Compensating for non-response in a study estimating the incidence of mental disorders in long-term sickness absence by a two-phased design. *Scand J Public Health*. 2010 Aug;38(6):625-32.

Sokoya OO, Baiyewu O. Geriatric depression in Nigerian primary care attendees. *Int J Geriatr Psychiatry*. 2003 Jun;18(6):506-10.

Solberg LI, Fischer LR, Wei F, Rush WA, Conboy KS, Davis TF, et al. A CQI intervention to change the care of depression: A controlled study. *Effect Clin Pract*. 2001 Nov;4(6):239-49.

Solomon DA, Keller MB, Leon AC, Mueller TI, Shea MT, Warshaw M, et al. Recovery from major depression. A 10-year prospective follow-up across multiple episodes. *Arch Gen Psychiatry*. 1997 Nov;54(11):1001-6.

Solomon DA, Leon AC, Maser JD, Truman CJ, Coryell W, Endicott J, et al. Distinguishing bipolar major depression from unipolar major depression with the screening assessment of depression-polarity (SAD-P). *J Clin Psychiatr*. 2006 Mar;67(3):434-42.

Sonnenberg CM, Beekman AT, Deeg DJ, and Tilburg V. Drug treatment in depressed elderly in the Dutch community. *Int J Geriatr Psychiatry*. 2003 Feb;18(2):99-104.

Sorenson DS, Tschetter L. Prevalence of negative birth perception, disaffirmation, perinatal trauma symptoms, and depression among postpartum women. *Psychiatr Care*. 2010 Jan;46(1):14-25.

Steadman HJ, Osher FC, Robbins PC, Case B, Samuels S. Prevalence of serious mental illness among jail inmates. *Psychiatr Serv*. 2009 Jun;60(6):761-5.

Steffens DC, Skoog I, Norton MC, Hart AD, Tschanz JT, Plassman BL, et al. Prevalence of depression and its treatment in an elderly population: The Cache County study. *Arch Gen Psychiatry*. 2000 Jun;57(6):601-7.

Stein MB, Kirk P, Prabhu V, Grott M, Terepa M. Mixed anxiety-depression in a primary-care clinic. *J Affect Disord*. 1995 May 17;34(2):79-84.

Steiner M. Postnatal depression: A few simple questions. *Fam Pract*. 2002 Oct;19(5):469-70.

Stek ML, Vinkers DJ, Gussekloo J, Van der Mast RC, Beekman AT, Westendorp RG. Natural history of depression in the oldest old: Population-based prospective study. *Br J Psychiatry*. 2006 Jan;188:65-9.

Stokes SC, Thompson LW, Murphy S, Gallagher-Thompson D. Screening for depression in immigrant Chinese-American elders: Results of a pilot study. *J Gerontol Soc Work*. 2001;36(1-2):27-44.

Strickland PL, Deakin JFW, Percival C, Dixon J, Gater RA, Goldberg DP. Bio-social origins of depression in the community: Interactions between social adversity, cortisol and serotonin neurotransmission. *Br J Psychiatry*. 2002;180(FEB.):168-73.

Stuart S, Couser G, Schilder K, O'Hara MW, Gorman L. Postpartum anxiety and depression: Onset and comorbidity in a community sample. *J Nervous Ment Dis*. 1998 Jul;186(7):420-4.

Suen LJ, Tusaie K. Is somatization a significant depressive symptom in older Taiwanese Americans? *Geriatr Nurs*. 2004 May;25(3):157-63.

Suhr JA, Gunstad J. Postconcussive symptom report: The relative influence of head injury and depression. *J Clin Exp Neuropsychol*. 2002;24(8):981-93.

Sutter AL, Leroy V, Dallay D, Verdoux H, Bourgeois M. Post-partum blues and mild depressive symptomatology at days three and five after delivery, A French cross sectional study. *J Affect Disord.* 1997 Jun;44(1):1-4.

Suvisaari J, Aalto-Setälä T, Tuulio-Henriksson A, Harkanen T, Saarni SI, Perala J, et al. Mental disorders in young adulthood. *Psychol Med.* 2009;39(2):287-99.

Swalm D, Brooks J, Doherty D, Nathan E, Jacques A. Using the Edinburgh Postnatal Depression Scale to screen for perinatal anxiety. *Arch Womens Ment Health.* 2010;13(6):515-22.

Swardfager W, Herrmann N, Dowlati Y, Oh P, Kiss A, Lanctot KL. Relationship between cardiopulmonary fitness and depressive symptoms in cardiac rehabilitation patients with coronary artery disease. *J Rehabil Med.* 2008 Mar;40(3):213-8.

Taha AB, Lee PY, Lin K, Siti RG, Hassan S. Anxiety and depression in the primary health care clinics and the use of complementary and alternative medicine. *Int Med J.* 2005;12(2):93-7.

Tamburrino MB, Lynch DJ, Nagel RW, Stadler N, Pauling T. Screening women in family practice settings: Association between depression and smoking cigarettes. *Fam Pract Res J.* 1994 Dec;14(4):333-7.

Tang GW, Dennis S, Comino E, Zwar N. Anxiety and depression in Chinese patients attending an Australian GP clinic. *Aust Fam Phys.* 2009 Jul;38(7):552-5.

Tanriverdi D, Okanli A, Sezgin S, Ekinçi M. Quality of life in patients with multiple sclerosis in Turkey: Relationship to depression and fatigue. *J Neurosci Nurs.* 2010 Oct;42(5):267-73.

Tashakori A, Shanesaz A, Rezapour A. Assessment of some potential risk factors of postpartum depression. *Pakistan J Med Sci.* 2009;25(2):261-4.

Teissedre F, Chabol H. Postnatal depression: A study of the predictive effects of postnatal anxiety. *Ir J Psychol Med.* 2003;20(4):111-4.

Teychenne M, Ball K, Salmon J. Physical activity, sedentary behavior and depression among disadvantaged women. *Health Educ Res.* 2010 Aug;25(4):632-44.

Thavichachart N, Intoh P, Thavichachart T, Meksupa O, Tangwongchai S, Sughondhabirrom A, et al. Epidemiological survey of mental disorders and knowledge attitude practice upon mental health among people in Bangkok Metropolis. *J Med Assoc Thailand.* 2001 Jun;84:Suppl-26.

Theison A-K, Geisthoff UW, Forstl H, Schroder SG. Agitation in the morning: Symptom of depression in dementia? *Int J Geriatr Psychiatry.* 2009;24(4):335-40.

Thomas JL, Wilk JE, Riviere LA, McGurk D, Castro CA, Hoge CW. Prevalence of mental health problems and functional impairment among active component and national guard soldiers 3 and 12 months following combat in Iraq. *Arch Gen Psychiatry.* 2010;67(6):614-23.

Thombs BD, Hudson M, Taillefer SS, Baron M, Markland J, Pope J, et al. Prevalence and clinical correlates of symptoms of depression in patients with systemic sclerosis. *Arthritis Care Res.* 2008;59(4):504-9.

Thompson MA, Unutzer J, Katon WJ, Geyman JP, Gimlett D, Walker EA. Detection and treatment of depressive syndromes in a rural island clinic. *J Am Board Fam Pract.* 1999 Mar;12(2):120-7.

Thongtang O, Sukhatunga K, Ngamthipwatthana T, Chulakadabba S, Vuthiganond S, Pooviboonsuk P, et al. Prevalence and incidence of depression in the Thai elderly. *J Med Assoc Thailand.* 2002 May;85(5):540-4.

Tiemeier H, van Tuijl HR, Hofman A, Kiliaan AJ, Breteler MM. Plasma fatty acid composition and depression are associated in the elderly: The Rotterdam Study. *Am J Clin Nutr.* 2003 Jul;78(1):40-6.

Tiemens BG, VonKorff M, Lin EH. Diagnosis of depression by primary care physicians versus a structured diagnostic interview: Understanding discordance. *Gen Hosp Psychiatry.* 1999;21:87-96.

Topbas M, Yaris F, Can G, Kapuca M, Sayar K. The effect of sociodemographic factors and medical status on depressive symptoms of the inhabitants of nursing home and community-dwelling elders in Trabzon-Turkey. *Klin Psikofarmacol Bulteni.* 2004;14(2):57-60.

Trestman RL, Ford J, Zhang W, Wiesbrock V. Current and lifetime psychiatric illness among inmates not identified as acutely mentally ill at intake in Connecticut's jails. *J Am Acad Psychiatr Law.* 2007;35(4):490-500.

Tsai K-W, Chiang J-K, Lee C-S. Undiagnosed depression in patients with type 2 diabetes and its associated factors. *Tzu Chi Med J.* 2008;20(1):44-8.

Tseng YF, Hsu CM, Liu YH, Chen CH. The experiences of prenatal depression among Taiwanese women. *J Adv Nurs.* 2008 Dec;64(5):495-503.

Tsunoda T, Yamashita R, Kojima Y, Takahara S. Risk factors for depression after kidney transplantation. *Transplant Proceed.* 2010;42(5):1679-81.

Tucer B, Yalcin BM, Ozturk A, Mazicioglu MM, Yilmaz Y, Kaya M. Risk factors for low back pain and its relation with pain related disability and depression in a Turkish sample. *Turk Neurosurg.* 2009 Oct;19(4):327-32.

Turner K, Piazzini A, Franza A, Fumarola C, Chifari R, Marconi AM, et al. Postpartum depression in women with epilepsy versus women without epilepsy. *Epilepsy Behav.* 2006;9(2):293-7.

Turner K, Piazzini A, Franza A, Marconi AM, Canger R, Canevini MP. Epilepsy and postpartum depression. *Epilepsia.* 2009;50(SUPPL. 1):24-7.

Turrina C, Caruso R, Este R, Lucchi F, Fazzari G, Dewey ME, et al. Affective disorders among elderly general practice patients. A two-phase survey in Brescia, Italy. *Br J Psychiatry.* 1994 Oct;165(4):533-7.

Tylee A. Depression in Europe: Experience from the DEPRES II survey. *Eur Neuropsychopharmacol.* 2000;10(SUPPL. 4):S445-S448.

Ugarriza DN. A descriptive study of postpartum depression. *Psychiatr Care.* 1995 Jul;31(3):25-9.

UK National Screening Committee. Second report of the UK National Screening Committee. London (UK): Departments of Health for England, Scotland, Northern Ireland and Wales; 2000.

Unutzer J, Patrick DL, Diehr P, Simon G, Grembowski D, Katon W. Quality adjusted life years in older adults with depressive symptoms and chronic medical disorders. *Int Psychogeriatr.* 2000 Mar;12(1):15-33.

Unutzer J, Patrick DL, Marmon T, Simon GE, Katon WJ. Depressive symptoms and mortality in a prospective study of 2,558 older adults. *Am J Geriatr Psychiatr.* 2002 Sep;10(5):521-30.

Upton MW, Evans M, Goldberg DP, Sharp DJ. Evaluation of ICD-10 PHC mental health guidelines in detecting and managing depression within primary care. *Br J Psychiatry.* 1999 Nov;175:476-82.

Valkamo M, Hintikka J, Niskanen L, Koivumaa-Honkanen H, Honkalampi K, Viinamaki H. Depression and associated factors in coronary heart disease. *Scand Cardiovasc J.* 2001;35(4):259-63.

Vamos EP, Mucsi I, Keszei A, Kopp MS, Novak M. Comorbid depression is associated with increased healthcare utilization and lost productivity in persons with diabetes: A large nationally representative Hungarian population survey. *Psychosom Med.* 2009 Jun;71(5):501-7.

Van Bastelaar KMP, Pouwer F, Geelhoed-Duijvestijn PHLM, Tack CJ, Bazelmans E, Beekman AT, et al. Diabetes-specific emotional distress mediates the association between depressive symptoms and glycaemic control in Type 1 and Type 2 diabetes. *Diabetic Med.* 2010;27(7):798-803.

Van der Weele GM, Gussekloo J, de Waal MW, de Craen AJ, Van der Mast RC. Co-occurrence of depression and anxiety in elderly subjects aged 90 years and its relationship with functional status, quality of life and mortality. *Int J Geriatr Psychiatry.* 2009 Jun;24(6):595-601.

van Haastregt JC, Zijlstra GA, van Rossum E, van Eijk JT, Kempen GI. Feelings of anxiety and symptoms of depression in community-living older persons who avoid activity for fear of falling. *Am J Geriatr Psychiatr.* 2008 Mar;16(3):186-93.

van Steenbergen-Weijnenburg KM, Van Puffelen AL, Horn EK, Nuyen J, Sytze Van DP, van Benthem TB, et al. More co-morbid depression in patients with Type-2 diabetes with multiple complications. An observational study at a specialized outpatient clinic. *Diabetic Med.* 2011;28(1):86-9.

Vazquez FL, Blanco V. Symptoms of depression and related factors among Spanish university students. *Psychol Rep.* 2006 Oct;99(2):583-90.

Vazquez FL, Blanco V. Prevalence of DSM-IV major depression among Spanish university students. *J Am Coll Health.* 2008 Sep;57(2):165-71.

Verger P, Lions C, Ventelou B. Is depression associated with health risk-related behaviour clusters in adults. *Eur J Public Health*. 2009;19(6):618-24.

Verkerk GJ, Denollet J, Van Heck GL, Van Son MJ, Pop VJ. Personality factors as determinants of depression in postpartum women: A prospective 1-year follow-up study. *Psychosom Med*. 2005 Jul;67(4):632-7.

Verney SP, Jervis LL, Fickenscher A, Roubideaux Y, Bogart A, Goldberg J. Symptoms of depression and cognitive functioning in older American Indians. *Aging Ment Health*. 2008 Jan;12(1):108-15.

Vik K, Aass IM, Willumsen AB, Hafting M. "It's about focusing on the mother's mental health": Screening for postnatal depression seen from the health visitors' perspective--A qualitative study. *Scand J Public Health*. 2009 May;37(3):239-45.

Vilalta-Franch J, Garre-Olmo J, Lopez-Pousa S, Turon-Estrada A, Lozano-Gallego M, Hernandez-Ferrandiz M, et al. Comparison of different clinical diagnostic criteria for depression in Alzheimer disease. *Am J Geriatr Psychiatry*. 2006;14(7):589-97.

Vinkers DJ, Gussekloo J, Stek ML, Westendorp RG, Van der Mast RC. The 15-item Geriatric Depression Scale (GDS-15) detects changes in depressive symptoms after a major negative life event. The Leiden 85-plus Study. *Int J Geriatr Psychiatry*. 2004 Jan;19(1):80-4.

von Heideken W, Gustavsson JM, Lundin-Olsson L, Kallin K, Nygren B, Lundman B, et al. Health status in the oldest old. Age and sex differences in the Umea 85+ Study. *Aging Clin Exp Res*. 2006 Apr;18(2):116-26.

von Knorring L, Akerblad AC, Bengtsson F, Carlsson A, Ekselius L. Cost of depression: Effect of adherence and treatment response. *Eur Psychiatry*. 2006 Sep;21(6):349-54.

Voros V, Osvath P, Kovacs L, Varga J, Fekete S, Kovacs A. Screening for suicidal behaviour and mental disorders with Prime-MD questionnaire in general practice. *Prim Care Community Psychiatry*. 2006;11(4):193-6.

Vuorilehto M, Melartin T, Isometsa E. Depressive disorders in primary care: Recurrent, chronic, and comorbid. *Psychol Med*. 2005 May;35(5):673-82.

Wada T, Ishine M, Sakagami T, Okumiya K, Fujisawa M, Murakami S, et al. Depression in Japanese community-dwelling elderly - Prevalence and association with ADL and QOL. *Arch Gerontol Geriatr*. 2004;39(1):15-23.

Wada T, Ishine M, Sakagami T, Kita T, Okumiya K, Mizuno K, et al. Depression, activities of daily living, and quality of life of community-dwelling elderly in three Asian countries: Indonesia, Vietnam, and Japan. *Arch Gerontol Geriatr*. 2005;41(3):271-80.

Walker LG, Cordiner CM, Gilbert FJ, Needham G, Deans HE, Affleck IR, et al. How distressing is attendance for routine breast screening? *Psycho Oncol*. 1994;3(4):299-304.

Walters BAJ, Hays RD, Spritzer KL, Fridman M, Carter WB. Health-related quality of life, depressive symptoms, anemia, and malnutrition at hemodialysis initiation. *Am J Kidney Dis.* 2002;40(6):1185-94.

Wang J, Hausermann M, Ajdacic-Gross V, Aggleton P, Weiss MG. High prevalence of mental disorders and comorbidity in the Geneva Gay Men's Health Study. *Soc Psychiatry Psychiatr Epidemiol.* 2007 May;42(5):414-20.

Wang SJ, Liu HC, Fuh JL, Liu CY, Wang PN, Lu SR. Comorbidity of headaches and depression in the elderly. *Pain.* 1999 Sep;82(3):239-43.

Warner CM, Warner CH, Breitbach J, Rachal J, Matuszak T, Grieger TA. Depression in entry-level military personnel. *Mil Med.* 2007;172(8):795-9.

Warner R, Appleby L, Whitton A, Faragher B. Demographic and obstetric risk factors for postnatal psychiatric morbidity. *Br J Psychiatry.* 1996 May;168(5):607-11.

Wasserman GA, McReynolds LS, Ko SJ, Katz LM, Cauffman E, Haxton W, et al. Screening for emergent risk and service needs among incarcerated youth: Comparing MAYSI-2 and Voice DISC-IV. *J Am Acad Child Adolesc Psychiatry.* 2004;43(5):629-39.

Watts SC, Bhutani GE, Stout IH, Ducker GM, Cleator PJ, McGarry J, et al. Mental health in older adult recipients of primary care services: Is depression the key issue? Identification, treatment and the general practitioner. *Int J Geriatr Psychiatry.* 2002 May;17(5):427-37.

Webber AP, Martin JL, Harker JO, Josephson KR, Rubenstein LZ, Alessi CA. Depression in older patients admitted for postacute nursing home rehabilitation. *J Am Geriatr Soc.* 2005;53(6):1017-22.

Webster ML, Thompson JM, Mitchell EA, Werry JS. Postnatal depression in a community cohort. *Aust NZ J Psychiatry.* 1994 Mar;28(1):42-9.

Wedding U, Koch A, Rohrig B, Pientka L, Sauer H, Hoffken K, et al. Requestioning depression in patients with cancer: Contribution of somatic and affective symptoms to Beck's Depression Inventory. *Ann Oncol.* 2007 Nov;18(11):1875-81.

Wedding U, Koch A, Rohrig B, Pientka L, Sauer H, Hoffken K, et al. Depression and functional impairment independently contribute to decreased quality of life in cancer patients prior to chemotherapy. *Acta Oncol.* 2008;47(1):56-62.

Weingardt KR, Hsu J, Dunn ME. Brief screening for psychological and substance abuse disorders in veterans with long-term spinal cord injury. *Rehabil Psychol.* 2001;46(3):271-8.

Weintraub D, Xie S, Karlawish J, Siderowf A. Differences in depression symptoms in patients with Alzheimer's and Parkinson's diseases: Evidence from the 15-item Geriatric Depression Scale (GDS-15). *Int J Geriatr Psychiatry.* 2007;22(10):1025-30.

Weiss D. The prevalence of depression in male sex addicts residing in the United States. *Sex Addict Compuls*. 2004;11(1-2):57-69.

Wetherell JL, Arean PA. Psychometric evaluation of the Beck Anxiety Inventory with older medical patients. *Psychol Assess*. 1997;9(2):136-44.

White ML, Groh CJ. Depression and quality of life in women after a myocardial infarction. *J Cardiovasc Nurs*. 2007;22(2):138-44.

Wickberg B, Hwang CP. Screening for postnatal depression in a population-based Swedish sample. *Acta Psychiatrica Scandinavica*. 1997 Jan;95(1):62-6.

Williams JBW, Ellis A, Middleton A, Kobak KA. Primary care patients in psychiatric clinical trials: A pilot study using videoconferencing. *Ann Geriatr Psychiatr*. 2007;6 , 2007. Article Number: 24. Date of Publication: 04 Oct 2007.

Williams JWJ, Mulrow CD, Kroenke K, Dhanda R, Badgett RG, Omori D, and Lee S. Case-finding for depression in primary care: A randomized trial. *Am J Med*. 1999;106(1):36-43.

Willrich A, Pinzur M, McNeil M, Juknelis D, Lavery L. Health related quality of life, cognitive function, and depression in diabetic patients with foot ulcer or amputation. A preliminary study. *Foot Ankle Int*. 2005;26(2):128-34.

Winrow AM, Holmes JD. Old age medical patients screening positive for depression. *Ir J Psychol Med*. 2005;22(4):124-7.

Withall A, Harris LM, Cumming SR. The relationship between cognitive function and clinical and functional outcomes in major depressive disorder. *Psychol Med*. 2009;39(3):393-402.

Wittchen H-U, Kessler RC, Beesdo K, Krause P, Hofler M, Hoyer J. Generalized anxiety and depression in primary care: Prevalence, recognition, and management. *J Clin Psychiatr*. 2002;63(SUPPL. 8):24-34.

Wittink MN, Bogner HR. Primary care physicians' assessments of older patients' health and psychological status and recommendation of mammography. *J Am Board Fam Med*. 2008 Jan;21(1):17-23.

Wong KW, Chao DVK, Fung R, Lam PT. The prevalence of depressive symptoms in a regional geriatric day hospital. *Hong Kong Pract*. 2004;26(4):172-9.

Wong SY, Woo J, Hong AW, Leung JC, Leung PC. Clinically relevant depressive symptoms and peripheral arterial disease in elderly men and women. Results from a large cohort study in Southern China. *J Psychosom Res*. 2007 Nov;63(5):471-6.

Wu B, Chi I, Plassman BL, Guo M. Depressive symptoms and health problems among Chinese immigrant elders in the US and Chinese elders in China. *Aging Ment Health*. 2010 Aug;14(6):695-704.

Wu LR, Parkerson GR, Jr., Doraiswamy PM. Health perception, pain, and disability as correlates of anxiety and depression symptoms in primary care patients. *J Am Board Fam Pract.* 2002 May;15(3):183-90.

Yamamoto Y, Hayashino Y, Akiba T, Akizawa T, Asano Y, Saito A, et al. Depressive symptoms predict the subsequent risk of bodily pain in dialysis patients: Japan dialysis outcomes and practice patterns study. *Pain Med.* 2009;10(5):883-9.

Yeung A, Chan R, Mischoulon D, Sonawalla S, Wong E, Nierenberg AA, et al. Prevalence of major depressive disorder among Chinese-Americans in primary care. *Gen Hosp Psychiatry.* 2004 Jan;26(1):24-30.

Ying DG, Jiang S, Yang H, Zhu S. Frequency of generalized anxiety disorder in Chinese primary care. *Postgrad Med.* 2010 Jul;122(4):32-8.

Yonkers KA, Ramin SM, Rush AJ, Navarrete CA, Carmody T, March D, et al. Onset and persistence of postpartum depression in an inner-city maternal health clinic system. *Am J Psychiatry.* 2001 Nov;158(11):1856-63.

Yoshida K, Yamashita H, Ueda M, Tashiro N. Postnatal depression in Japanese mothers and the reconsideration of 'Satogaeri bunben'. *Pediatr Int.* 2001 Apr;43(2):189-93.

Yoshida S, Hirai M, Suzuki S, Awata S, Oka Y. Neuropathy is associated with depression independently of health-related quality of life in Japanese patients with diabetes. *Psychiatry Clin Neurosci.* 2009 Feb;63(1):65-72.

Zayas LH, Jankowski KRB, McKee MD. Prenatal and postpartum depression among low-income Dominican and Puerto Rican women. *Hispanic J Behav Sci.* 2003 Aug;25(3):370-85.

Zelikovsky N, Schast AP, Jean-Francois D. Parent stress and coping: Waiting for a child to receive a kidney transplant. *J Clin Psychol Med Set.* 2007;14(4):320-9.

Zelkowitz P, Milet TH. Screening for post-partum depression in a community sample. *Can J Psychiatry.* 1995 Mar;40(2):80-6.

Zelkowitz P, Saucier JF, Wang T, Katofsky L, Valenzuela M, Westreich R. Stability and change in depressive symptoms from pregnancy to two months postpartum in childbearing immigrant women. *Arch Womens Ment Health.* 2008;11(1):1-11.

Zivin K, Eisenberg D, Gollust SE, Golberstein E. Persistence of mental health problems and needs in a college student population. *J Affect Disord.* 2009 Oct;117(3):180-5.

Zoger S, Svedlund J, Holgers KM. Relationship between tinnitus severity and psychiatric disorders. *Psychosomatics.* 2006;47(4):282-8.

Zubaran C, Foresti K, Thorell MR, Franceschini P, Homero W. Depressive symptoms in crack and inhalant users in Southern Brazil. *J Ethn Subst Abuse*. 2010 Jul;9(3):221-36.

Zunzunegui MV, Beland F, Llacer A, Leon V. Gender differences in depressive symptoms among Spanish elderly. *Soc Psychiatry Psychiatr Epidemiol*. 1998 May;33(5):195-205.

Zuvekas S. Prescription drugs and the changing patterns of treatment for mental disorders, 1996-2001. *Health Affairs*. 2005;24:195-205.

Excluded: Not an eligible study design

Adult screening for depression. *Nurse Pract*. 1986; 21(5):82-98. PMID:8734628.

Screening identifies individuals with untreated depression. *J Psychosoc Nurs Ment Health Serv*. 1994; 32(1):45-6. PMID:8145198.

The diagnosis is depression. *Tecnologica*. 1995; 1-6. PMID:10168301.

New NMHA Web site features depression screening tool. *J Psychosoc Nurs Ment Health Serv*. 2000; 38(8):10.

Screening for depression with two questions. *S Afr Med J*. 2004; 94(2):100.

Screening for and treating male depression. *Geriatrics*. 2004; 59(10):46. PMID:15508556.

Screening can uncover depression in heart failure patients. *Cardiol Rev*. 2005; 22(2):10.

Accuracy of depression evaluation in hemodialysis patients. *Nat Clin Pract Nephrol*. 2006; 2(8):409-10.

Psychosocial risk factors: Perinatal screening and intervention. *Obstetrics and Gynecology*. 2006; 108(2):469-77.

Summaries for patients. Screening for depression in adults: U.S. Preventive Services Task Force recommendations. *Ann Intern Med*. 2009; 151(11):I56.

Depression screening effective only with follow-up: USPSTF. *J Natl Med Assoc*. 2010; 102(3):261-2.

Abas M. Depression and anxiety among older Caribbean people in the UK: Screening, unmet need and the provision of appropriate services. *Int J Geriatr Psychiatry*. 1996; 11(4):377-82.

Adkins B, Titus-Howard T, Massey V, Washburn N, Molinaro J, Lang B, Bruning WL, Simmons S, Gerken P, Sommer V, LaNoue K, Rogers B, and Wilson B. Recognizing depression in cancer outpatients. *Community Oncol*. 2005; 2(6):528-33.

Albani C, Bailer H, Blaser G, Brahler E, Geyer M, and Grulke N. Two-question depression-screeners - The solution to all problems? *Wien Med Wochenschr.* 2006; 156(7-8):185-8. PMID:16823535.

Albert C. The dark days of postpartum depression. Primary care screening is essential. *Adv Nurse Pract.* 2002; 10(6):67-70. PMID:12400366.

Albus C, Jordan J, and Herrmann-Lingen C. Screening for psychosocial risk factors in patients with coronary heart disease-Recommendations for clinical practice. *Eur J Cardiovasc Prev Rehabil.* 2004; 11(1):75-9.

Alexopoulos GS, Bruce ML, Schulberg HC, Katz I, TenHave T, Brown GK, Reynolds CF, and Mulsant BH. Interventions for depressed elderly primary care patients. *Int J Geriatr Psychiatry.* 2001; 16(6):553-9.

Alexopoulos GS, Borson S, Cuthbert BN, Devanand DP, Mulsant BH, Olin JT, and Oslin DW. Assessment of late life depression. *Biol Psychiatry.* 2002; 52(3):164-74. PMID:12182923.

Ali BS and Amanullah S. A comparative review of two screening instruments: The Aga Khan University Anxiety and Depression Scale and the self reporting questionnaire. *JPMA.* 1998; 48(3):79-82.

Amadori K, Herrmann E, and Pullen RK. Comparison of the 15-item geriatric depression scale (GDS-15) and the GDS-4 during screening for depression in an in-patient geriatric patient group. *J Am Geriatr Soc.* 2011; 59(1):171-2. PMID:21226694.

Aubry JM, Weber RB, Miguez M, Dubuis V, Bizzini V, Gex-Fabry M, Bondolfi G, and Gervasoni N. Performance of the mood disorder questionnaire (MDQ) according to bipolar subtype and symptom severity. *Bipolar Disord.* 2009; Conference: 8th International Conference on Bipolar Disorder Pittsburgh, PA United States. Conference Start: 20090625 Conference End: 20090627. Conference: 8th International Conference on Bipolar Disorder Pittsburgh, PA United States. Conference Start: 20090625 Conference End: 20090627. Conference Publication:(var.pagings):16-7.

Baldessarini RJ and Viguera AC. Perinatal screening for depression. *Am J Obstet Gynecol.* 2010; 203(2):e16. PMID:20417471.

Barbui C and Tansella M. Identification and management of depression in primary care settings. A meta-review of evidence. *Epidemiologia e Psichiatria Sociale.* 2006; 15(4):276-83.

Barclay L and Sydney BL. EDPS and clinical depression. *Aust J Adv Nurs.* 1994; 12(1):4. PMID:7786455.

Barrie MA. Objective screening tools to assess cognitive impairment and depression. *Top Geriatr Rehabil.* 2002; 18(2):28-46.

Beacham TD, May M, Williams PR, Schenk L, Askew R, Walker J, and Norwood A. Assessing postpartum depression in women: The home health nurse perspective. *Home Healthcare Nurse.* 561; 26(9):552-60. PMID:18849723.

Beck CT. State of the science on postpartum depression: What nurse researchers have contributed-part 2. *Am J Matern Child Nurs.* 157; 33(3):151-6. PMID:18453904.

Beck CT. Screening methods for postpartum depression. *JOGNN.* 1995; 24(4):308-12. PMID:7643261.

Beck CT. Postpartum depression. Stopping the thief that steals motherhood. *AWHONN Lifelines.* 1999; 3(4):41-4. PMID:10690011.

Beck CT. Recognizing and screening for postpartum depression in mothers of NICU infants. *Adv Neonatal Care.* 2003; 3(1):37-46. PMID:12882180.

Beck CT. Postpartum depression: It isn't just the blues. *Am J Nurs.* 2006; 106(5):40-50.

Bennett HA and Einarson TR. Depressive symptoms among women screened in obstetrics settings. *J Womens Health.* 2004; 13(1):119-Feb. PMID:15006284.

Bertram S, Wollan P, Kurland M, Palmer S, and Yawn B. Which U.S. women are most likely to have postpartum depression? *J Affect Disord.* 2010; Conference: 5th Biennial Meeting of the International Society for Affective Disorders, ISAD Vancouver, BC Canada. Conference Start: 20100416 Conference End: 20100419 Sponsor: World Psychiatric Association. Conference: 5th Biennial Meeting of the International Society for Affective Disorders, ISAD Vancouver, BC Canada. Conference Start: 20100416 Conference End: 20100419 Sponsor: World Psychiatric Association. Conference Publication:(var.pagings):S55-S56.

Bishop KK. Utilization of the Stetler model: Evaluating the scientific evidence on screening for postpartum depression risk factors in a primary care setting. *Kentucky Nurse.* 2007; 55(1):7-Mar. PMID:17348601.

Blazer DG. Psychiatry and the oldest old. *Am J Psychiatry.* 2000; 157(12):1915-24.

Bogen D. Electronic reminder systems increase screening for postpartum depression. *J Pediatr.* 2009; 155(5):758-9.

Bond S. Edinburgh Postnatal Depression Scale more efficient than others when screening for postpartum depression. *J Midwifery Womens Health.* 2008; 53(5):473-4.

Bonin-Guillaume S, Clement JP, Chassain AP, and Leger JM. Psychometric evaluation of depression in the elderly subject: Which instruments? What are the future perspectives? *Encephale.* 1995; 21(1):25-34. PMID:7720619.

Boyd RC, Le HN, and Somberg R. Review of screening instruments for postpartum depression. *Arch Womens Ment Health.* 2005; 8(3):141-53. PMID:16133785.

Brasic JR. Monitoring people treated with isotretinoin for depression. *Psychol Rep.* 2007; 100(3 II):1312-4.

Brody DS. Improving the management of depression in primary care: Recent accomplishments and ongoing challenges. *Dis Manag Health Outcome*. 2003; 11(1):21-31.

Bromberger JT. A psychosocial understanding of depression in women: For the primary care physician. *J Am Med Womens Assoc*. 2004; 59(3):198-206. PMID:15354373.

Brown E, Kaiser RM, and Gellis ZD. Screening and assessment of late-life depression in home healthcare: Issues and challenges. *Ann Long Term Care*. 2007; 15(10):27-32.

Bruhl KG, Luijendijk HJ, and Muller MT. Nurses' and nursing assistants' recognition of depression in elderly who depend on long-term care. *J Am Med Dir Assoc*. 2007; 8(7):441-5. PMID:17845946.

Brunell WE and Wells KB. Treatment of dysthymia with antidepressants. *Am J Psychiatry*. 1995; 152(7):1104-5.

Buist A. Mental health in pregnancy: The sleeping giant. *Australas Psychiatr*. 2002; 10(3):203-6.

Bunevicius A, Kusminskas L, and Bunevicius R. Validity of the Edinburgh postnatal depression scale. *Eur Psychiatry*. 2009; Conference: 17th European Psychiatric Association, EPA Congress Lisbon Portugal. Conference Start: 20090124 Conference End: 20090128. Conference: 17th European Psychiatric Association, EPA Congress Lisbon Portugal. Conference Start: 20090124 Conference End: 20090128. Conference Publication:(var.pagings):S896.

Campbell LC, Clauw DJ, and Keefe FJ. Persistent pain and depression: A biopsychosocial perspective. *Biol Psychiatr*. 2003; 54(3):399-409. PMID:12893114.

Cantwell R and Smith S. Prediction and prevention of perinatal mental illness. *Psychiatry*. 2009; 8(1):21-7.

Carson VB and Vanderhorst KJ. OASIS-C, depression screening, and M1730: Additional screening is necessary: The value of using standardized assessments. *Home Healthcare Nurse*. 2010; 28(3):183-90. PMID:20308812.

Chaudron LH. Postpartum depression: What pediatricians need to know. *Pediatr Rev*. 2003; 24(5):154-60.

Choi, J. Development and evaluation of a computer-interpretable guideline for depression screening and initial management in primary care. Jeeyae: Columbia U., US: Choi; 2006.

Citrome L and Flax JA. Depression screening. *Psychiatr Serv*. 1995; 46(2):186-7. PMID:7712262.

Clarke DM and McKenzie DP. Brief screening for depression. *J Am Geriatr Soc*. 1996; 44(2):212-3. PMID:8576517.

Clarkson P and Lynch S. Depression screening instruments. *Br J Gen Pract*. 1998; 48(429):1180-1. PMID:9687167.

Clayton A and Guico-Pabia C. Recognition of depression among women presenting with menopausal symptoms. *Menopause*. 2008; 15(4):758-67.

Coleman EA and Fox PD. Translating evidence-based geriatric care into practice: Lessons from managed care organizations part III: Depression and dementia. *Ann Long Term Care*. 2002; 10(11):33-40.

Coleman VH, Morgan MA, Zinberg S, and Schulkin J. Clinical approach to mental health issues among obstetrician-gynecologists: A review. *Obstetrical and Gynecological Survey*. 2006; 61(1):51-8.

Collins KA, Westra HA, Dozois DJA, and Burns DD. Gaps in accessing treatment for anxiety and depression: Challenges for the delivery of care. *Clin Psychol Rev*. 2004; 24(5):583-616.

Conwell Y, Van OK, and Caine ED. Suicide in older adults. *Psychiatr Clin North Am*. 2011; 34(2):451-68.

Copeland JR and Camus V. The use of screening tools in psychiatry of the elderly person. *Revue Medicale de la Suisse Romande*. 1999; 119(4):323-8. PMID:10361470.

Culpepper L, Clayton AH, Lieberman III JA, and Susman JL. Treating depression and anxiety in primary care. *Prim Care Comp J Clin Psychiatr*. 2008; 10(2):145-52.

Currie ML and Rademacher R. The pediatrician's role in recognizing and intervening in postpartum depression. *Pediatr Clin North Am*. 2004; 51(3):785-801. PMID:15157598.

Davies M. Should diabetes services routinely screen for psychological distress? *Practical Diabetes Int*. 2007; 24(9):483-6.

Davis JM and Gershtein CM. Screening for depression in patients with chronic illness: Why and how? *Dis Manag Health Outcome*. 2003; 11(6):375-8.

Dejesus RS, Vickers KS, Melin GJ, and Williams MD. A system-based approach to depression management in primary care using the Patient Health Questionnaire-9. *Mayo Clin Proceed*. 2007; 82(11):1395-402. PMID:17976360.

Dietrich AJ, Oxman TE, Williams JW, Jr., Kroenke K, Schulberg HC, Bruce M, and Barry SL. Going to scale: Re-engineering systems for primary care treatment of depression. *Ann Fam Med*. 2004; 2(4):301-4. PMID:15335127.

Ehde DM and Bombardier CH. Depression in persons with multiple sclerosis. *Phys Med Rehabil Clin North Am*. 2005; 16(2):437-48.

El-Sakka AI. Erectile dysfunction, depression, and ischemic heart disease: Does the existence of one component of this triad necessitate inquiring the other two? *J Sex Med*. 2011; 8(4):937-40.

Elliott SA and Leverton TJ. Is the EPDS a magic wand?: 2. 'Myths' and the evidence base. *J Reprod Infant Psychol*. 2000; 18(4):297-307.

Epker J and Block AR. Presurgical psychological screening in back pain patients: A review. *Clin J Pain*. 2001; 17(3):200-5.

Epstein-Lubow G, Gaudiano BA, Hinckley M, Salloway S, and Miller IW. Evidence for the validity of the American Medical Association's caregiver self-assessment questionnaire as a screening measure for depression. *J Am Geriatr Soc*. 2010; 58(2):387-8. PMID:20370867.

Erlangsen A, Nordentoft M, Conwell Y, Waern M, De LD, Lindner R, Oyama H, Sakashita T, Andersen-Ranberg K, Quinnett P, Draper B, Lapierre S, and International Research Group on Suicide Among the Elderly. Key considerations for preventing suicide in older adults: Consensus opinions of an expert panel. *Crisis*. 2011; 32(2):106-9. PMID:21616757.

Evins GG and Theofrastous JP. Postpartum depression: A review of postpartum screening. *Prim Care Update Ob/Gyns*. 1997; 4(6):241-6.

Farrell MK. Using functional assessment and screening tools with frail older adults. *Top Geriatr Rehabil*. 2004; 20(1):14-20.

Feinberg E, Smith MV, Morales MJ, Claussen AH, Smith DC, and Perou R. Improving women's health during internatal periods: Developing an evidenced-based approach to addressing maternal depression in pediatric settings. *J Womens Health*. 2006; 15(6):692-703.

Forbes W, Tracy DE, Hamilton RJ, and Eagles JM. A depression screening, day far the elderly in a general practice setting. *Psychiatr Bull*. 1995; 19(4):226-7.

Freedland KE, Carney RM, and Jaffe AS. Depression screening in patients with heart disease. *JAMA*. 2009; 301(13):1337.

Freeman MP. Screening and treatment for women with mood disorders associated with reproductive events. *J Clin Psychiatr*. 2007; 68(12):1946.

Fry LR. Prenatal screening. *Prim Care Clin Office Pract*. 2000; 27(1):55-69.

Ganguli M and Hendrie HC. Screening for cognitive impairment and depression in ethnically diverse older populations. *Alzheimer Dis Assoc Disord*. 2005; 19(4):275-8.

Garriock HA and Moreno FA. Genetics of depression: Implications for clinical practice yet? *Clin Neuropsychiatry*. 2011; 8(1):37-46.

Geddes JR, Freemantle N, Mason J, Eccles MP, and Boynton J. WITHDRAWN: Selective serotonin reuptake inhibitors (SSRIs) versus other antidepressants for depression. *Cochrane Database Sys Rev*. 2006; (3):CD001851. PM:17636689.

Gelenberg AJ. Using assessment tools to screen for, diagnose, and treat major depressive disorder in clinical practice. *J Clin Psychiatr*. 2010; 71(Suppl E1):e01. PMID:20371028.

Gilbody SM, Whitty PM, Grimshaw JM, and Thomas RE. Improving the detection and management of depression in primary care. *Qual Saf Health Care*. 2003; 12(2):149-55. PMID:12679514.

Gilliam FG. Diagnosis and treatment of mood disorders in persons with epilepsy. *Curr Opin Neurol*. 2005; 18(2):129-33.

Glogoski-Williams C. Recognition of depression in the older adult. *Occup Ther Ment Health*. 2000; 15(2):17-34.

Goldberg D. Psychological disorders in general medical settings. *Soc Psychiatr*. 1996; 31(1):1-2.

Goldberg JF. Ruling in and ruling out 'caseness' in the bipolar spectrum: Implications for initial screening. *Bipolar Disord*. 2010; 12(5):539-40.

Greenberg SA. The geriatric depression scale: Short form. *Am J Nurs*. 2007; 107(10):60-9.

Greiver M. Screening and long-term follow-up of depression in my practice. *Can Fam Physician*. 2007; 53(9):1445.

Gutknecht DR. Review: Ultra-short screening tests are not highly accurate for detecting depression in primary care. *Evid Base Med*. 2007; 12(4):120.

Hackley B, Sharma C, Kedzior A, and Sreenivasan S. Managing mental health conditions in primary care settings. *J Midwifery Womens Health*. 2010; 55(1):9-19.

Haddad M, Walters P, and Tylee A. Mood disorders in primary care. *Psychiatry*. 2009; 8(2):71-5.

Halbreich U. Prevalence of mood symptoms and depressions during pregnancy: Implications for clinical practice and research. *CNS Spectrums*. 2004; 9(3):177-84.

Harpole LH and Williams J. Assessment and management of depression in older adults. *Prim Psychiatry*. 2004; 11(5):31-6.

Hauser P. Neuropsychiatric side effects of HCV therapy and their treatment: Focus on IFN alpha-induced depression. *Gastroenterol Clin*. 2004; 33(1 Suppl):S35-S50.

Haythornthwaite JA. IMMPACT recommendations for clinical trials: Opportunities for the RDC/TMD. *JOR*. 2010; 37(10):799-806.

Hegadoren KM. Assessment and treatment of depression: A review in high-risk populations. *J Wound Ostomy Continence Nurs*. 2010; 31(5):259-72. PMID:15867724.

Henkel V, Moehrensclager M, Hegerl U, Moeller HJ, Ring J, and Worret WI. Screening for depression in adult acne vulgaris patients: tools for the dermatologist. *J Cosmet Dermatol*. 2002; 1(4):202-7. PMID:17147540.

Henshaw C. Clinical and biological aspects of postpartum blues and depression. *Curr Opin Psychiatr.* 2000; 13(6):635-8.

Herrmann C. Screening for major depression in a group of diabetic patients. *Psychosom Med.* 1997; 59(5):559-60. PMID:9316191.

Herrmann C. International experiences with the Hospital Anxiety and Depression Scale--A review of validation data and clinical results. *J Psychosom Res.* 1997; 42(1):17-41. PMID:9055211.

Hewitt CE, Gilbody SM, Mann R, and Brealey S. Instruments to identify post-natal depression: Which methods have been the most extensively validated, in what setting and in which language? *Int J Psychiatry Clin Pract.* 2010; 14(1):72-6.

Hirschfeld RMA. Mood disorder questionnaire: It's impact on the field. *Depress Anxiety.* 2010; 27(7):627-30.

Hirst KP and Moutier CY. Postpartum major depression. *Am Fam Physician.* 2010; 82(8):926-33. PMID:20949886.

Holmwood C, Jones DS, and Jackson-Bowers E. The depression puzzle. Do the pieces fit? *Aust Fam Phys.* 2001; 30(5):489-92. PMID:11432023.

Horwath E, Cohen R, and Weissman M. Epidemiology of depressive and anxiety disorders. Tsuang M, Tohen M, editors, In: *Textbook in Psychiatric Epidemiology.* 2nd ed ed. Hoeboken, NJ: John Wiley & Sons, Inc.; 2002. Chapter 16, p. 389-426.

Inott T. Is it delirium, dementia, or depression? *Nursing.* 2007; 37(11):65. PMID:17968283.

Isaac C and Schlife J. More than the blues: Perinatal depression. *Nebraska Nurse.* 1923; 40(3):8-13.

Jacobsen PB. Screening for psychological distress in cancer patients: Challenges and opportunities. *J Clin Oncol.* 2007; 25(29):4526-7. PMID:17846452.

Janati A. Suicidality, depression screening, and antiepileptic drugs: Reaction to the fda alert. *Neurology.* 2009; 73(20):1710.

Jolley SN and Betrus P. Comparing postpartum depression and major depressive disorder: Issues in assessment. *Issues Ment Health Nurs.* 2007; 28(7):765-80.

Jorm AF. The epidemiology of depressive states in the elderly: Implications for recognition, intervention and prevention. *Soc Psychiatry Psychiatr Epidemiol.* 1995; 30(2):53-9. PMID:7754416.

Katon W, Von Korff M, Lin E, Unutzer J, Simon G, Walker E, Ludman E, and Bush T. Population-based care of depression: Effective disease management strategies to decrease prevalence. *Gen Hosp Psychiatry.* 1997; 19(3):169-78. PMID:9218985.

Katon W and Ciechanowski P. Impact of major depression on chronic medical illness. *J Psychosom Res.* 2002; 53(4):859-63. PM:12377294.

Katona C and Livingston G. Impact of screening old people with physical illness for depression? *Lancet.* 2000; 356(9224):91-2. PMID:10963242.

Kemper KJ, Kelleher K, and Olson AL. Implementing maternal depression screening. *Pediatrics.*120(2):448-9. PMID:17671074.

Kemper KJ and Kelleher KJ. Rationale for family psychosocial screening. *Ambul Child Health.* 1996; 1(4):311-24.

Kemper KJ and Kelleher KJ. Family psychosocial screening: Instruments and techniques. *Ambul Child Health.* 1996; 1(4):325-39.

Kerr LK and Kerr LD, Jr. Screening tools for depression in primary care: The effects of culture, gender, and somatic symptoms on the detection of depression. *West J Med.* 2001; 175(5):349-52. PMID:11694495.

Keshen A and MacDonald JL. Postpartum depression: Making the case for routine screening. *McGill J Med.* 2002; 7(1):66-70.

Kessler R. Depression screening. *J Fam Pract.* 2003; 52(6):466. PMID:12836648.

Kimball AB, Gladman D, Gelfand JM, Gordon K, Horn EJ, Korman NJ, Korver G, Krueger GG, Strober BE, and Lebwohl MG. National Psoriasis Foundation clinical consensus on psoriasis comorbidities and recommendations for screening. *J Am Acad Dermatol.* 2008; 58(6):1031-42.

Kirchner JE, Curran GM, and Aikens J. Detecting depression in VA primary care clinics. *Psychiatr Serv.* 2004; 55(4):350.

Kirkcaldy RD and Tynes LL. Depression screening in a VA primary care clinic. *Psychiatr Serv.* 2006; 57(12):1694-6. PMID:17158481.

Klinkman MS, Schwenk TL, and Coyne JC. Depression in primary care - More like asthma than appendicitis: The Michigan depression project. *Can J Psychiatry.* 1997; 42(9):966-73.

Knight D. Health care screening for men who have sex with men. *Am Fam Physician.* 2004; 69(9):2149-56.

Knight D. CPD: Health care screening for men who have sex with men. *S Afr Fam Pract.* 2005; 47(4):34-9.

Koenig HG, Ford SM, and Sibert TE. Screening for depression in elderly patients. 1995; 119-130.

Kojima H. Treatment of depression in patients with coronary heart disease. *Am J Med.* 2007; 120(9):e13.

Koopmans RT, Zuidema SU, Leontjevas R, and Gerritsen DL. Comprehensive assessment of depression and behavioral problems in long-term care. *Int Psychogeriatr*. 2010; 22(7):1054-62. PMID:20843390.

Kornstein SG. Chronic depression in women. *J Clin Psychiatr*. 2002; 63(7):602-9.

Korsen N. Translating a guideline into practice: The USPSTF recommendations on screening for depression in adults. *Am Fam Physician*. 2010; 82(8):891. PMID:20949882.

Kowalenko T and Khare RK. Should we screen for depression in the emergency department? *Acad Emerg Med*. 2004; 11(2):177-8. PMID:14759961.

Krantz I, Eriksson B, Lundquist-Persson C, Ahlberg BM, and Nilstun T. Screening for postpartum depression with the Edinburgh Postnatal Depression Scale (EPDS): An ethical analysis. *Scand J Public Health*. 2008; 36(2):211-6. PMID:18519287.

Kripke C. Screening for depression. *Am Fam Physician*. 2006; 73(9):1550-1. PMID:16719246.

Krishnan KR, DeLong M, Kraemer H, Carney R, Spiegel D, Gordon C, McDonald W, Dew M, Alexopoulos G, Buckwalter K, Cohen PD, Evans D, Kaufmann PG, Olin J, Otey E, and Wainscott C. Comorbidity of depression with other medical diseases in the elderly. *Biol Psychiatr*. 2002; 52(6):559-88. PM:12361669.

Kurlowicz L. The Geriatric Depression Scale (GDS). *Director*. 1999; 7(4):134-5. PMID:10703343.

Kurlowicz LH and Streim JE. Measuring depression in hospitalized, medically ill, older adults. *Arch Psychiatr Nurs*. 1998; 12(4):209-18. PMID:9714940.

Lako I, Taxis K, Bruggeman R, Knegtering R, Wiersma D, and Slooff C. Prevalence of depressive symptoms and the effectiveness of antidepressants in routine clinical practice of schizophrenia. *Schizophrenia Res*. 2010; Conference: 2nd Schizophrenia International Research Society Conference, SIRS 2010 Florence Italy. Conference Start: 20100410 Conference End: 20100414. Conference: 2nd Schizophrenia International Research Society Conference, SIRS 2010 Florence Italy. Conference Start: 20100410 Conference End: 20100414. Conference Publication:(var.pagings):534-5.

Lange K, Coutant R, Danne T, Kapellen T, Pankowska E, Rami B, Aschemeier B, Blasig S, Hartmann R, Krug N, Marquardt E, Remus K, and Kordonouri O. High quality of life in children and psychological wellbeing in mothers 12 month after diabetes onset: Results of the paediatric onset-trial of sensor-enhanced CSII. *Pediatr Diabetes*. 2010; Conference: 36th Annual Meeting of the International Society for Pediatric and Adolescent Diabetes, ISPAD Buenos Aires Argentina. Conference Start: 20101027 Conference End: 20101030. Conference Publication:(var.pagings):101.

Le HN and Boyd RC. Prevention of major depression: Early detection and early intervention in the general population. *Clin Neuropsychiatry*. 2006; 3(1):6-22.

Liberto TL. Screening for depression and help-seeking in postpartum women during well-baby pediatric visits: An integrated review. *J Pediatr Health Care*. 2012; 26(2):109-17.

Lieberman III JA. Ask the professor: Depression/anxiety: Strategies for effective primary care. *Consultant*. 2001; 41(10):1469-73.

Lin KW and Lam C. Screening for depression in adults. *Am Fam Physician*. 2010; 82(8):985-6. PMID:20949894.

Lloyd-Williams M. Screening for depression in palliative care. *Am J Hospice Palliat Med*. 2001; 18(2):79-80. PMID:11407132.

Lloyd-Williams M. Screening for depression in palliative care patients: A review. *Eur J Canc Care*. 2001; 10(1):31-5. PMID:11827265.

Lo D, Paniagua MA, and Thomas DR. Missed opportunities: Nursing home providers' ability to identify and recognize depressive symptoms. *J Am Med Dir Assoc*. 2010; 11(9):680-3.

Lynch S, Clarkson P, Blenkiron P, and Fraser J. Scale-based protocols for the detection and management of depression. *Prim Care Psychiatr*. 2002; 8(3):77-84.

MacKnight C, Boydon A, and Rockwood K. Screening instruments for depression in elderly subjects. *Age Ageing*. 1998; 27(3):411.

Manoj H and Sivan M. Study to establish the usefulness of the one question 'Are you depressed?' in screening people with multiple sclerosis for depression. *Clin Rehab*. 2007; 21(12):1143. PMID:18042609.

Marcus SM and Heringhausen JE. Depression in childbearing women: When depression complicates pregnancy. *Prim Care Clin Office Pract*. 2009; 36(1):151-65.

Marsay C and Szabo CP. Screening for metabolic syndrome-Adherence to guidelines. *Afr J Psychiatr*. 2011; 14(1):64-6.

Marshall GN. Screening for psychiatric problems in the orofacial trauma setting. *Oral Maxillofac Surg Clin*. 2010; 22(2):225-9.

Martin CR. What does the Hospital Anxiety and Depression Scale (HADS) really measure in liaison psychiatry settings? *Curr Psychiatr Rev*. 2005; 1(1):69-73.

Martin CR and Cheng KKF. Issues in the psychiatric screening of cancer patients. *Curr Psychiatr Rev*. 2006; 2(1):3-10.

Martin CR and Thompson DR. Depression in coronary heart disease patients: Etiological and screening issues. *Curr Psychiatr Rev*. 2006; 2(2):245-54.

Masters N. Depression screening instrument. *Br J Gen Pract*. 1995; 45(397):439. PMID:7576851.

Matthey S. Detection and treatment of postnatal depression (perinatal depression or anxiety). *Curr Opin Psychiatr*. 2004; 17(1):21-9.

Matthey S. Are we overpathologising motherhood? *J Affect Disord.* 2010; 120(1-3):263-6. PMID:19628282.

Maurer D and Colt R. An evidence-based approach to the management of depression. *Prim Care Clin Office Pract.* 2006; 33(4):923-41.

McAlpine DD and Wilson AR. Screening for depression in primary care: What do we still need to know? *Depress Anxiety.* 2004; 19(3):137-45. PMID:15129415.

McDonald WM. New research in low-tech solutions to a significant public health problem: Assessment of depression in the elderly. *Am J Geriatr Psychiatr.* 2009; 17(7):537-41. PMID:19546652.

McElhaney JE. Depression in the elderly: Could you be missing the signs? *Consultant.* 2003; 43(4):566-73.

McWilliams S, Enudi W, Tessema H, Bhriain SN, and Swanwick G. Insight into moderate cognitive impairment is associated with depression in the elderly. *Int J Geriatr Psychiatry.* 2010; 25(2):214-5.

Mian AI. Depression in pregnancy and the postpartum period: Balancing adverse effects of untreated illness with treatment risks. *J Psychiatr Pract.* 2005; 11(6):389-96.

Mitchell AJ, Lord K, and Symonds P. A symptom level exploration of the PHQ9 in a UK cancer setting: Which is the optimal single question that predicts DSMIV major depression? *Psycho Oncol.* 2010; Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference Publication:(var.pagings):S282.

Mitchell AJ. Depression screening and management programmes with staff assistance in primary care increase response and remission rates, but programmes without staff assistance do not show benefits. *Evid Base Med.* 2010; 15(2):49-50.

Mitchell AJ, Lord K, Baker-Glenn E, and Symonds P. A large scale validation of the emotion thermometers as a screening tool for major depression in an ethnically diverse cancer population. *Psycho Oncol.* 2010; Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference Publication:(var.pagings):S97.

Mitchell AJ, Carter G, and Meader N. A definitive meta-analysis to ascertain the optimal screening method for depression in cancer and palliative care. Part I - Single test application. *Psycho Oncol.* 2010; Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference Publication:(var.pagings):S163.

Moghabghab R, Adler L, Banez C, Boutcher F, Perivolaris A, Rancoeur D-M, Spevakow D, Tully S, Wallace S, and Woo K. The experience of implementing nursing best practice guidelines for the screening of delirium, dementia and depression in the older adult. *Geriatr Aging*. 2003; 6(10):39-41.

Moran MB. Screening for depression. *J Am Geriatr Soc*. 1995; 43(4):457. PMID:7706644.

Morley KI, Hall WD, and Carter L. Genetic screening for susceptibility to depression: Can we and should we? *Aust NZ J Psychiatry*. 2004; 38(1-2):73-80. PMID:14731197.

Mui AC, Burnette D, and Chen LM. Cross-cultural assessment of geriatric depression: A review of the CES-D and the GDS. *J Ment Health Aging*. 2001; 7(1):137-64.

Musich SA, Schultz AB, Burton WN, and Edington DW. Overview of disease management approaches: Implications for corporate-sponsored programs. *Dis Manag Health Outcome*. 2004; 12(5):299-326.

Musselman DL, McDaniel JS, Porter MR, and Nemeroff CB. Screening for depression. *J Med Assoc Ga*. 1999; 88(1):47-53. PMID:10087715.

Muzik M, Marcus SM, Flynn H, and Rosenblum KL. Depression during pregnancy: Detection, comorbidity and treatment. *Asia Pacific Psychiatr*. 2010; 2(1):7-18.

Nardone DA and Smith SP. Screening for depression. *J Gen Intern Med*. 1997; 12(12):789-90. PMID:9436902.

Nitz, N. M. Older adults and depressive symptoms. Nicole Mariko: U Minnesota, US: Nitz; 2005.

Nylen KJ, Segre LS, and O'Hara MW. Public health implications of postpartum depression. *Psychiatr Ann*. 2005; 35(7):599-609.

O'Donoghue DL, Boissonneault GA, Brenneman AE, Essary AC, Fortier F, Heinan ML, Leger MM, McNellis R, Moreau T, Zarbock S, Clinical and Scientific Affairs Council, and AAPA. Major depression. Screening and treatment. *J Am Acad Phys Assist*. 2009; 22(11):18-9. PMID:19999171.

O'Hara MW and Gorman LL. Can postpartum depression be predicted? *Prim Psychiatry*. 2004; 11(3):42-7.

Ong PS. Late-life depression: Current issues and new challenges. *Ann Acad Med Singapore*. 2003; 32(6):764-70.

Pangilinan J, Kelly BM, and Pangilinan JM. Depression in the patient with brain cancer. *Community Oncol*. 2007; 4(9):533-9.

Panzer PE. It's your life ... take control Depression Screening Project. *Delaware Med J*. 2007; 79(9):357-9. PMID:17941419.

Parveen A, Ali I, Husain M, Farhan S, Bhatia MR, Rehman R, Ansari S, Jaffery F, Rahman A, Chaudhry IB, and Husain N. Prevalence of maternal depression in a paediatric clinic: A study from a low income

country. Eur Psychiatry. 2010; Conference: 18th European Congress of Psychiatry Munich Germany. Conference Start: 20100227 Conference End: 20100302. Conference: 18th European Congress of Psychiatry Munich Germany. Conference Start: 20100227 Conference End: 20100302. Conference Publication:(var.pagings).

Patten SB, Bilsker D, and Goldner E. The evolving understanding of major depression epidemiology: Implications for practice and policy. Can J Psychiatry. 2008; 53(10):689-95. PMID:18940037.

Patten SB. Major depression prevalence is very high, but the syndrome is a poor proxy for community populations' clinical treatment needs. Can J Psychiatry. 2008; 53(7):411-9. PMID:18674395.

Peach J, Koob JJ, and Kraus MJ. Psychometric evaluation of the Geriatric Depression Scale (GDS): Supporting its use in health care settings. Clin Gerontol. 2001; 23(3-4):57-68.

Pignone M, Gaynes B, Lohr K, Rushton JL, and Mulrow C. Screening for depression in adults. Ann Intern Med. 2003; 138:767-8.

Piwonski J, Sygnowska E, and Piwonska A. Cardiovascular risk factors in persons with and without depressive symptoms. Atherosclerosis Suppl. 2009; Conference: 15th International Symposium on Atherosclerosis Boston, MA United States. Conference Start: 20090614 Conference End: 20090618. Conference: 15th International Symposium on Atherosclerosis Boston, MA United States. Conference Start: 20090614 Conference End: 20090618. Conference Publication:(var.pagings).

Pomerantz JM. Screening for depression in primary care. Drug Benefit Trends. 2005; 17(6):273-4.

Pritchard E and Dewing J. Screening for dementia and depression in older people. Nursing Standard. 1999; 14(5):46-52. PMID:11061177.

Rait G and Burns A. Screening for depression and cognitive impairment in older people from ethnic minorities. Age Ageing. 1998; 27(3):271-5.

Raj YP. How primary care doctors can screen depression. Med Econ. 2008; 85(14):28-33. PMID:18756998.

Raphel S. New recommendations on screening and treatment for major depressive disorder in children and adolescents. J Child Adolesc Psychiatr Nurs. 2009; 22(3):170-1. PMID:19702972.

Rife JC. Assessing confusion and depression within the context of extended unemployment: The use of clinical screening instruments. Clin Gerontol. 1995; 15(2):67-70.

Riggs RM, McCarthy JM, and Kaufman L. The utility of the beck depression inventory in the evaluation of chronic pelvic pain patients. J Gynecol Surg. 2007; 23(4):127-32.

Rubin RR, Ciechanowski P, Egede LE, Lin EHB, and Lustman PJ. Recognizing and treating depression in patients with diabetes. Curr Diabetes Rep. 2004; 4(2):119-25.

Rush AJ, Polatin P, and Gatchel RJ. Depression and chronic low back pain: Establishing priorities in treatment. *Spine*. 2000; 25(20):2566-71.

Sahr N. Assessment and diagnosis of elderly depression. *Clin Excellence Nurs Pract*. 1999; 3(3):158-64. PMID:10646410.

Saks K, Tut E, Kaarik E, and Jaanson K. Depressive symptoms in older Estonians: Prevalence and models. *J Am Geriatr Soc*. 2002; 50(6):1164-5.

Sanderson K and Andrews G. Common mental disorders in the workforce: Recent findings from descriptive and social epidemiology. *Can J Psychiatry*. 2006; 51(2):63-75.

Saperia J, Ashby D, and Gunnell D. Suicidal behaviour and SSRIs: Updated meta-analysis. *Br Med J*. 2006; 332(7555):1453. PM:16777898.

Sarton, C. Postpartum depression: Timing, location of residence, and perceived stress. Cherylann: State U New York At Binghamton, US: Sarton; 2006.

Sato Y, Tanda H, Kato S, Onishi S, Nakajima H, Nitta T, Akagashi K, Hanzawa T, and Tobe M. Clinical analysis of major depressive disorder and serum testosterone level and aging males'f symptom score in self-referred patients in a late onset of hypogonadism clinic. *J Sex Med*. 2009; Conference: Joint Meeting of the European(ESSM):79-80.

Sauvanet JP. Diabetes and depression, from screening to intervention. *Soins - Psychiatrie*. 2010; (268):11-Jun. PMID:20540385.

Schade CP, Jones ER, Jr., and Wittlin BJ. A ten-year review of the validity and clinical utility of depression screening. *Psychiatr Serv*. 1998; 49(1):55-61. PMID:9444681.

Schwenk TL. Screening for depression in primary care. A disease in search of a test. *J Gen Intern Med*. 1996; 11(7):437-9. PMID:8842939.

Schwenk TL. Screening for depression in primary care. *JAMA*. 2000; 284(11):1379-80. PMID:10989389.

Segre LS, Stuart S, and O'Hara MW. Interpersonal psychotherapy for antenatal and postpartum depression. *Prim Psychiatry*. 2004; 11(3):52-6+66.

Sela RA. Screening for depression in palliative cancer patients attending a pain and symptom control clinic. *Palliat Support Care*. 2007; 5(3):207-17. PMID:17969824.

Sellick SM. Identifying and treating depression among older adults with cancer. *Geriatr Aging*. 2007; 10(5):288-91.

Sengupta S. Diagnosis of depression in general practice. *Indian J Med Sci*. 2005; 59(5):217-25.

Serby M and Yu M. Overview: depression in the elderly. *Mount Sinai J Med*. 2003; 70(1):38-44. PM:12516008.

Seritan AL, McCloud MK, and Hinton L. Geriatric depression - Review for primary care. *Curr Psychiatr Rev.* 2009; 5(2):137-42.

Sharma V, Burt VK, and Ritchie HL. Assessment and treatment of bipolar II postpartum depression: A review. *J Affect Disord.* 2010; 125(1-3):18-26.

Sharp BA, Meikle SF, James MD, Steiner C, and Remus D. NHQR/NHDR measures for women of reproductive age. *Medical Care.* 2005; 43(3 Suppl):I64-I71.

Sharp LK and Lipsky MS. Screening for depression across the lifespan: A review of measures for use in primary care settings. *Am Fam Physician.* 2002; 66(6):1001-8. PMID:12358212.

Shenassa ED. Delivering the goods: The importance of screening accuracy for effective community intervention and prevention. *J Community Psychol.* 2002; 30(2):197-210.

Sherman SE. Review: screening for depression reduces persistent depression. *ACP J Club.* 2002; 137(3):100-Dec.

Simon GE. Treating depression in patients with chronic disease: recognition and treatment are crucial; Depression worsens the course of a chronic illness. *West J Med.* 2001; 175(5):292-3. PM:11694462.

Simon GE. Evidence review: Efficacy and effectiveness of antidepressant treatment in primary care. *Gen Hosp Psychiatry.* 2002; 24(4):213-24. PM:12100832.

Singh AR and Newton WP. Several depression screening tools work equally well. *J Fam Pract.* 2002; 51(6):511.

Sit DK, Flint C, Svidergol D, White J, Wimer M, Bish B, and Wisner KL. Best practices: an emerging best practice model for perinatal depression care. *Psychiatr Serv.* 2009; 60(11):1429-31. PMID:19880455.

Smolderen KG, Buchanan DM, Amin AA, Gosch K, Nugent K, Riggs L, Seavey G, and Spertus JA. Real-world lessons from the implementation of a depression screening protocol in acute myocardial infarction patients implications for the american heart association depression screening advisory. *Circulation.* 2011; 4(3):283-92.

Snowden M, Steinman L, Frederick J, and Wilson N. Screening for depression in older adults: Recommended instruments and considerations for community-based practice. *Clin Geriatr.* 2009; 17(9):26-32.

Sobey WS. Barriers to postpartum depression prevention and treatment: A policy analysis. *J Midwifery Womens Health.* 2002; 47(5):331-6. PMID:12361344.

Solberg LI, Trangle MA, and Wineman AP. Follow-up and follow-through of depressed patients in primary care: The critical missing components of quality care. *J Am Board Fam Pract.* 2005; 18(6):520-7.

Soler-Lopez M and Reid M. Population screening for postnatal depression in Scotland. *MEDIFAM.* 2000; 10(8):481-7.

Sorocco KH and McCallum TJ. Mental health promotion in older adults: Addressing treatment approaches and available screening tools. *Geriatrics*. 2006; 61(1):19-22.

Staab JP, Datto CJ, Weinrieb RM, Gariti P, Rynn M, and Evans DL. Detection and diagnosis of psychiatric disorders in primary medical care settings. *Med Clin North Am*. 2001; 85(3):579-96. PMID:11349474.

Stacey BG. Assessment of anxiety and depression in primary care. *Med J Aust*. 1998; 168(7):368.

Stevens JC and Diehl SJ. Ob/Gyn residents as primary care providers: Implementing a new curriculum for diagnosing and treating depression and anxiety. *Prim Care Update Ob/Gyns*. 2003; 10(6):297-9.

Stewart D. Depression during pregnancy. *Can Fam Physician*. 2005; 51:1061-7. PMID:16121822.

Stiles PG and McGarrahan JF. The Geriatric Depression Scale: A comprehensive review. *J Clin Geropsychol*. 1998; 4(2):89-110.

Sukhwani P, Prasankumar RP, Hameed FA, Widmer KL, Winfrey KL, and Rodenhauer P. Public outreach depression screening: Collaboration between medical education and integrative primary care. *Southern Med J*. 2006; 99(9):1016-7. PMID:17004544.

Svandra P. Psychological problems and hospitalization: Recognizing in order to treat better. *Revue de L'Infirmiere*. 1998; (42):64-70. PMID:9847847.

Swanwick GR and Wrigley M. Screening for depression in elderly patients. *Am J Psychiatry*. 1998; 155(7):994-5. PMID:9659876.

Szygula-Jurkiewicz B, Duszanska A, and Polonski L. Is depression a problem in patients with chronic heart failure? *Polskie Archiwum Medycyny Wewnetrznej*. 2008; 118(1-2):52-6.

Taleb M, Rouillon F, Hegerl U, Hamdani N, and Gorwood P. Programmes against depression. *Encephale*. 2006; 32(1):9-15.

Thakur SS and Supiano MA. Screening for common clinical conditions in older women. *Clin Obstet Gynecol*. 2007; 50(3):767-75.

Thombs BD and Ziegelstein RC. The effect of a primary care practice-based depression intervention on mortality in older adults. *Ann Intern Med*. 2008; 148(3):244-5.

Timonen M and Liukkonen T. Management of depression in adults. *BMJ*. 2008; 336(7641):435-9.

Tylee A. Identifying and managing depression in primary care in the United Kingdom. *J Clin Psychiatr*. 2006; 67(SUPPL. 6):41-5.

Valente SM. Detecting and evaluating depression among elderly patients in home health. *Home Health Care Manag Pract*. 2005; 17(2):101-8.

Valente SM and Saunders J. Screening for depression and suicide: Self-report instruments that work. *J Psychosoc Nurs Ment Health Serv.* 2005; 43(11):22-31. PMID:16350912.

Vannotti M, Jacot-Sadovski I, and Cornuz J. The screening of major depression in primary care: Recommendations and clinical stakes. *Medecine et Hygiene.* 2002; 60(2411):2022-6.

Vieira T. When joy becomes grief. Screening tools for postpartum depression. *AWHONN Lifelines.* 2002; 6(6):506-13. PMID:12593259.

Vodermaier A, Linden W, Greig D, and Regina M. Prevalence rates of anxiety and depressive disorders in adult cancer patients - A consecutive complete assessment at the time point of cancer diagnosis. *Psycho Oncol.* 2010; Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference: IPOS 12th World Congress of Psycho-Oncology Quebec City, QC Canada. Conference Start: 20100525 Conference End: 20100529. Conference Publication:(var.pagings):S78.

von Knorring L. Review of Depression in women with focus on the postpartum period. *Nordic Journal of Psychiatry.* 2003; 57(5):390.

Wada T, Ishine M, Kita T, Fujisawa M, and Matsubayashi K. Depression screening of elderly community-dwelling Japanese. *J Am Geriatr Soc.* 2003; 51(9):1328-9. PMID:12919256.

Walker, K. E. Correlates of the scales of a modified screening version of the Multidimensional Pain Inventory with depression and anxiety on a chronic pain sample. Katherine Elise: U North Texas, US: Walker; 2010.

Wang P. Beyond screening: Improving the quality of treatment for depression. *Harv Rev Psychiatr.* 1999; 7(3):177-82. PMID:10483937.

Watson LC, Pignone MP, and Iliffe S. Review: Screening instruments had sensitivities of 67-100% and specificities of 53-98% for detecting major depression in older primary care patients. *Evid Base Med.* 2004; 9(4):124.

Weeks SK, Andreson MA, Harmon LS, and Michaels TK. Getting inside depression and suicide ideation. One comprehensive screening approach targets patients 4 years and older. *Holistic Nurs Pract.* 2005; 19(1):5-9. PMID:15789503.

Weinberger T, Forrester A, Markov D, Chism K, and Kunkel EJ. Women at a dangerous intersection: Diagnosis and treatment of depression and related disorders in patients with breast cancer. *Psychiatr Clin North Am.* 2010; 33(2):409-22. PMID:20385344.

Whelan PJ, Gaughran F, Walwyn R, Chatterton K, and Macdonald A. 'Do you feel that your life is empty?' The clinical utility of a one-off question for detecting depression in elderly care home residents. *Age Ageing.* 2008; 37(4):475-8.

Wilens JM and Mounts KO. Women with depression - "You can't tell by looking". *Matern Child Health J.* 2006; 10(SUPPL. 7):183-7.

Wincata J, Weiss M, Marquart B, and Alexandrowicz R. Screening instruments for general hospital and primary care patients. *Adv Psychosom Med.* 2004; 26:74-97. PMID:15326863.

Wright AF. Should general practitioners be testing for depression? *Br J Gen Pract.* 1994; 44(380):132-5. PMID:8204322.

Wright IA. Monitoring depression in patients undergoing alpha-interferon and ribavirin therapy for hepatitis C. *Gastroenterol Nurs.* 2000; 23(6):275-80.

Wright KM, Huffman AH, Adler AB, and Castro CA. Psychological screening program overview. *Mil Med.* 2002; 167(10):853-61.

Wu CM and Kelley LS. Choosing an appropriate depression assessment tool for Chinese older adults: A review of 11 instruments. The best tools take into account cultural differences. *J Gerontol Nurs.* 2007; 33(8):12-22. PMID:17718374.

Wylie L, Hollins Martin CJ, Marland G, Martin CR, and Rankin J. The enigma of post-natal depression: An update. *J Psychiatr Ment Health Nurs.* 2011; 18(1):48-58.

Yamamoto Y, Tanioka M, Hayashino Y, Mishina H, Kato M, Fukuhara S, Utani A, and Miyachi Y. Application of a two-question screening instrument to detect depressive symptoms in patients with vitiligo: A pilot study. *J Am Acad Dermatol.* 2011; 64(5):e69-e70. PMID:21496683.

Yohannes AM, Baldwin RC, and Connolly MJ. Depression and anxiety in elderly patients with chronic obstructive pulmonary disease. *Age Ageing.* 2006; 35(5):457-9.

Zalai DM and Novak M. Depressive disorders in patients with chronic kidney disease. *Prim Psychiatry.* 2008; 15(1):66-72.

Zamani BA and Ghoreishi FS. Prevalence of depressive disorder in rheumatoid arthritis patients in 2007-2009. *Int J Rheum Dis.* 2010; Conference: 14th Congress of Asia Pacific League of Associations for Rheumatology, APLAR 2010 Hong Kong Hong Kong. Conference Start: 20100711 Conference End: 20100715. Conference: 14th Congress of Asia Pacific League of Associations for Rheumatology, APLAR 2010 Hong Kong Hong Kong. Conference Start: 20100711 Conference End: 20100715. Conference Publication:(var.pagings):74.

Ziegelstein RC. Depression in older survivors of myocardial infarction. *Geriatr Aging.* 2003; 6(2):59-62.

Zimmer KP and Minkovitz CS. Maternal depression: An old problem that merits increased recognition by child healthcare practitioners. *Curr Opin Pediatr.* 2003; 15(6):636-40. PMID:14631212.

Excluded: Not an eligible population

Abiodun OA. Postnatal depression in primary care populations in Nigeria. *Gen Hosp Psychiatry*. 2006 Mar;28(2):133-6.

Ackermann RT, Rosenman MB, Downs SM, Holmes AM, Katz BP, Li J, et al. Telephonic case-finding of major depression in a Medicaid chronic disease management program for diabetes and heart failure. *Gen Hosp Psychiatry*. 2005 Sep;27(5):338-43.

Adewuya AO. Early postpartum mood as a risk factor for postnatal depression in Nigerian women. *Am J Psychiatry*. 2006 Aug;163(8):1435-7.

Adewuya AO, Ola BA, Dada AO, Fasoto OO. Validation of the Edinburgh Postnatal Depression Scale as a screening tool for depression in late pregnancy among Nigerian women. *J Psychosom Obstet Gynaecol*. 2006 Dec;27(4):267-72.

Ahmed K, Kelshiker A, Jenner C. The screening and treatment of undiagnosed depression in patients with chronic obstructive pulmonary disease (COPD) in a general practice. *Prim Care Respir J*. 2007 Aug;16(4):249-51.

Al Haddad MK. Depression in elderly primary care attendees in Bahrain. *Arab J Psychiatr*. 2000 May;11(1):48-55.

Alexander S, Palmer C, Stone PC. Evaluation of screening instruments for depression and anxiety in breast cancer survivors. *Breast Canc Res Treat*. 2010 Jul;122(2):573-8.

Allen K, Cull A, Sharpe M. Diagnosing major depression in medical outpatients: Acceptability of telephone interviews. *J Psychosom Res*. 2003 Oct;55(4):385-7.

Annunziato RA, Rubinstein D, Sheikh S, Maurer M, Cotter G, McKay MM, et al. Site matters: Winning the hearts and minds of patients in a cardiology clinic. *Psychosomatics*. 2008 Sep;49(5):386-91.

Armstrong S, Small R. Screening for postnatal depression: Not a simple task. *Aust NZ J Public Health*. 2007 Feb;31(1):57-61.

Aujla N, Abrams KR, Davies MJ, Taub N, Skinner TC, Khunti K. The prevalence of depression in white-European and South-Asian people with impaired glucose regulation and screen-detected type 2 diabetes mellitus. *PLoS ONE*. 2009;4(11):e7755.

Aujla N, Skinner TC, Khunti K, Davies MJ. The prevalence of depressive symptoms in a white European and South Asian population with impaired glucose regulation and screen-detected Type 2 diabetes mellitus: A comparison of two screening tools. *Diabetic Med*. 2010;27(8):896-905.

Austin M-P, Hadzi-Pavlovic D, Priest SR, Reilly N, Wilhelm K, Saint K, et al. Depressive and anxiety disorders in the postpartum period: How prevalent are they and can we improve their detection? *Arch Womens Ment Health*. 2010;13(5):395-401.

Awata S, Bech P, Yoshida S, Hirai M, Suzuki S, Yamashita M, et al. Reliability and validity of the Japanese version of the World Health Organization-Five Well-Being Index in the context of detecting depression in diabetic patients. *Psychiatry Clin Neurosci*. 2007 Feb;61(1):112-9.

Axford J, Butt A, Heron C, Hammond J, Morgan J, Alavi A, et al. Prevalence of anxiety and depression in osteoarthritis: Use of the Hospital Anxiety and Depression Scale as a screening tool. *Clin Rheumatol*. 2010 Nov;29(11):1277-83.

Aydin IO, Ulusahin A. Depression, anxiety comorbidity, and disability in tuberculosis and chronic obstructive pulmonary disease patients: Applicability of GHQ-12. *Gen Hosp Psychiatry*. 2001;23(2):77-83.

Baas KD, Wittkamp KA, van Weert HC, Lucassen P, Huyser J, van den Hoogen H, et al. Screening for depression in high-risk groups: Prospective cohort study in general practice. *Br J Psychiatry*. 2009 May;194(5):399-403.

Bagedahl-Strindlund M, Monsen BK. Postnatal depression: A hidden illness. *Acta Psychiatrica Scandinavica*. 1998 Oct;98(4):272-5.

Baker L, Oswald K. Screening for postpartum depression in a rural community. *Community Ment Health J*. 2008 Jun;44(3):171-80.

Ballestrem CL, Strauss M, Kachele H. Contribution to the epidemiology of postnatal depression in Germany--Implications for the utilization of treatment. *Arch Womens Ment Health*. 2005 May;8(1):29-35.

Barnett B, Matthey S, Gyaneshwar R. Screening for postnatal depression in women of non-English speaking background. *Arch Womens Ment Health*. 1999;2(2):67-74.

Bass JK, Ryder RW, Lammers MC, Mukaba TN, Bolton PA. Post-partum depression in Kinshasa, Democratic Republic of Congo: Validation of a concept using a mixed-methods cross-cultural approach. *Trop Med Int Health*. 2008 Dec;13(12):1534-42.

Beard JR, Dietrich UC, Brooks LO, Brooks RT, Heathcote K, Kelly B. Incidence and outcomes of mental disorders in a regional population: The northern rivers mental health study. *Aust NZ J Psychiatry*. 2006;40(8):674-82.

Berle JO, Aarre TF, Mykletun A, Dahl AA, Holsten F. Screening for postnatal depression. Validation of the Norwegian version of the Edinburgh Postnatal Depression Scale, and assessment of risk factors for postnatal depression. *J Affect Disord*. 2003 Sep;76(1-3):151-6.

Boersma K, Linton SJ. Screening to identify patients at risk: Profiles of psychological risk factors for early intervention. *Clin J Pain*. 1969;21(1):38-43.

Bowen A, Muhajarine N. Prevalence of antenatal depression in women enrolled in an outreach program in Canada. *JOGNN*. 2006 Jul;35(4):491-8.

Brooks J, Nathan E, Speelman C, Swalm D, Jacques A, Doherty D. Tailoring screening protocols for perinatal depression: Prevalence of high risk across obstetric services in Western Australia. *Arch Womens Ment Health*. 2009 Apr;12(2):105-12.

Buist A, Bilszta J, Barnett B, Milgrom J, Ericksen J, Condon J, et al. Recognition and management of perinatal depression in general practice--A survey of GPs and postnatal women. *Aust Fam Phys*. 2005 Sep;34(9):787-90.

Buist A, Condon J, Brooks J, Speelman C, Milgrom J, Hayes B, et al. Acceptability of routine screening for perinatal depression. *J Affect Disord*. 2006 Jul;93(1-3):233-7.

Buist A, Ellwood D, Brooks J, Milgrom J, Hayes BA, Sved-Williams A, et al. National program for depression associated with childbirth: The Australian experience. *Best Pract Res Clin Obstet Gynaecol*. 2007 Apr;21(2):193-206.

Carter FA, Carter JD, Luty SE, Wilson DA, Frampton CMA, Joyce PR. Screening and treatment for depression during pregnancy: A cautionary note. *Aust NZ J Psychiatry*. 2005 Apr;39(4):255-61.

Chaaya M, Campbell OM, El Kak F, Shaar D, Harb H, Kaddour A. Postpartum depression: Prevalence and determinants in Lebanon. *Arch Womens Ment Health*. 2002 Oct;5(2):65-72.

Chabrol H, Teissedre F, Saint-Jean M, Teisseyre N, Roge B, Mullet E. Prevention and treatment of postpartum depression: A controlled randomized study on women at risk. *Psychol Med*. 2002 Aug;32(6):1039-47.

Chabrol H, Teissedre F, Saint-Jean M, Teisseyre N, Roge B. Prevention and treatment of post partum depression: A controlled study. *Devenir*. 2003;15(1):5-25.

Chaudron LH, Szilagyi PG, Kitzman HJ, Wadkins HI, Conwell Y. Detection of postpartum depressive symptoms by screening at well-child visits. *Pediatrics*. 2004 Mar;113(3:Pt 1):551-8.

Chilcot J, Wellsted D, Farrington K. Screening for depression while patients dialyse: An evaluation. *Nephrol Dial Transplant*. 2008 Aug;23(8):2653-9.

Clemmens D, Driscoll JW, Beck CT. Postpartum depression as profiled through the depression screening scale. *Am J Matern Child Nurs*. 2004 May;29(3):180-5.

Coleman R, Morison L, Paine K, Powell RA, Walraven G. Women's reproductive health and depression: A community survey in the Gambia, West Africa. *Soc Psychiatry Psychiatr Epidemiol*. 2006 Sep;41(9):720-7.

Crandall M, Sridharan L, Schermer C. Depression and thoughts of death among disadvantaged mothers: Risk factors and impact on maternal and child health. *Arch Suicide Res*. 2010 Jul;14(3):248-60.

Crotty F, Sheehan J. Prevalence and detection of postnatal depression in an Irish community sample. *Ir J Psychol Med*. 2004;21(4):117-21.

Curro V, De Rosa E, Maulucci S, Maulucci ML, Silvestri MT, Zambrano A. The use of Edinburgh Postnatal Depression Scale to identify postnatal depression symptoms at well child visit. *Ital J Pediatr.* 2009;35(32).

Daly EJ, Trivedi MH, Raskin P, Grannemann BG. Screening for depression in a diabetic outpatient population. *Int J Psychiatry Clin Pract.* 2007;11(4):268-72.

Davidson S, Koritsas S, O'Connors DW, Clarke D. The feasibility of a GP led screening intervention for depression among nursing home residents. *Int J Geriatr Psychiatry.* 2006 Nov;21(11):1026-30.

Davies BR, Howells S, Jenkins M. Early detection and treatment of postnatal depression in primary care. *J Adv Nurs.* 2003;44(3):248-55.

Davis S, Cross J, Lind BK. Exploring the Postpartum Adjustment Questionnaire as a predictor of postpartum depression. *JOGNN.* 2008 Nov;37(6):622-30.

Dennis CL. Can we identify mothers at risk for postpartum depression in the immediate postpartum period using the Edinburgh Postnatal Depression Scale? *J Affect Disord.* 2004 Feb;78(2):163-9.

Dennis CL, Ross LE. La symptomatologie depressive dans la periode postnatale immediate: Identifier les caracteristiques maternelles reliees aux scores de depistage de vrai et de faux positif. *Can J Psychiatry.* 2006 Apr;51(5):265-73.

Dillard DA, Christopher D. The Southcentral Foundation depression collaborative. *Int J Circumpolar Health.* 2007;66:Suppl-53.

Dixon L, Myers P, Johnson J, Corty E. Screening for mental illness with the addiction severity index. *Am J Addict.* 1996;5(4):301-7.

Donlan W, Lee J. Screening for depression among indigenous Mexican migrant farmworkers using the Patient Health Questionnaire-9. *Psychol Rep.* 2010 Apr;106(2):419-32.

Downie J, Wynaden D, McGowan S, Juliff D, Axten C, Fitzpatrick L, et al. Using the Edinburgh Postnatal Depression Scale to achieve best practice standards. *Nursing & Health Sciences.* 2003 Dec;5(4):283-7.

Dugan W, McDonald MV, Passik SD, Rosenfeld BD, Theobald D, Edgerton S. Use of the Zung Self-Rating Depression Scale in cancer patients: Feasibility as a screening tool. *Psycho Oncol.* 1998 Nov;7(6):483-93.

Dworkin SF, Sherman J, Mancl L, Ohrbach R, LeResche L, Truelove E. Reliability, validity, and clinical utility of the research diagnostic criteria for Temporomandibular Disorders Axis II Scales: Depression, non-specific physical symptoms, and graded chronic pain. *J Orofacial Pain.* 2002;16(3):207-20.

Ell K, Vourlekis B, Nissly J, Padgett D, Pineda D, Sarabia O, et al. Integrating mental health screening and abnormal cancer screening follow-up: An intervention to reach low-income women. *Community Ment Health J.* 2002 Aug;38(4):311-25.

Evins GG, Theofrastous JP, Galvin SL. Postpartum depression: A comparison of screening and routine clinical evaluation. *Am J Obstet Gynecol.* 2000 May;182(5):1080-2.

Fann JR, Berry DL, Wolpin S, Austin-Seymour M, Bush N, Halpenny B, et al. Depression screening using the Patient Health Questionnaire-9 administered on a touch screen computer. *Psycho Oncol.* 2009;18(1):14-22.

Feinberg E, Smith MV, Naik R. Ethnically diverse mothers' views on the acceptability of screening for maternal depressive symptoms during pediatric well-child visits. *J Health Care Poor Underserv.* 2009 Aug;20(3):780-97.

Figueira P, Correa H, Malloy-Diniz L, Romano-Silva MA. Edinburgh Postnatal Depression Scale for screening in the public health system. *Rev Saude Publica.* 2009 Aug;43:Suppl-84.

Filipic I, Popovic-Grle S, Marcinko D, Basic S, Hotujac L, Pavicic F, et al. Screening for depression disorders in patients with chronic somatic illness. *Collegium Antropol.* 2007 Mar;31(1):139-43.

Flynn HA, O'Mahen HA, Massey L, Marcus S. The impact of a brief obstetrics clinic-based intervention on treatment use for perinatal depression. *J Womens Health.* 2006;15(10):1195-204.

Fossey L, Papiernik E, Bydlowski M. Postpartum blues: a clinical syndrome and predictor of postnatal depression? *J Psychosom Obstet Gynaecol.* 1997 Mar;18(1):17-21.

Foster MA, Ragsdale K, Dunne B, Jones E, Ihnen GH, Lentz C, et al. Detection and treatment of depression in a VA primary care clinic. *Psychiatr Serv.* 1999 Nov;50(11):1494-5.

Francoeur RB. A flexible item to screen for depression in inner-city minorities during palliative care symptom assessment. *Am J Geriatr Psychiatr.* 2006 Mar;14(3):228-36.

Franken IHA, Hendriks VM. Screening and diagnosis of anxiety and mood disorders in substance abuse patients. *Am J Addict.* 2001;10(1):30-9.

Fuggle P, Haydon K. Estimating resources for setting up an HV service for postnatal depression. *Br J Community Nurs.* 2000;5(7):348-51.

Fuggle P, Glover L, Khan F, Haydon K. Screening for postnatal depression in Bengali women: Preliminary observations from using a translated version of the Edinburgh Postnatal Depression Scale (EPDS). *J Reprod Infant Psychol.* 2002;20(2):71-82.

Gao LL, Chan SW, Mao Q. Depression, perceived stress, and social support among first-time Chinese mothers and fathers in the postpartum period. *Res Nurs Health.* 2009 Feb;32(1):50-8.

Georgiopoulos AM, Bryan TL, Wollan P, Yawn BP. Routine screening for postpartum depression. *J Fam Pract.* 2001 Feb;50(2):117-22.

Gilmer TP, Walker C, Johnson ED, Philis-Tsimikas A, Unutzer J. Improving treatment of depression among Latinos with diabetes using project Dulce and IMPACT. *Diabetes Care.* 2008 Jul;31(7):1324-6.

Giordano M, Tirelli P, Ciarambino T, Gambardella A, Ferrara N, Signoriello G, et al. Screening of depressive symptoms in young-old hemodialysis patients: Relationship between Beck Depression Inventory and 15-item Geriatric Depression Scale. *Nephron*. 2007;106(4):c187-c192.

Glavin K, Ellefsen B, Erdal B. Norwegian public health nurses' experience using a screening protocol for postpartum depression. *Public Health Nurs*. 2010 May;27(3):255-62.

Gordon TE, Cardone IA, Kim JJ, Gordon SM, Silver RK. Universal perinatal depression screening in an Academic Medical Center. *Obstetrics and Gynecology*. 2006 Feb;107(2:Pt 1):342-7.

Haggman S, Maher CG, Refshauge KM. Screening for symptoms of depression by physical therapists managing low back pain. *Phys Ther*. 2004 Dec;84(12):1157-66.

Hanna B, Jarman H, Savage S. The clinical application of three screening tools for recognizing postpartum depression. *Int J Nurs Pract*. 2004 Apr;10(2):72-9.

Hanna B, Jarman H, Savage S, Layton K. The early detection of postpartum depression: Midwives and nurses trial a checklist. *JOGNN*. 2004;33(2):191-7.

Harrington AR, Greene-Harrington CC. Healthy Start screens for depression among urban pregnant, postpartum and interconceptional women. *J Natl Med Assoc*. 2007 Mar;99(3):226-31.

Hatton DC, Harrison-Hohner J, Matarazzo J, Edwards P, Lewy A, Davis L. Missed antenatal depression among high risk women: A secondary analysis. *Arch Womens Ment Health*. 2007;10(3):121-3.

Hauenstein EJ, Peddada SD. Prevalence of major depressive episodes in rural women using primary care. *J Health Care Poor Underserv*. 2007 Feb;18(1):185-202.

Haver B. Screening for psychiatric comorbidity among female alcoholics: The use of a questionnaire (SCL-90) among women early in their treatment programme. *Alcohol Alcohol*. 1997;32(6):725-30.

Hayes DK, Ta VM, Hurwitz EL, Mitchell-Box KM, Fuddy LJ. Disparities in self-reported postpartum depression among Asian, Hawaiian, and Pacific Islander Women in Hawaii: Pregnancy Risk Assessment Monitoring System (PRAMS), 2004-2007. *Matern Child Health J*. 2010 Sep;14(5):765-73.

Hearn G, Iloff A, Jones I, Kirby A, Ormiston P, Parr P, et al. Postnatal depression in the community. *Br J Gen Pract*. 1998 Mar;48(428):1064-6.

Heller K, Viken RJ, Swindle RW, Jr. Screening for depression in African American and Caucasian older women. *Aging Ment Health*. 2010 Apr;14(3):339-46.

Honey KL, Bennett P, Morgan M. Predicting postnatal depression. *J Affect Disord*. 2003 Sep;76(1-3):201-10.

Hopko DR, Bell JL, Armento MEA, Robertson SMC, Hunt MK, Wolf NJ, et al. The phenomenology and screening of clinical depression in cancer patients. *J Psychosoc Oncol*. 2007;26(1):31-51.

Horowitz JA. Community-based postpartum depression screening within the first month after delivery. *Contemp Nurse*. 2006 Feb;21(1):85-93.

Horowitz JA, Murphy CA, Gregory KE, Wojcik J. Community-based postpartum depression screening: Results from the CARE study. *Psychiatr Serv*. 2009;60(11):1432-4.

Ingadottir E, Thome M. Evaluation of a web-based course for community nurses on postpartum emotional distress. *Scand J Caring Sci*. 2006;20(1):86-92.

Jacob S, Berry M, Kneebone A, Delaney G, Fowler A, Behan S, et al. Computerised screening for anxiety, depression and radiation toxicity in cancer patients. *Cancer Forum*. 2004;28(3):143-6.

Jevitt C, Zapata L, Harrington M, Berry E. Screening for perinatal depression with limited psychiatric resources. *J Am Psychiatr Nurs Assoc*. 2005 Dec;11(6):359-63.

Kaaya SF, Mbwambo JK, Kilonzo GP, Van Den Borne H, Leshabari MT, Fawzi MC, et al. Socio-economic and partner relationship factors associated with antenatal depressive morbidity among pregnant women in Dar es Salaam, Tanzania. *Tanzan J Health Res*. 2010 Jan;12(1):23-35.

Kahn LS, Fox CH, McIntyre RS, Tumiel-Berhalter L, Berdine DE, Lyle H. Assessing the prevalence of depression among individuals with diabetes in a medicaid managed-care program. *Int J Psychiatr Med*. 2008;38(1):13-29.

Kaminsky LM, Carlo J, Muench MV, Nath C, Harrigan JT, Canterino J. Screening for postpartum depression with the Edinburgh Postnatal Depression Scale in an indigent population: Does a directed interview improve detection rates compared with the standard self-completed questionnaire? *J Matern Fetal Neonatal Med*. 2008 May;21(5):321-5.

Katon WJ, Von Korff M, Lin EH, Simon G, Ludman E, Russo J, et al. The Pathways Study: A randomized trial of collaborative care in patients with diabetes and depression. *Arch Gen Psychiatry*. 2004 Oct;61(10):1042-9.

Katz KS, Blake SM, Milligan RA, Sharps PW, White DB, Rodan MF, et al. The design, implementation and acceptability of an integrated intervention to address multiple behavioral and psychosocial risk factors among pregnant African American women. *BMC Pregnancy Childbirth*. 2008;8(1):22.

Katz MR, Kopek N, Waldron J, Devins GM, Tomlinson G. Screening for depression in head and neck cancer. *Psycho Oncol*. 2004 Apr;13(4):269-80.

Kim H, Bracha Y, Tipnis A. Automated depression screening in disadvantaged pregnant women in an urban obstetric clinic. *Arch Womens Ment Health*. 2007;10(4):163-9.

Kim JJ, Gordon TE, La Porte LM, Adams M, Kuendig JM, Silver RK. The utility of maternal depression screening in the third trimester. *Am J Obstet Gynecol*. 2008 Nov;199(5):509-5.

Kirpinar I, Gozum S, Pasinlioglu T. Prospective study of postpartum depression in eastern Turkey prevalence, socio-demographic and obstetric correlates, prenatal anxiety and early awareness. *J Clin Nurs*. 2010 Feb;19(3-4):422-31.

Knight DE, Draeger RW, Heaton PC, Patel NC. Pharmacist screening for depression among patients with diabetes in an urban primary care setting. *JAPhA*. 2008 Jul;48(4):518-21.

Koenig HG, George LK, Peterson BL. Religiosity and remission of depression in medically ill older patients. *Am J Psychiatry*. 1998 Apr;155(4):536-42.

Kuosmanen L, Vuorilehto M, Kumpuniemi S, Melartin T. Post-natal depression screening and treatment in maternity and child health clinics. *J Psychiatr Ment Health Nurs*. 2010 Aug;17(6):554-7.

Lau Y, Wong DFK, Chan KS. The utility of screening for perinatal depression in the second trimester among Chinese: A three-wave prospective longitudinal study. *Arch Womens Ment Health*. 2010;13(2):153-64.

Lau Y, Wang Y, Yin L, Chan KS, Guo X. Validation of the Mainland Chinese version of the Edinburgh Postnatal Depression Scale in Chengdu mothers. *Int J Nurs Stud*. 2010 Sep;47(9):1139-51.

Le HN, Perry DF, Sheng X. Using the internet to screen for postpartum depression. *Matern Child Health J*. 2009 Mar;13(2):213-21.

Le HN, Perry DF, Ortiz G. The Postpartum Depression Screening Scale-Spanish version: Examining the psychometric properties and prevalence of risk for postpartum depression. *J Immigrant Minor Health*. 2010 Apr;12(2):249-58.

Lee DT, Yip AS, Chan SS, Tsui MH, Wong WS, Chung TK. Postdelivery screening for postpartum depression. *Psychosom Med*. 2003 May;65(3):357-61.

Leigh B, Milgrom J. Acceptability of antenatal screening for depression in routine antenatal care. *Aust J Adv Nurs*. 2007 Mar;24(3):14-8.

Liabsuetrakul T, Vittayanont A, Pitanupong J. Clinical applications of anxiety, social support, stressors, and self-esteem measured during pregnancy and postpartum for screening postpartum depression in Thai women. *J Obstet Gynaecol Res*. 2007 Jun;33(3):333-40.

Lincoln A, Feyerharm R, Damron P, DeVault M, Lorenz D, Dooley S. Maternal depression after delivery in Oklahoma. *J Oklahoma State Med Assoc*. 2008 Dec;101(12):307-11.

Lloyd-Williams M. Is it appropriate to screen palliative care patients for depression? *Am J Hospice Palliat Med*. 2002 Mar;19(2):112-4.

Lloyd-Williams M, Riddleston H. The stability of depression scores in patients who are receiving palliative care. *J Pain Symptom Manag*. 2002 Dec;24(6):593-7.

Luoma I, Tamminen T, Kaukonen P, Laippala P, Puura K, Salmelin R, et al. Longitudinal study of maternal depressive symptoms and child well-being. *J Am Acad Child Adolesc Psychiatr*. 2001 Dec;40(12):1367-74.

MacArthur C, Winter HR, Bick DE, Knowles H, Lilford R, Henderson C, et al. Effects of redesigned community postnatal care on womens' health 4 months after birth: A cluster randomised controlled trial. *Lancet*. 2002 Feb 2;359(9304):378-85.

Mancini F, Carlson C, Albers L. Use of the Postpartum Depression Screening Scale in a collaborative obstetric practice. *J Midwifery Womens Health*. 2007 Sep;52(5):429-34.

Mark TL, Fortner B, Johnson G. Evaluation of a tablet PC technology to screen and educate oncology patients. *Support Care Canc*. 2008;16(4):371-8.

Matsudaira T, Igarashi H, Kikuchi H, Kano R, Mitoma H, Ohuchi K, et al. Factor structure of the Hospital Anxiety and Depression Scale in Japanese psychiatric outpatient and student populations. *Health Qual Life Outcomes*. 2009;7:42.

Matthey S, Barnett B, Howie P, Kavanagh DJ. Diagnosing postpartum depression in mothers and fathers: Whatever happened to anxiety? *J Affect Disord*. 2003;74(2):139-47.

Matthey S, Phillips J, White T, Glossop P, Hopper U, Panasetis P, et al. Routine psychosocial assessment of women in the antenatal period: Frequency of risk factors and implications for clinical services. *Arch Womens Ment Health*. 2004 Oct;7(4):223-9.

Mauri M, Oppo A, Montagnani MS, Borri C, Banti S, Camilleri V, et al. Beyond "postpartum depressions": Specific anxiety diagnoses during pregnancy predict different outcomes: Results from PND-ReScU. *J Affect Disord*. 2010;127(1-3):177-84.

Mayberry LJ, Horowitz JA, Declercq E. Depression symptom prevalence and demographic risk factors among U.S. women during the first 2 years postpartum. *JOGNN*. 2007 Nov;36(6):542-9.

Meyer HA, Sinnott C, Seed PT. Depressive symptoms in advanced cancer. Part 1. Assessing depression: The Mood Evaluation Questionnaire. *Palliat Med*. 2003 Oct;17(7):596-603.

Milgrom J, Negri LM, Gemmill AW, McNeil M, Martin PR. A randomized controlled trial of psychological interventions for postnatal depression. *Br J Clin Psychol*. 2005 Nov;44(4):529-42.

Miller L, Shade M, Vasireddy V. Beyond screening: Assessment of perinatal depression in a perinatal care setting. *Arch Womens Ment Health*. 2009 Oct;12(5):329-34.

Miller LS, Boyd BJ, Chernov AJ. Improving the identification and treatment of postpartum depression in a managed care organization. *J Clin Outcome Manag*. 2004;11(3):157-61.

Mishina H, Hayashino Y, Takayama JI, Kasahara M, Fukuhara S. Can pediatricians accurately identify maternal depression at well-child visits? *Pediatr Int*. 2010 Apr;52(2):284-9.

Morris-Rush JK, Freda MC, Bernstein PS. Screening for postpartum depression in an inner-city population. *Am J Obstet Gynecol*. 2003 May;188(5):1217-9.

Mosack V, Shore ER. Screening for depression among pregnant and postpartum women. *J Community Health Nurs*. 2006;23(1):37-47.

Murray D, Cox JL, Chapman G, Jones P. Childbirth: life event or start of a long-term difficulty? Further data from the Stoke-on-Trent controlled study of postnatal depression. *Br J Psychiatry*. 1995 May;166(5):595-600.

Na YM, Kim KS, Lee KU, Chae JH, Kim JH, Kim DJ, et al. The relationship between depressive symptoms in outpatients with chronic illness and health care costs. *Yonsei Med J*. 2007 Oct 31;48(5):787-94.

Nakku JE, Nakasi G, Mirembe F. Postpartum major depression at six weeks in primary health care: Prevalence and associated factors. *Afr Health Sci*. 2006 Dec;6(4):207-14.

Navarro B, Andres F, Parraga I, Morena S, Latorre JM, Lopez-Torres J. Approach to major depression in old people. *Int Psychogeriatr*. 2010 Aug;22(5):733-8.

Neron S, Correa JA, Dajczman E, Kasymjanova G, Kreisman H, Small D. Screening for depressive symptoms in patients with unresectable lung cancer. *Support Care Canc*. 2007 Oct;15(10):1207-12.

Nguyen TN, Faulkner D, Allen S, Hauck YL, Frayne J, Rock D, et al. Managing pregnant women with serious mental illness: Using the Edinburgh Postnatal Depression Scale as a marker of anxiety and depressive symptoms. *Aust NZ J Psychiatry*. 2010;44(11):1036-42.

Nishizono-Maher A, Kishimoto J, Yoshida H, Urayama K, Miyato M, Otsuka Y, et al. The role of self-report questionnaire in the screening of postnatal depression- A community sample survey in central Tokyo. *Soc Psychiatry Psychiatr Epidemiol*. 2004 Mar;39(3):185-90.

O'Mahen HA, Flynn HA. Preferences and perceived barriers to treatment for depression during the perinatal period. *J Womens Health*. 2008 Oct;17(8):1301-9.

Olson AL, Dietrich AJ, Prazar G, Hurley J, Tuddenham A, Hedberg V, et al. Two approaches to maternal depression screening during well child visits. *J Dev Behav Pediatr*. 2005 Jun;26(3):169-76.

Olson AL, Dietrich AJ, Prazar G, Hurley J. Brief maternal depression screening at well-child visits. *Pediatrics*. 2006 Jul;118(1):207-16.

Oppo A, Mauri M, Ramacciotti D, Camilleri V, Banti S, Borri C, et al. Risk factors for postpartum depression: The role of the Postpartum Depression Predictors Inventory-Revised (PDPI-R). Results from the Perinatal Depression-Research & Screening Unit (PNDRScU) study. *Arch Womens Ment Health*. 2009 Aug;12(4):239-49.

Ostergaard SD, Foldager L, Allgulander C, Dahl AA, Huuhtanen MT, Rasmussen I, et al. Psychiatric caseness is a marker of major depressive episode in general practice. *Scand J Prim Health Care*. 2010 Dec;28(4):211-5.

Ouedraogo A, Kere M, Lankoande J, Pale K, Bonkian AE, Kone B. Screening of post-partum depressive states in the Yalgado Ouedraogo National Hospital Center maternity ward in Ouagadougou Burkina Faso. [French]. *J Gynecol Obstet Biol Reprod*. 1998;27(6):611-6.

Oyama H, Fujita M, Goto M, Shibuya H, Sakashita T. Outcomes of community-based screening for depression and suicide prevention among Japanese elders. *Gerontologist*. 2006 Dec;46(6):821-6.

Oyama H, Sakashita T, Hojo K, Ono Y, Watanabe N, Takizawa T, et al. A community-based survey and screening for depression in the elderly: The short-term effect on suicide risk in Japan. *Crisis*. 2010;31(2):100-8.

Pallant JF, Bailey CM. Assessment of the structure of the Hospital Anxiety and Depression Scale in musculoskeletal patients. *Health Qual Life Outcomes*. 2005;3:82.

Panthangi V, West P, Savoy-Moore RT, Geeta M, Reickert E. Is seasonal variation another risk factor for postpartum depression? *J Am Board Fam Med*. 2009 Sep;22(5):492-7.

Pasquini M, Biondi M, Costantini A, Cairoli F, Ferrarese G, Picardi A, et al. Detection and treatment of depressive and anxiety disorders among cancer patients: Feasibility and preliminary findings from a liaison service in an oncology division. *Depress Anxiety*. 2006;23(7):441-8.

Passik SD, Kirsh KL, Theobald D, Donaghy K, Holtsclaw E, Edgerton S, et al. Use of a depression screening tool and a fluoxetine-based algorithm to improve the recognition and treatment of depression in cancer patients. A demonstration project. *J Pain Symptom Manag*. 2002 Sep;24(3):318-27.

Patel D, Sharpe L, Thewes B, Rickard J, Schnieden V, Lewis C. Feasibility of using risk factors to screen for psychological disorder during routine breast care nurse consultations. *Canc Nurs*. 2010 Jan;33(1):19-27.

Payne DK, Hoffman RG, Theodoulou M, Dosik M, Massie MJ. Screening for anxiety and depression in women with breast cancer. *Psychiatry and medical oncology gear up for managed care. Psychosomatics*. 1999 Jan;40(1):64-9.

Pedersen SS, Denollet J, de Jonge P, Simsek C, Serruys PW, van Domburg RT. Brief depression screening with the PHQ-2 associated with prognosis following percutaneous coronary intervention with paclitaxel-eluting stenting. *J Gen Intern Med*. 2009;24(12):1359.

Pereira PK, Lovisi GM, Pilowsky DL, Lima LA, Legay LF. Depression during pregnancy: Prevalence and risk factors among women attending a public health clinic in Rio de Janeiro, Brazil. *Cadernos de Saude Publica*. 2009 Dec;25(12):2725-36.

Peyrot M, Rubin RR. Persistence of depressive symptoms in diabetic adults. *Diabetes Care*. 1999 Mar;22(3):448-52.

Pietrzak RH, Johnson DC, Goldstein MB, Malley JC, Rivers AJ, Morgan CA, et al. Psychosocial buffers of traumatic stress, depressive symptoms, and psychosocial difficulties in veterans of Operations Enduring Freedom and Iraqi Freedom: The role of resilience, unit support, and postdeployment social support. *J Spec Oper Med*. 2009;9(3):74-8.

Pitanupong J, Liabsuetrakul T, Vittayanont A. Validation of the Thai Edinburgh Postnatal Depression Scale for screening postpartum depression. *Psychiatry Res*. 2007 Jan 15;149(1-3):253-9.

Poinso F, Samuelian JC, Delzenne V, Huiart L, Sparrow J, Rufo M. Post-partum depressions: Delimitation of a high risk group immediately after birth: A prospective evaluation and mother-baby relations. *La Psychiatrie de l'Enfant*. 2001;44(2):379-413.

Pollock JI, Manaseki-Holland S, Patel V. Depression in Mongolian women over the first 2 months after childbirth: Prevalence and risk factors. *J Affect Disord*. 2009;116(1-2):126-33.

Poole H, White S, Blake C, Murphy P, Bramwell R. Depression in chronic pain patients: Prevalence and measurement. *Pain Pract*. 2009 May;9(3):173-80.

Ragland D, Payakachat N, Hays EB, Banken J, Dajani NK, Ott RE. Depression and diabetes: Establishing the pharmacist's role in detecting comorbidity in pregnant women. *JAPhA*. 2010 Mar 1;50(2):195-9.

Reck C, Struben K, Backenstrass M, Stefenelli U, Reinig K, Fuchs T, et al. Prevalence, onset and comorbidity of postpartum anxiety and depressive disorders. *Acta Psychiatrica Scandinavica*. 2008 Dec;118(6):459-68.

Reeve J, Lloyd-Williams M, Dowrick C. Depression in terminal illness: The need for primary care-specific research. *Fam Pract*. 2007 Jun;24(3):263-8.

Regmi S, Sligl W, Carter D, Grut W, Seear M. A controlled study of postpartum depression among Nepalese women: Validation of the Edinburgh Postpartum Depression Scale in Kathmandu. *Trop Med Int Health*. 2002;7(4):378-82.

Reme SE, Eriksen HR. Is one question enough to screen for depression? *Scand J Public Health*. 2010 Aug;38(6):618-24.

Rodgers J, Martin CR, Morse RC, Kendell K, Verrill M. An investigation into the psychometric properties of the Hospital Anxiety and Depression Scale in patients with breast cancer. *Health Qual Life Outcomes*. 2005;3:41.

Rychnovsky J, Beck CT. Screening for postpartum depression in military women with the Postpartum Depression Screening Scale. *Mil Med*. 2006 Nov;171(11):1100-4.

Schaper AM, Rooney BL, Kay NR, Silva PD. Use of the Edinburgh Postnatal Depression Scale to identify postpartum depression in a clinical setting. *J Reprod Med*. 1994 Aug;39(8):620-4.

Seehusen DA, Baldwin LM, Runkle GP, Clark G. Are family physicians appropriately screening for postpartum depression? *J Am Board Fam Pract.* 2005 Mar;18(2):104-12.

Segre LS, O'Hara MW, Arndt S, Beck CT. Screening and counseling for postpartum depression by nurses: The women's views. *Am J Matern Child Nurs.* 2010 Sep;35(5):280-5.

Shakespeare J, Blake F, Garcia J. A qualitative study of the acceptability of routine screening of postnatal women using the Edinburgh Postnatal Depression Scale. *Br J Gen Pract.* 2003 Aug;53(493):614-9.

Sharan H, Kaplan B, Weizer N, Sulkes J, Merlob P. Early screening of postpartum depression using the Edinburgh Postnatal Depression Scale. *Int J Risk Saf Med.* 2006;18(4):213-8.

Sharpe M, Strong V, Allen K, Rush R, Postma K, Tulloh A, et al. Major depression in outpatients attending a regional cancer centre: Screening and unmet treatment needs. *Br J Cancer.* 2004 Jan 26;90(2):314-20.

Sheeder J, Kabir K, Stafford B. Screening for postpartum depression at well-child visits: Is once enough during the first 6 months of life? *Pediatrics.* 2009 Jun;123(6):e982-e988.

Shemesh E, Annunziato RA, Rubinstein D, Sultan S, Malhotra J, Santra M, et al. Screening for depression and suicidality in patients with cardiovascular illnesses. *Am J Cardiol.* 2009 Nov 1;104(9):1194-7.

Shimizu K, Akechi T, Okamura M, Oba A, Fujimori M, Akizuki N, et al. Usefulness of the nurse-assisted screening and psychiatric referral program. *Cancer.* 2005;103(9):1949-56.

Shimizu K, Akizuki N, Akechi T, Okamura M, Oba A, Shimamoto M, et al. Clinical experience of the modified nurse-assisted screening and psychiatric referral program. *Palliat Support Care.* 2008 Mar;6(1):29-32.

Sit DKY, Flint C, Svidergol D, White J, Wimer M, Bish B, et al. An emerging best practice model for perinatal depression care. *Psychiatr Serv.* 2009;60(11):1429-31.

Smith MV, Rosenheck RA, Cavaleri MA, Howell HB, Poschman K, Yonkers KA. Screening for and detection of depression, panic disorder, and PTSD in public-sector obstetric clinics. *Psychiatr Serv.* 2004 Apr;55(4):407-14.

Stamp GE, Williams AS, Crowther CA. Predicting postnatal depression among pregnant women. *Birth.* 1996 Dec;23(4):218-23.

Stowe ZN, Hostetter AL, Newport DJ. The onset of postpartum depression: Implications for clinical screening in obstetrical and primary care. *Am J Obstet Gynecol.* 2005 Feb;192(2):522-6.

Subramanian DN, Hopayian K. An audit of the first year of screening for depression in patients with diabetes and ischaemic heart disease under the Quality and Outcomes Framework. *Qual Prim Care.* 2008;16(5):341-4.

Sulaiman N, Hamdan A, Tamim H, Mahmood DA, Young D. The prevalence and correlates of depression and anxiety in a sample of diabetic patients in Sharjah, United Arab Emirates. *BMC Fam Pract*. 2010;11:80.

Tam LW, Newton RP, Dern M, Parry BL. Screening women for postpartum depression at well baby visits: Resistance encountered and recommendations. *Arch Womens Ment Health*. 2002 Oct;5(2):79-82.

Teissedre F, Chabrol H. A study of the Edinburgh Postnatal Depression Scale (EPDS) on 859 mothers: Detection of mothers at risk for postpartum depression. *Encephale*. 2004;30(4):376-81.

Teissedre F, Chabrol H. Screening, prevention and postpartum treatment: A randomized comparative study on 450 women. *Neuropsychiatrie de l'Enfance et de l'Adolescence*. 2004;52(5):266-73.

Teng HW, Hsu CS, Shih SM, Lu ML, Pan JJ, Shen WW. Screening postpartum depression with the Taiwanese version of the Edinburgh Postnatal Depression scale. *Compr Psychiatr*. 2005 Jul;46(4):261-5.

Tuohy A, McVey C. Subscales measuring symptoms of non-specific depression, anhedonia, and anxiety in the Edinburgh Postnatal Depression Scale. *Br J Clin Psychol*. 2008 Jun;47(Pt:2):2-69.

Turvey CL, Willyard D, Hickman DH, Klein DM, Kukoyi O. Telehealth screen for depression in a chronic illness care management program. *Telemed J E Health*. 2007 Feb;13(1):51-6.

Ugarriza DN. Screening for postpartum depression. *J Psychosoc Nurs Ment Health Serv*. 2000 Dec;38(12):44-51.

Uwakwe R, Okonkwo JE. Affective (depressive) morbidity in puerperal Nigerian women: Validation of the Edinburgh Postnatal Depression Scale. *Acta Psychiatrica Scandinavica*. 2003 Apr;107(4):251-9.

van Steenberg-Weijnenburg KM, de VL, Ploeger RR, Brals JW, Vloedveld MG, Veneman TF, et al. Validation of the PHQ-9 as a screening instrument for depression in diabetes patients in specialized outpatient clinics. *BMC Health Serv Res*. 2010;10:235.

Verkerk GJ, Pop VJ, Van Son MJ, Van Heck GL. Prediction of depression in the postpartum period: A longitudinal follow-up study in high-risk and low-risk women. *J Affect Disord*. 2003 Nov;77(2):159-66.

Walker J, Postma K, McHugh GS, Rush R, Coyle B, Strong V, et al. Performance of the Hospital Anxiety and Depression Scale as a screening tool for major depressive disorder in cancer patients. *J Psychosom Res*. 2007 Jul;63(1):83-91.

Walsh RM, Abbott JH. Screening for depression in a musculoskeletal outpatient physical therapy clinic: Point prevalence and comparison of two instruments. *J Back Musculoskel Rehabil*. 2008;21(3):171-4.

Watt S, Sword W, Krueger P, Sheehan D, Ontario Mother & Infant Survey. A cross-sectional study of early identification of postpartum depression: Implications for primary care providers from The Ontario Mother & Infant Survey. *BMC Fam Pract*. 2002 Apr 11;3:5.

Webster J, Linnane J, Dibley L, Starrenburg S, Roberts J, Hinson J. The impact of screening for risk factors associated with postnatal depression at the first prenatal visit. *J Qual Clin Pract.* 1997 Jun;17(2):65-71.

Webster J, Linnane JW, Dibley LM, Pritchard M. Improving antenatal recognition of women at risk for postnatal depression. *Aust NZ J Obstet Gynaecol.* 2000 Nov;40(4):409-12.

Webster J, Linnane J, Roberts J, Starrenburg S, Hinson J, Dibley L. IDentify, Educate and Alert (IDEA) trial: An intervention to reduce postnatal depression. *Int J Obstet Gynaecol.* 2003 Sep;110(9):842-6.

Wickberg B, Hwang CP. Counselling of postnatal depression: A controlled study on a population based Swedish sample. *J Affect Disord.* 1996;39(3):209-16.

Wittkampf K, van Ravesteijn H, Baas K, van de Hoogen H, Schene A, Bindels P, et al. The accuracy of Patient Health Questionnaire-9 in detecting depression and measuring depression severity in high-risk groups in primary care. *Gen Hosp Psychiatry.* 2009 Sep;31(5):451-9.

Wojnar M, Drozd W, Araszkievicz A, Szymanski W, Nawacka-Pawlaczyk D, Urbanski R, et al. Assessment and prevalence of depression in women 45-55 years of age visiting gynecological clinics in Poland: Screening for depression among midlife gynecologic patients. *Arch Womens Ment Health.* 2003;6(3):193-201.

Yamashita H, Yoshida K, Nakano H, Tashiro N. Postnatal depression in Japanese women. Detecting the early onset of postnatal depression by closely monitoring the postpartum mood. *J Affect Disord.* 2000 May;58(2):145-54.

Yonkers KA, Smith MV, Lin H, Howell HB, Shao L, Rosenheck RA. Depression screening of perinatal women: An evaluation of the healthy start depression initiative. *Psychiatr Serv.* 2009 Mar;60(3):322-8.

Yonkers KA, Smith MV, Gotman N, Belanger K. Typical somatic symptoms of pregnancy and their impact on a diagnosis of major depressive disorder. *Gen Hosp Psychiatry.* 2009 Jul;31(4):327-33.

Yoshida K, Marks MN, Kibe N, Kumar R, Nakano H, Tashiro N. Postnatal depression in Japanese women who have given birth in England. *J Affect Disord.* 1997 Mar;43(1):69-77.

Zubaran C, Foresti K, Schumacher MV, Amoretti AL, Muller LC, Thorell MR, et al. Validation of a screening instrument for postpartum depression in Southern Brazil. *J Psychosom Obstet Gynaecol.* 2009 Dec;30(4):244-54.

Zubaran C, Foresti K, Schumacher MV, Amoretti AL, Thorell MR, Muller LC. The correlation between postpartum depression and health status. *Matern Child Health J.* 2010 Sep;14(5):751-7.

Excluded: No comparators

Angst J, Gamma A, Sellaro R, Zhang H, Merikangas K. Toward validation of atypical depression in the community: Results of the Zurich cohort study. *J Affect Disord.* 2002 Nov;72(2):125-38.

Aragones E, Pinol JL, Labad A, Folch S, Melich N. Detection and management of depressive disorders in primary care in Spain. *Int J Psychiatr Med.* 2004;34(4):331-43.

Baer L, Jacobs DG, Cukor P, O'Laughlen J, Coyle JT, Magruder KM. Automated telephone screening survey for depression. *JAMA.* 1995;273(24):1943-4.

Baiyewu O, Smith-Gamble V, Lane KA, Gureje O, Gao S, Ogunniyi A, et al. Prevalence estimates of depression in elderly community-dwelling African Americans in Indianapolis and Yoruba in Ibadan, Nigeria. *Int Psychogeriatr.* 2007 Aug;19(4):679-89.

Barkow K, Heun R, Ustun TB, Berger M, Bermejo I, Gaebel W, et al. Identification of somatic and anxiety symptoms which contribute to the detection of depression in primary health care. *Eur Psychiatry.* 2004 Aug;19(5):250-7.

Beausang P, Syeed R. Screening for anxiety and depression in adult general medical in-patients in a Scottish District General Hospital. *Scottish Med J.* 1998 Dec;43(6):177-80.

Becker SM. Detection of somatization and depression in primary care in Saudi Arabia. *Soc Psychiatry Psychiatr Epidemiol.* 2004 Dec;39(12):962-6.

Bellantuono C, Mazzi MA, Tansella M, Rizzo R, Goldberg D. The identification of depression and the coverage of antidepressant drug prescriptions in Italian general practice. *J Affect Disord.* 2002 Oct;72(1):53-9.

Bermejo I, Niebling W, Berger M, Harter M. Patients' and physicians' evaluation of the PHQ-D for depression screening. *Prim Care Community Psychiatr.* 2005;10(4):125-31.

Bijl D, Van Marwijk HWJ, Beekman ATF, de Haan M, van Tilburg W. A randomized controlled trial to improve the recognition, diagnosis and treatment of major depression in elderly people in general practice: Design, first results and feasibility of the West Friesland Study. *Prim Care Psychiatr.* 2003;8(4):135-40.

Blanchard MR, Waterreus A, Mann AH. The nature of depression among older people in inner London, and the contact with primary care. *Br J Psychiatry.* 1994 Mar;164(3):396-402.

Cameron IM, Lawton K, Reid IC. Appropriateness of antidepressant prescribing: An observational study in a Scottish primary-care setting. *Br J Gen Pract.* 2009;59(566):644-9.

Christensen KS, Sokolowski I, Olesen F. Case-finding and risk-group screening for depression in primary care. *Scand J Prim Health Care.* 2011 Jun;29(2):80-4.

Chung H, Teresi J, Guarnaccia P, Meyers BS, Holmes D, Bobrowitz T, et al. Depressive symptoms and psychiatric distress in low income Asian and Latino primary care patients: Prevalence and recognition. *Community Ment Health J.* 2003 Feb;39(1):33-46.

Costa E, Barreto SM, Uchoa E, Lima-Costa MFF. Agreement between the Geriatric Depression Scale and the General Health Questionnaire in a population-based elderly cohort: The Bambui health & ageing study (BHAS). *Clin Gerontol.* 2003;26(3-4):69-82.

Diminic-Lisica I, Franiskovic T, Baricev-Novakovic Z, Bukmir L, Grkovic J, Moro L. The prevalence and detection of depressive disorders in a Croatian primary care setting. *Prim Care Community Psychiatr.* 2005;10(1):19-25.

Doetch TM, Alger BH, Glasser M, Levenstein J. Detecting depression in elderly outpatients: Findings from depression symptom scales and the Dartmouth COOP charts. *Fam Med.* 1994 Sep;26(8):519-23.

Donnelly PL, Kim KS. The Patient Health Questionnaire (PHQ-9K) to screen for depressive disorders among immigrant Korean American elderly. *J Cult Divers.* 2008;15(1):24-9.

Dumont P, Andreoli A, Borgacci S, Carballeira Y, Rentsch D, de Tonnac N, et al. Quick detection of depression: A significant clinical issue. *Revue Medicale Suisse.* 2005;1(5):344-6.

Edwards JB, Tudiver F. Women's preventive screening in rural health clinics. *Womens Health Iss.* 2008;18(3):155-66.

Ell K, Unutzer J, Aranda M, Sanchez K, Lee PJ. Routine PHQ-9 depression screening in home health care: Depression, prevalence, clinical and treatment characteristics and screening implementation. *Home Health Care Serv Q.* 2005;24(4):1-19.

Eskelinen M, Ollonen P. Beck Depression Inventory (BDI) in patients with breast disease and breast cancer: A prospective case-control study. *In Vivo.* 2011 Jan;25(1):111-6.

Fechner-Bates S, Coyne JC, Schwenk TL. The relationship of self-reported distress to depressive disorders and other psychopathology. *J Consult Clin Psychol.* 1994 Jun;62(3):550-9.

Freeman MP, Wright R, Watchman M, Wahl RA, Sisk DJ, Fraleigh L, et al. Postpartum depression assessments at well-baby visits: Screening feasibility, prevalence, and risk factors. *J Womens Health.* 2005 Dec;14(10):929-35.

Freudenstein U, Arthur A, Matthews R, Jagger C. Can routine information improve case finding of depression among 65 to 74 year olds in primary care? *Fam Pract.* 2002 Oct;19(5):520-2.

Gallo JJ, Bogner HR, Straton JB, Margo K, Lesho P, Rabins PV, et al. Patient characteristics associated with participation in a practice-based study of depression in late life: The Spectrum study. *Int J Psychiatr Med.* 2005;35(1):41-57.

Garlow SJ, Rosenberg J, Moore JD, Haas AP, Koestner B, Hendin H, et al. Depression, desperation, and suicidal ideation in college students: Results from the American Foundation for Suicide Prevention College Screening Project at Emory University. *Depress Anxiety*. 2008;25(6):482-8.

Gega L, Kenwright M, Mataix-Cols D, Cameron R, Marks IM. Screening people with anxiety/depression for suitability for guided self-help. *Cognit Behav Ther*. 2005;34(1):16-21.

Gellis ZD. Depression screening in medically ill homecare elderly. *Best Pract Ment Health*. 2010 Jan;6(1):1-16.

Gonzalez GM, Costello CR, La Tourette TR, Joyce LK, Valenzuela M. Bilingual telephone-assisted computerized speech-recognition assessment: Is a voice-activated computer program a culturally and linguistically appropriate tool for screening depression in English and Spanish? *Cult Divers Ment Health*. 1997;3(2):93-111.

Gottfries GG, Noltorp S, Norgaard N. Experience with a Swedish version of the Geriatric Depression Scale in primary care centres. *Int J Geriatr Psychiatry*. 1997 Oct;12(10):1029-34.

Hamid WA, Wykes T, Stansfeld S. The social disablement of men in hostels for homeless people. I. Reliability and prevalence. *Br J Psychiatry*. 1995;166(JUNE):806-8.

Hammond MF. Doctors' and nurses' observations on the Geriatric Depression Rating Scale. *Age Ageing*. 2004 Mar;33(2):189-92.

Heisel MJ, Flett GL, Duberstein PR, Lyness JM. Does the geriatric depression scale (GDS) distinguish between older adults with high versus low levels of suicidal ideation? *Am J Geriatr Psychiatry*. 2005 Oct;13(10):876-83.

Hirshfield S, Wolitski RJ, Chiasson MA, Remien RH, Humberstone M, Wong T. Screening for depressive symptoms in an online sample of men who have sex with men. *AIDS Care*. 2008 Sep;20(8):904-10.

Houston TK, Cooper LA, Vu HT, Kahn J, Toser J, Ford DE. Screening the public for depression through the Internet. *Psychiatr Serv*. 2001 Mar;52(3):362-7.

Janssen J, Beekman AT, Comijs HC, Deeg DJ, Heeren TJ. Late-life depression: The differences between early- and late-onset illness in a community-based sample. *Int J Geriatr Psychiatry*. 2006 Jan;21(1):86-93.

Johanson R, Chapman G, Murray D, Johnson I, Cox J. The North Staffordshire Maternity Hospital prospective study of pregnancy-associated depression. *J Psychosom Obstet Gynaecol*. 2000 Jun;21(2):93-7.

Johnston L, Reid A, Wilson J, Levesque J, Driver B. Detecting depression in the aged: Is there concordance between screening tools and the perceptions of nursing home staff and residents? A pilot study in a rural aged care facility. *Aust J Rural Health*. 2007 Aug;15(4):252-6.

Kasantikul D, Karnjanathanalers N, Limsuwan N, Thongtang O, Vuthiganond S, Khuangsirikul V, et al. Health-related self-report (HRSR) scale: The diagnostic screening test for depression in Thai population. *J Med Assoc Thailand*. 1997 Oct;80(10):647-57.

Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003 Jun 18;289(23):3095-105.

King M, Walker C, Levy G, Bottomley C, Royston P, Weich S, et al. Development and validation of an international risk prediction algorithm for episodes of major depression in general practice attendees: The predictD study. *Arch Gen Psychiatry*. 2008;65(12):1368-76.

Knox ED, Dopheide JA, Wincor MZ, Han PK. Depression screening in a university campus pharmacy: A pilot project. *JAPhA*. 2006 Jul;46(4):502-6.

Koponen H, Jokelainen J, Keinanen-Kiukaanniemi S, Vanhala M. Depressive symptoms and 10-year risk for cardiovascular morbidity and mortality. *World J Biol Psychiatr*. 2010 Sep;11(6):834-9.

Lagerberg D, Magnusson M, Sundelin C. Drawing the line in the Edinburgh Postnatal Depression Scale (EPDS): A vital decision. *Int J Adolesc Med Health*. 2011;23(1):27-32.

Lagrue G, Dupont P, Fakhfakh R. Anxiety and depressive disorders in tobacco dependence. *Encephale*. 2002;28(4):374-7.

Lamoureux BE, Linardatos E, Fresco DM, Bartko D, Logue E, Milo L. Using the QIDS-SR16 to identify major depressive disorder in primary care medical patients. *Behav Ther*. 2010 Sep;41(3):423-31.

Lee AG, Beaver HA, Jogerst G, Daly JM. Screening elderly patients in an outpatient ophthalmology clinic for dementia, depression, and functional impairment. *Ophthalmology*. 2007;114(4):651-7.

Leung SS, Leung C, Lam TH, Hung SF, Chan R, Yeung T, et al. Outcome of a postnatal depression screening programme using the Edinburgh Postnatal Depression Scale: A randomized controlled trial. *J Public Health*. 2011 Jun;33(2):292-301.

Levkoff SE, Chen H, Coakley E, McDonel Herr EC, Oslin DW, Katz I, et al. Design and sample characteristics of the PRISM-E multisite randomized trial to improve behavioral health care for the elderly. *J Aging Health*. 2004;16(1):3-27.

Lewis G, Sharp D, Bartholomew J, et al. Computerized assessment of common mental disorders in primary care: effect on clinical outcome. *Fam Pract* 1996;13:120-6

Licht-Strunk E, Beekman AT, de Haan M, van Marwijk HW. The prognosis of undetected depression in older general practice patients. A one year follow-up study. *J Affect Disord*. 2009 Apr;114(1-3):310-5.

Lotfi L, Flyckt L, Krakau I, Martensson B, Nilsson GH. Undetected depression in primary healthcare: Occurrence, severity and co-morbidity in a two-stage procedure of opportunistic screening. *Nordic Journal of Psychiatry*. 2010;64(6):421-7.

Lowe B, Kroenke K, Grafe K. Detecting and monitoring depression with a two-item questionnaire (PHQ-2). *J Psychosom Res*. 2005 Feb;58(2):163-71.

Lubetkin EI, Jia H, Gold MR. Depression, anxiety, and associated health status in low-income Chinese patients. *Am J Prev Med*. 2003 May;24(4):354-60.

Lutz W, Stahl SM, Howard KI, Grissom GR, Joske R. Some relationships among assessments of depression. *J Clin Psychol*. 2002;58(12):1545-53.

Magruder KM, Norquist GS, Feil MB, Kopans B, Jacobs D. Who comes to a voluntary depression screening program? *Am J Psychiatry*. 1995 Nov;152(11):1615-22.

Mancuso CA, Westermann H, Choi TN, Wenderoth S, Briggs WM, Charlson ME. Psychological and somatic symptoms in screening for depression in asthma patients. *J Asthma*. 2008 Apr;45(3):221-5.

Marcus SM, Flynn HA, Blow FC, Barry KL. Depressive symptoms among pregnant women screened in obstetrics settings. *J Womens Health*. 2003 May;12(4):373-80.

Maruyama A. Mental health problems in primary care: In the context of general health service in Japan. *Japan Med Assoc J*. 2006;49(1):3-13.

McCrea D, Arnold E, Marchevsky D, Kaufman BM. The prevalence of depression in geriatric medical outpatients. *Age Ageing*. 1994 Nov;23(6):465-7.

McCusker J, Cole M, Latimer E, Sewitch M, Ciampi A, Cepoiu M, et al. Recognition of depression in older medical inpatients discharged to ambulatory care settings: A longitudinal study. *Gen Hosp Psychiatry*. 2008 May;30(3):245-51.

McGrady A, Lynch DJ, Nagel RW, Tamburrino M. Coherence between physician diagnosis and patient self reports of anxiety and depression in primary care. *J Nervous Ment Dis*. 2010;198(6):420-4.

McQuaid JR, Stein MB, McCahill M, Laffaye C, Ramel W. Use of brief psychiatric screening measures in a primary care sample. *Depress Anxiety*. 2000;12(1):21-9.

Mergl R, Seidscheck I, Allgaier AK, Moller HJ, Hegerl U, Henkel V. Depressive, anxiety, and somatoform disorders in primary care: Prevalence and recognition. *Depress Anxiety*. 2007;24(3):185-95.

Mitchell AM, Mittelstaedt ME, Schott-Baer D. Postpartum depression: The reliability of telephone screening. *Am J Matern Child Nurs*. 2006 Nov;31(6):382-7.

Morrow-Howell N, Proctor E, Choi S, Lawrence L, Brooks A, Hasche L, et al. Depression in public community long-term care: Implications for intervention development. *J Behav Health Serv Res*. 2008 Jan;35(1):37-51.

Moutier C, Norcross W, Jong P, Norman M, Kirby B, McGuire T, et al. The suicide prevention and depression awareness program at the University of California, San Diego School of Medicine. *Acad Med.* 2012 Mar;87(3):320-6.

Muhwezi WW, Agren H, Musisi S. Detection of major depression in Ugandan primary health care settings using simple questions from a subjective well-being (SWB) subscale. *Soc Psychiatry Psychiatr Epidemiol.* 2007 Jan;42(1):61-9.

Mullan E, Katona P, D'Ath P, Katona C. Screening, detection and management of depression in elderly primary care attenders. II: Detection and fitness for treatment: a case record study. *Fam Pract.* 1994 Sep;11(3):267-70.

Mumford DB, Nazir M, Jilani FU, Baig IY. Stress and psychiatric disorder in the Hindu Kush: A community survey of mountain villages in Chitral, Pakistan. *Br J Psychiatry.* 1996 Mar;168(3):299-307.

Munoz RF, McQuaid JR, Gonzalez GM, Dimas J, Rosales VA. Depression screening in a women's clinic: Using automated Spanish- and English-language voice recognition. *J Consult Clin Psychol.* 1999 Aug;67(4):502-10.

Nadeem E, Lange JM, Edge D, Fongwa M, Belin T, Miranda J. Does stigma keep poor young immigrant and U.S.-born black and Latina women from seeking mental health care? *Psychiatr Serv.* 2007;58(12):1547-54.

Nakao M, Yano E. Somatic symptoms for predicting depression: One-year follow-up study in annual health examinations. *Psychiatry Clin Neurosci.* 2006 Apr;60(2):219-25.

Neal RM, Baldwin RC. Screening for anxiety and depression in elderly medical outpatients. *Age Ageing.* 1994 Nov;23(6):461-4.

Nease DE, Klinkman MS, Volk RJ. Improved detection of depression in primary care through severity evaluation. *J Fam Pract.* 2002 Dec;51(12):1065-70.

Nguyen NL, Hunt DD, Scott CS. Screening for depression in a primary care setting in Vietnam. *J Nervous Ment Dis.* 2005 Feb;193(2):144-7.

Nuevo R, Dunn G, Dowrick C, Vazquez-Barquero JL, Casey P, Dalgard OS, et al. Cross-cultural equivalence of the Beck Depression Inventory: A five-country analysis from the ODIN study. *J Affect Disord.* 2009 Apr;114(1-3):156-62.

Nunn R, Annells M, Sims J. Screening for depression in older district nursing clients: Feasibility study. *Br J Community Nurs.* 2007;12(3):108-14.

O'Connor DW, Parslow RA. Differences in older people's responses to CIDI's depression screening and diagnostic questions may point to age-related bias. *J Affect Disord.* 2010 Sep;125(1-3):361-4.

Ongel K, Mergen H, Tan S, Kisioglu AN, Katirci E, Cadirci D, et al. Psychometric properties of depression in university students of Turkey. *Biomed Res.* 2010;21(3):265-71.

Ormel J, VonKorff M, Oldehinkel AJ, Simon G, Tiemens BG, Ustun TB. Onset of disability in depressed and non-depressed primary care patients. *Psychol Med.* 1999 Jul;29(4):847-53.

Parashos IA, Stamouli S, Rogakou E, Theodotou R, Nikas I, Mougias A. Recognition of depressive symptoms in the elderly: What can help the patient and the doctor. *Depress Anxiety.* 2002;15(3):111-6.

Patten SB, Schopflocher D. Longitudinal epidemiology of major depression as assessed by the Brief Patient Health Questionnaire (PHQ-9). *Compr Psychiatr.* 2009 Jan;50(1):26-33.

Pelissolo A, Maniere F, Boutges B, Allouche M, Richard-Berthe C, Corruble E. Anxiety and depressive disorders in 4,425 long term benzodiazepine users in general practice. *Encephale.* 2007 Jan;33(1):32-8.

Pepersack T. Minimum geriatric screening tools to detect common geriatric problems. *J Nutr Health Aging.* 2008;12(5):348-52.

Posse M, Hallstrom T. Depressive disorders among somatizing patients in primary health care. *Acta Psychiatrica Scandinavica.* 1998 Sep;98(3):187-92.

Pouget R, Yersin B, Wietlisbach V, Bumand B, Bula CJ. Depressed mood in a cohort of elderly medical inpatients: Prevalence, clinical correlates and recognition rate. *Aging Clin Exp Res.* 2000 Aug;12(4):301-7.

Poutanen O, Koivisto AM, Joukamaa M, Mattila A, Salokangas RK. The Depression Scale as a screening instrument for a subsequent depressive episode in primary healthcare patients. *Br J Psychiatry.* 2007 Jul;191:50-4.

Poutanen O, Mattila A, Seppala NH, Groth L, Koivisto AM, Salokangas RK. Seven-year outcome of depression in primary and psychiatric outpatient care: Results of the TADEP (Tampere Depression) II Study. *Nordic Journal of Psychiatry.* 2007;61(1):62-70.

Poutanen O, Koivisto AM, Mattila A, Joukamaa M, Salokangas RK. Gender differences in the symptoms of major depression and in the level of social functioning in public primary care patients. *Eur J Gen Pract.* 2009;15(3):161-7.

Prakash O, Gupta LN, Singh VB, Nagarajarao G. Applicability of 15-item Geriatric Depression Scale to detect depression in elderly medical outpatients. *Asian J Psychiatr.* 2009;2(2):63-5.

Rapaport MH, Suminski M. National Depression Screening Day: The San Diego experience. *Hosp Community Psychiatr.* 1994 Oct;45(10):1042-3.

Rigler SK, Belt-Marchesi J. Depression screening in congregate meal sites for older adults. *Clin Gerontol.* 2005;28(3):71-80.

Rogers WH, Adler DA, Bungay KM, Wilson IB. Depression screening instruments made good severity measures in a cross-sectional analysis. *J Clin Epidemiol.* 2005 Apr;58(4):370-7.

Rojahn J, Warren VJ, Ohringer S. A comparison of assessment methods for depression in mental retardation. *J Autism Dev Disord.* 1994 Jun;24(3):305-13.

Rouch-Leroyer I, Sourgen C, Barberger-Gateau P, Fuhrer R, Dartigues JF. Detection of depressive symptomatology in elderly people: A short version of the CES-D scale. *Aging Clin Exp Res.* 2000 Jun;12(3):228-33.

Rovner BW, Shmueli-Dulitzki Y. Screening for depression in low-vision elderly. *Int J Geriatr Psychiatry.* 1997 Sep;12(9):955-9.

Runkewitz K, Kirchmann H, Strauss B. Anxiety and depression in primary care patients: Predictors of symptom severity and developmental correlates. *J Psychosom Res.* 2006;60(5):445-53.

Sanchez-Villegas A, Schlatter J, Ortuno F, Lahortiga F, Pla J, Benito S, et al. Validity of a self-reported diagnosis of depression among participants in a cohort study using the Structured Clinical Interview for DSM-IV (SCID-I). *BMC Psychiatry.* 2008;8:43.

Sartorius N, Ustun TB, Lecrubier Y, Wittchen HU. Depression comorbid with anxiety: Results from the WHO study on psychological disorders in primary health care. *Br J Psychiatry.* 1996 Jun;(30):38-43.

Schnall R, Currie LM, Jia H, John RM, Lee NJ, Velez O, et al. Predictors of depression screening rates of nurses receiving a personal digital assistant-based reminder to screen. *J Urban Health.* 2010 Jul;87(4):703-12.

Sears SFJ, Danda CE, Evans GD. PRIME-MD and rural primary care: Detecting depression in a low-income rural population. *Prof Psychol Res Pract.* 1999 Aug;30(4):357-60.

Segre LS, O'Hara MW, Brock RL, Taylor D. Depression screening of perinatal women by the des moines healthy start project: Program description and evaluation. *Psychiatr Serv.* 2012;63(3):250-5.

Shea TL, Tennant A, Pallant JF. Rasch model analysis of the Depression, Anxiety and Stress Scales (DASS). *BMC Psychiatry.* 2009;9:21.

Sherina MS, Rampal L, Kaneson N. Psychological stress among undergraduate medical students. *Med J Malaysia.* 2004 Jun;59(2):207-11.

Stecker T, Fortney JC, Sherbourne CD. An intervention to increase mental health treatment engagement among OIF veterans: A pilot trial. *Mil Med.* 2011 Jun;176(6):613-9.

Swartz RJ, de Moor C, Cook KF, Fouladi RT, Basen-Engquist K, Eng C, et al. Mode effects in the center for epidemiologic studies depression (CES-D) scale: Personal digital assistant vs. paper and pencil administration. *Qual Life Res.* 2007 Jun;16(5):803-13.

Taylor JK, Schoenbaum M, Katon WJ, Pincus HA, Hogan DM, Unutzer J. Strategies for identifying and channeling patients for depression care management. *Am J Manag Care.* 2008 Aug;14(8):497-504.

Terluin B, Van Hout HPJ, Van Marwijk HWJ, Ader HJ, van der Meer K, de Haan M, et al. Reliability and validity of the assessment of depression in general practice: The Short Depression Interview. *Gen Hosp Psychiatry*. 2002 Nov;24(6):396-401.

Thompson C, Ostler K, Peveler RC, Baker N, Kinmonth AL. Dimensional perspective on the recognition of depressive symptoms in primary care: The Hampshire Depression Project 3. *Br J Psychiatry*. 2001 Oct;179:317-23.

Tiemens BG, Ormel J, Simon GE. Occurrence, recognition, and outcome of psychological disorders in primary care. *Am J Psychiatry*. 1996 May;153(5):636-44.

van den Berg M, Smit F, Vos T, van Baal PH. Cost-effectiveness of opportunistic screening and minimal contact psychotherapy to prevent depression in primary care patients. *PLoS ONE*. 2011;6(8):e22884.

Van Der Veen WJ, van der Meer K, Penninx BW. Screening for depression and anxiety: Correlates of non-response and cohort attrition in the Netherlands study of depression and anxiety (NESDA). *Int J Meth Psychiatr Res*. 2009 Dec;18(4):229-39.

Van der Weele GM, de JR, de Waal MW, Spinhoven P, Rooze HA, Reis R, et al. Response to an unsolicited intervention offer to persons aged ≥ 75 years after screening positive for depressive symptoms: A qualitative study. *Int Psychogeriatr*. 2012 Feb;24(2):270-7.

Van AM, Mancini C, Simpson W, Patterson B. Potential use of Internet-based screening for anxiety disorders: A pilot study. *Depress Anxiety*. 2010;27(11):1006-10.

Vaughan R, Cushman LF, Nye A. The New York City Community Outreach Study: Biomedical and mental health status among a community sample of urban Hispanics. *Ethn Dis*. 2007;17(1):99-105.

Volkers AC, Nuyen J, Verhaak PF, Schellevis FG. The problem of diagnosing major depression in elderly primary care patients. *J Affect Disord*. 2004 Oct 15;82(2):259-63.

Vuorilehto MS, Melartin TK, Isometsa ET. Course and outcome of depressive disorders in primary care: A prospective 18-month study. *Psychol Med*. 2009 Oct;39(10):1697-707.

Wagner HR, Burns BJ, Broadhead WE, Yarnall KS, Sigmon A, Gaynes BN. Minor depression in family practice: Functional morbidity, co-morbidity, service utilization and outcomes. *Psychol Med*. 2000 Nov;30(6):1377-90.

Weissman MM, Feder A, Pilowsky DJ, Olfson M, Fuentes M, Blanco C, et al. Depressed mothers coming to primary care: Maternal reports of problems with their children. *J Affect Disord*. 2004 Feb;78(2):93-100.

Weissman MM, Neria Y, Gameroff MJ, Pilowsky DJ, Wickramaratne P, Lantigua R, et al. Positive screens for psychiatric disorders in primary care: A long-term follow-up of patients who were not in treatment. *Psychiatr Serv*. 2010 Feb;61(2):151-9.

Williams JW, Jr., Kerber CA, Mulrow CD, Medina A, Aguilar C. Depressive disorders in primary care: Prevalence, functional disability, and identification. *J Gen Intern Med.* 1995 Jan;10(1):7-12.

Wittkamp KA, van Zwieten M, Smits FT, Schene AH, Huyser J, van Weert HC. Patients' view on screening for depression in general practice. *Fam Pract.* 2008 Dec;25(6):438-44.

Wong MT, Ho TP, Ho MY, Yu CS, Wong YH, Lee SY. Development and inter-rater reliability of a standardized verbal instruction manual for the Chinese Geriatric Depression Scale-short form. *Int J Geriatr Psychiatry.* 2002 May;17(5):459-63.

Wyshak G, Barsky AJ. Relationship between patient self-ratings and physician ratings of general health, depression, and anxiety. *Arch Fam Med.* 1994 May;3(5):419-24.

Yeung A, Yu SC, Fung F, Vorono S, Fava M. Recognizing and engaging depressed Chinese Americans in treatment in a primary care setting. *Int J Geriatr Psychiatry.* 2006 Sep;21(9):819-23.

ZamZam R, Thambu M, Midin M, Omar K, Kaur P. Psychiatric morbidity among adult patients in a semi-urban primary care setting in Malaysia. *Int J Ment Health Sys.* 2009;3 , 2009. Article Number: 13. Date of Publication: 18 Jun 2009.

Zimmerman M, Lish JD, Farber NJ, Hartung J, Lush D, Kuzma MA, et al. Screening for depression in medical patients. Is the focus too narrow? *Gen Hosp Psychiatry.* 1994 Nov;16(6):388-96.

Excluded: Not an eligible setting

Abas MA, Broadhead JC. Depression and anxiety among women in an urban setting in Zimbabwe. *Psychol Med.* 1997 Jan;27(1):59-71.

Alexopoulos GS, Sirey JA, Raue PJ, Kanellopoulos D, Clark TE, Novitch RS. Outcomes of depressed patients undergoing inpatient pulmonary rehabilitation. *Am J Geriatr Psychiatry.* 2006;14(5):466-75.

Allen H, Hyworon Z, Colombi A. Using self-reports of symptom severity to measure and manage workplace depression. *J Occup Environ Med.* 2010 Apr;52(4):363-74.

Arnold N, Kafetz K. Elderly medical inpatients screening positive for depression may show spontaneous improvement after discharge. *Age Ageing.* 2002 Sep;31(5):397-400.

Baker FM, Okwumabua J, Philipose V, Wong S. Screening African-American elderly for the presence of depressive symptoms: A preliminary investigation. *J Geriatr Psychiatr Neurol.* 1996 Jul;9(3):127-32.

Bambauer KZ, Locke SE, Aupont O, Mullan MG, McLaughlin TJ. Using the Hospital Anxiety and Depression Scale to screen for depression in cardiac patients. *Gen Hosp Psychiatry.* 2005 Jul;27(4):275-84.

Barth J, Martin CR. Factor structure of the Hospital Anxiety and Depression Scale (HADS) in German coronary heart disease patients. *Health Qual Life Outcomes*. 2005;3:15.

Barua A, Kar N. Screening for depression in elderly Indian population. *Indian J Psychiatr*. 2010;52(2):150-3.

Biros MH, Mann J, Hanson R, Cen YY. Unsuspected or unacknowledged depressive symptoms in young adult emergency department patients. *Acad Emerg Med*. 2009 Apr;16(4):288-94.

Booth BM, Kirchner JE, Hamilton G, Harrell R, Smith GR. Diagnosing depression in the medically ill: Validity of a lay-administered structured diagnostic interview. *J Psychiatr Res*. 1998 Nov;32(6):353-60.

Boothby JL, Durham TW. Screening for depression in prisoners using the Beck Depression Inventory. *Crim Justice Behav*. 1999 Mar;26(1):107-24.

Boudreaux ED, Clark S, Camargo CA, Jr. Mood disorder screening among adult emergency department patients: A multicenter study of prevalence, associations and interest in treatment. *Gen Hosp Psychiatry*. 2008 Jan;30(1):4-13.

Buchwald D, Manson SM, Brennenman DL, Dinges NG, Keane EM, Beals J, et al. Screening for depression among newly arrived Vietnamese refugees in primary care settings. *West J Med*. 1995 Oct;163(4):341-5.

Bushnell J, McLeod D, Dowell A, Salmond C, Ramage S, Cave R, et al. The nature and prevalence of psychological problems in New Zealand primary healthcare: A report on Mental Health and General Practice Investigation (MaGPIe). *NZ Med J*. 2003;116(1171).

Cauffman E. A statewide screening of mental health symptoms among juvenile offenders in detention. *J Am Acad Child Adolesc Psychiatry*. 2004;43(4):430-9.

Cohen CI, Hyland K, Kimhy D. The utility of mandatory depression screening of dementia patients in nursing homes. *Am J Psychiatry*. 2003 Nov;160(11):2012-7.

Cole MG, McCusker J, Elie M, Dendukuri N, Latimer E, Belzile E. Systematic detection and multidisciplinary care of depression in older medical inpatients: a randomized trial. *Can Med Assoc J*. 2006 Jan 3;174(1):38-44.

Cullum S, Tucker S, Todd C, Brayne C. Screening for depression in older medical inpatients. *Int J Geriatr Psychiatry*. 2006 May;21(5):469-76.

Cutler CB, Legano LA, Dreyer BP, Fierman AH, Berkule SB, Lusskin SI, et al. Screening for maternal depression in a low education population using a two item questionnaire. *Arch Womens Ment Health*. 2007;10(6):277-83.

Ell K, Unutzer J, Aranda M, Gibbs NE, Lee PJ, Xie B. Managing depression in home health care: A randomized clinical trial. *Home Health Care Serv Q*. 2007;26(3):81-104.

Engel CC, Oxman T, Yamamoto C, Gould D, Barry S, Stewart P, et al. RESPECT-Mil: Feasibility of a systems-level collaborative care approach to depression and post-traumatic stress disorder in military primary care. *Mil Med.* 2008 Oct;173(10):935-40.

Fabre X, Moiron L, Cannamela A, Renoux F. Depression in elderly emergency room patients aged 75 years and over. Prevalence, medical and social features, management schemes. *JEUR.* 2002;15(2):83-8.

Farragher B, Wrigley M, Donohoe O, Duggan J. Screening for depression in hospitalised elderly medical patients. *Ir J Psychol Med.* 1996;13(4):144-6.

Friedman DE, Kung DH, Laowattana S, Kass JS, Hrachovy RA, Levin HS. Identifying depression in epilepsy in a busy clinical setting is enhanced with systematic screening. *Seizure.* 2009 Jul;18(6):429-33.

Gerber MR, Leiter KS, Hermann RC, Bor DH. How and why community hospital clinicians document a positive screen for intimate partner violence: A cross-sectional study. *BMC Fam Pract.* 2005;6, 2005. Article Number: 48. Date of Publication: 19 Nov 2005.

Goldsmith ME. Postpartum depression screening by family nurse practitioners. *J Am Acad Nurse Pract.* 2007 Jun;19(6):321-7.

Harris MF, Silove D, Kehag E, Barratt A, Manicavasagar V, Pan J, et al. Anxiety and depression in general practice patients: Prevalence and management. *Med J Aust.* 1996 May 6;164(9):526-9.

Hedayati SS, Bosworth HB, Kuchibhatla M, Kimmel PL, Szczech LA. The predictive value of self-report scales compared with physician diagnosis of depression in hemodialysis patients. *Kidney Int.* 2006 May;69(9):1662-8.

Hermanns N, Kulzer B, Krichbaum M, Kubiak T, Haak T. How to screen for depression and emotional problems in patients with diabetes: Comparison of screening characteristics of depression questionnaires, measurement of diabetes-specific emotional problems and standard clinical assessment. *Diabetologia.* 2006 Mar;49(3):469-77.

Hollifield M, Katon W, Morojele N. Anxiety and depression in an outpatient clinic in Lesotho, Africa. *Int J Psychiatr Med.* 1994;24(2):179-88.

Huffman JC, Smith FA, Blais MA, Beiser ME, Januzzi JL, Fricchione GL. Recognition and treatment of depression and anxiety in patients with acute myocardial infarction. *Am J Cardiol.* 2006 Aug 1;98(3):319-24.

Hustey FM. The use of a brief depression screen in older emergency department patients. *Acad Emerg Med.* 2005 Sep;12(9):905-8.

Hustey FM, Smith MD. A depression screen and intervention for older ED patients. *Am J Emerg Med.* 2007 Feb;25(2):133-7.

Innamorati M, Pompili M, Serafini G, Lester D, Erbuto D, Amore M, et al. Psychometric properties of the suicidal history self-rating screening scale. *Arch Suicide Res.* 2011 Jan;15(1):87-92.

Jones LE, Doebbeling CC. Suboptimal depression screening following cancer diagnosis. *Gen Hosp Psychiatry.* 2007 Nov;29(6):547-54.

Keller R, Rigardetto R, Vaccarino P, Maggioni M, Iannocari G, Teriaca MJ. Screening anxious-depressive symptoms and pain in medical inpatients. *Panminerva Med.* 2008 Sep;50(3):217-20.

Kominski G, Andersen R, Bastani R, Gould R, Hackman C, Huang D, et al. UPBEAT: The impact of a psychogeriatric intervention in VA medical centers. *Medical Care.* 2001 May;39(5):500-12.

Krukowski RA, Friedman KE, Applegate KL. The utility of the Beck Depression Inventory in a bariatric surgery population. *Obesity Surgery.* 2010 Apr;20(4):426-31.

Kurlowicz LH. Benefits of psychiatric consultation-liaison nurse interventions for older hospitalized patients and their nurses. *Arch Psychiatr Nurs.* 2001 Apr;15(2):53-61.

Lafortune D. Prevalence and screening of mental disorders in short-term correctional facilities. *Int J Law Psychiatr.* 2010 Mar;33(2):94-100.

Lavretsky H, Bastani R, Gould R, Huang D, Llorente M, Maxwell A, et al. Predictors of two-year mortality in a prospective "UPBEAT" study of elderly veterans with comorbid medical and psychiatric symptoms. *Am J Geriatr Psychiatr.* 2002 Jul;10(4):458-68.

Lee AM, Chu LW, Chong CS, Chan SY, Tam S, Lam KS, et al. Relationship between symptoms of androgen deficiency and psychological factors and quality of life among Chinese men. *Int J Androl.* 2010 Oct 1;33(5):755-63.

Lee SJ, Loberiza FR, Antin JH, Kirkpatrick T, Prokop L, Alyea EP, et al. Routine screening for psychosocial distress following hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 2005;35(1):77-83.

Loke B, Nicklason F, Burvill P. Screening for depression: Clinical validation of geriatricians' diagnosis, the brief assessment schedule depression cards and the 5-item version of the symptom check list among non-demented geriatric inpatients. *Int J Geriatr Psychiatry.* 1996;11(5):461-5.

Lopes AA, Albert JM, Young EW, Satayathum S, Pisoni RL, Andreucci VE, et al. Screening for depression in hemodialysis patients: Associations with diagnosis, treatment, and outcomes in the DOPPS. *Kidney Int.* 2004 Nov;66(5):2047-53.

Manne S, Nereo N, DuHamel K, Ostroff J, Parsons S, Martini R, et al. Anxiety and depression in mothers of children undergoing bone marrow transplant: Symptom prevalence and use of the Beck depression and Beck anxiety inventories as screening instruments. *J Consult Clin Psychol.* 2001 Dec;69(6):1037-47.

Mao L, Kidd MR, Rogers G, Andrews G, Newman CE, Booth A, et al. Social factors associated with Major Depressive Disorder in homosexually active, gay men attending general practices in urban Australia. *Aust NZ J Public Health*. 2009 Feb;33(1):83-6.

Marchesi C, Brusamonti E, Borghi C, Giannini A, Di Ruvo R, Minneo F, et al. Anxiety and depressive disorders in an emergency department ward of a general hospital: A control study. *Emerg Med J*. 2004 Mar;21(2):175-9.

Mason L, Poole H. Healthcare professionals' views of screening for postnatal depression. *Community Pract*. 2008 Apr;81(4):30-3.

Massoudi P, Wickberg B, Hwang P. Screening for postnatal depression in Swedish child health care. *Acta Paediatr*. 2007 Jun;96(6):897-901.

McCusker J, Bellavance F, Cardin S, Trepanier S, Afilalo M, Belanger L, et al. Screening for geriatric problems in the emergency department: Reliability and validity. *Acad Emerg Med*. 1998;5(9):883-93.

McLearn AM, Ryba NL. Identifying severely mentally ill inmates: Can small jails comply with detection standards? *J Offender Rehabil*. 2003;37(1):25-40.

Meldon SW, Emerman CL, Schubert DS. Recognition of depression in geriatric ED patients by emergency physicians. *Ann Emerg Med*. 1997 Oct;30(4):442-7.

Milliken CS, Auchterlonie JL, Hoge CW. Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq war. *JAMA*. 2007 Nov 14;298(18):2141-8.

Mills JF, Kroner DG. Screening for suicide risk factors in prison inmates: Evaluating the efficiency of the Depression, Hopelessness and Suicide Screening Form (DHS). *Legal Criminol Psychol*. 2005;10(1):1-12.

Needlman R, Walders N, Kelly S, Higgins J, Sofranko K, Drotar D, et al. Impact of screening for maternal depression in a pediatric clinic: An exploratory study. *Ambul Child Health*. 1999;5(1):61-71.

Nguyen HV, Inderjeeth CA, Tang E, Barnabas L, Merriam M. Screening for depression in hospitalised and community-dwelling elderly: The use of the 4-item, 5-item and 15-item geriatric depression scales. *Australas J Ageing*. 2006 Dec;25(4):204-8.

Olfson M, Broadhead WE, Weissman MM, Leon AC, Farber L, Hoven C, et al. Subthreshold psychiatric symptoms in a primary care group practice. *Arch Gen Psychiatry*. 1996 Oct;53(10):880-6.

Osterman RL, Ribak J, Bohn CM, Fargo JD, Sommers MS. Screening for hazardous/harmful drinking and depressive symptoms in an at-risk emergency department population. *J Addict Nurs*. 2009 Mar;20(1):34-40.

Pailler ME, Cronholm PF, Barg FK, Wintersteen MB, Diamond GS, Fein JA. Patients' and caregivers' beliefs about depression screening and referral in the emergency department. *Pediatr Emerg Care*. 2009;25(11):721-7.

Patten SB, Lavorato DH, Metz LM. Clinical correlates of CES-D depressive symptom ratings in an MS population. *Gen Hosp Psychiatry*. 2005 Nov;27(6):439-45.

Picot MC, Abbar M, Chevalier A, Chastang JF, Luce D, Bonenfant S. Psychotropic drug use and associated factors in a professional environment. *Encephale*. 1996 Nov;22(6):403-8.

Price SK, Proctor EK. A rural perspective on perinatal depression: Prevalence, correlates, and implications for help-seeking among low-income women. *J Rural Health*. 2009;25(2):158-66.

Rumble S, Swartz L, Parry C, Zwarenstein M. Prevalence of psychiatric morbidity in the adult population of a rural South African village. *Psychol Med*. 1996;26(5):997-1007.

Scott EG, Luxmore B, Alexander H, Fenn RL, Christopher NC. Screening for adolescent depression in a pediatric emergency department. *Acad Emerg Med*. 2006 May;13(5):537-42.

Seal KH, Bertenthal D, Maguen S, Gima K, Chu A, Marmar CR. Getting beyond "Don't ask; don't tell": An evaluation of US Veterans Administration postdeployment mental health screening of veterans returning from Iraq and Afghanistan. *Am J Public Health*. 2008 Apr;98(4):714-20.

Shah A, Odutoye K, De T. Depression in acutely medically ill elderly inpatients: A pilot study of early identification and intervention by formal psychogeriatric consultation. *J Affect Disord*. 2001 Feb;62(3):233-40.

Sibitz I, Berger P, Freidl M, Topitz A, Krautgartner M, Spiegel W, et al. ICD-10 or DSM-IV? Anhedonia, fatigue and depressed mood as screening symptoms for diagnosing a current depressive episode in physically ill patients in general hospital. *J Affect Disord*. 2010 Oct;126(1-2):245-51.

Snijders AH, Robertson MM, Orth M. Beck Depression Inventory is a useful screening tool for major depressive disorder in Gilles de la Tourette syndrome. *J Neurol Neurosurg Ps*. 2006 Jun;77(6):787-9.

Soares-Filho GL, Freire RC, Biancha K, Pacheco T, Volschan A, Valenca AM, et al. Use of the Hospital Anxiety and Depression Scale (HADS) in a cardiac emergency room: Chest pain unit. *Clinics*. 2009;64(3):209-14.

Sogaard HJ, Bech P. Psychiatric disorders in long-term sickness absence -- A population-based cross-sectional study. *Scand J Public Health*. 2009 Sep;37(7):682-9.

Sorsdahl K, Flisher AJ, Ward C, Mertens J, Bresick G, Sterling S, et al. The time is now: Missed opportunities to address patient needs in community clinics in Cape Town, South Africa. *Trop Med Int Health*. 2010;15(10):1218-26.

Sowden G, Mastromauro CA, Januzzi JL, Fricchione GL, Huffman JC. Detection of depression in cardiac inpatients: Feasibility and results of systematic screening. *Am Heart J*. 2010 May;159(5):780-7.

Thompson DA, Leimig R, Gower G, Winsett RP. Assessment of depressive symptoms during post-transplant follow-up care performed via telehealth. *Telemed J E Health*. 2009 Sep;15(7):700-6.

Upshur C, Weinreb L. A survey of primary care provider attitudes and behaviors regarding treatment of adult depression: What changes after a collaborative care intervention? *Prim Care Comp J Clin Psychiatr*. 2008;10(3):182-6.

Vahter L, Kreegipuu M, Talvik T, Gross-Paju K. One question as a screening instrument for depression in people with multiple sclerosis. *Clin Rehab*. 2007;21(5):460-4.

Vahter L, Braschinsky M, Haldre S, Gross-Paju K. The prevalence of depression in hereditary spastic paraplegia. *Clin Rehab*. 2009 Sep;23(9):857-61.

Wada K, Tanaka K, Theriault G, Moriyama M, Satoh T, Aizawa Y. Application of the stratum-specific likelihood ratio (SSLR) analysis to results of a depressive symptoms screening survey among Japanese workers. *Soc Psychiatry Psychiatr Epidemiol*. 2007 May;42(5):410-3.

Wilson B, Spittal J, Heidenheim P, Herman M, Leonard M, Johnston A, et al. Screening for depression in chronic hemodialysis patients: Comparison of the Beck Depression Inventory, primary nurse, and nephrology team. *Hemodial Int*. 2006 Jan;10(1):35-41.

Yackel EE, McKennan MS, Fox-Deise A. A nurse-facilitated depression screening program in an Army primary care clinic: An evidence-based project. *Nursing Research*. 2010 Jan;59(1:Suppl):Suppl-65.

Yohannes AM, Baldwin RC, Connolly MJ. Depression and anxiety in elderly outpatients with chronic obstructive pulmonary disease: Prevalence, and validation of the BASDEC screening questionnaire. *Int J Geriatr Psychiatry*. 2000 Dec;15(12):1090-6.

Excluded: Only about harms of treatment

Report of the CSM expert working group on the safety of selective serotonin reuptake inhibitor antidepressants. Medicines and Healthcare Products Regulatory Agency (MHRA); 2004.

Ammerman RT, Putnam FW, Stevens J, Bosse NR, Short JA, Bodley AL, et al. An open trial of in-home CBT for depressed mothers in home visitation. *Matern Child Health J*. 2011 Nov;15(8):1333-41.

Blitzer LE, Atchison-Nevel DJ, Kenny MC. Using acupuncture to treat major depressive disorder: A pilot investigation. *Clin Acupuncture Oriental Med*. 2003;4(4):144-7.

Corey-Lisle PK, Nash R, Stang P, Swindle R. Response, partial response, and nonresponse in primary care treatment of depression. *Arch Intern Med*. 2004 Jun 14;164(11):1197-204.

Dalton SO, Sorensen HT, Johansen C. SSRIs and upper gastrointestinal bleeding: What is known and how should it influence prescribing? *CNS Drugs*. 2006;20(2):143-51.

Dobscha SK, Gerrity MS, Corson K, Bahr A, Cuilwik NM. Measuring adherence to depression treatment guidelines in a VA primary care clinic. *Gen Hosp Psychiatry*. 2003 Jul;25(4):230-7.

Dodd S, Malhi GS, Tiller J, Schweitzer I, Hickie I, Khoo JP, et al. A consensus statement for safety monitoring guidelines of treatments for major depressive disorder. *Aust NZ J Psychiatry*. 2011 Sep;45(9):712-25.

Fergusson D, Doucette S, Glass KC, Shapiro S, Healy D, Hebert P, et al. Association between suicide attempts and selective serotonin reuptake inhibitors: Systematic review of randomised controlled trials. *Br Med J*. 2005 Feb 19;330(7488):396.

Gaynes BN, Rush AJ, Trivedi MH, Wisniewski SR, Balasubramani GK, McGrath PJ, et al. Primary versus specialty care outcomes for depressed outpatients managed with measurement-based care: Results from STAR*D. *J Gen Intern Med*. 2008 May;23(5):551-60.

Gibbons RD, Brown CH, Hur K, Marcus SM, Bhaumik DK, Mann JJ. Relationship between antidepressants and suicide attempts: An analysis of the Veterans Health Administration data sets. *Am J Psychiatry*. 2007 Jul;164(7):1044-9.

Horowitz JA, Cousins A. Postpartum depression treatment rates for at-risk women. *Nursing Research*. 2006;55(2 Suppl):S23-S27.

Jick SS, Dean AD, Jick H. Antidepressants and suicide. *Br Med J*. 1995 Jan 28;310(6974):215-8.

Katon W, Russo J, Lin EHB, Schmittdiel J, Ciechanowski P, Ludman E, et al. Cost-effectiveness of a multicondition collaborative care intervention. *Arch Gen Psychiatry*. 2012;69(5):506-14.

Khan A, Khan S, Kolts R, Brown WA. Suicide rates in clinical trials of SSRIs, other antidepressants, and placebo: Analysis of FDA reports. *Am J Psychiatry*. 2003 Apr;160(4):790-2.

Kroenke K, West SL, Swindle R, Gilseman A, Eckert GJ, Dolor R, et al. Similar effectiveness of paroxetine, fluoxetine, and sertraline in primary care: A randomized trial. *JAMA*. 2001 Dec 19;286(23):2947-55.

Levenson M, Holland C. Statistical evaluation of suicidality in adults treated with antidepressants. 2006.

Lin HC, Lin YJ, Hsiao FH, Li CY. Prenatal care visits and associated costs for treatment-seeking women with depressive disorders. *Psychiatr Serv*. 2009 Sep;60(9):1261-4.

Lucock M, Kirby R, Wainwright N. A pragmatic randomized controlled trial of a guided self-help intervention versus a waiting list control in a routine primary care mental health service. *Br J Clin Psychol*. 2011 Sep;50(3):298-309.

Mackay FJ, Dunn NR, Mann RD. Antidepressants and the serotonin syndrome in general practice. *Br J Gen Pract*. 1999 Nov;49(448):871-4.

Martinez C, Rietbrock S, Wise L, Ashby D, Chick J, Moseley J, et al. Antidepressant treatment and the risk of fatal and non-fatal self harm in first episode depression: Nested case-control study. *Br Med J*. 2005 Feb 19;330(7488):389.

Mulrow CD, Williams JW, Jr., Chiquette E, Aguilar C, Hitchcock-Noel P, Lee S, et al. Efficacy of newer medications for treating depression in primary care patients. *Am J Med.* 2000 Jan;108(1):54-64.

Nyunt MS, Ko SM, Kumar R, Fones CC, Ng TP. Improving treatment access and primary care referrals for depression in a national community-based outreach programme for the elderly. *Int J Geriatr Psychiatry.* 2009 Nov;24(11):1267-76.

Olfson M, Marcus SC, Tedeschi M, Wan GJ. Continuity of antidepressant treatment for adults with depression in the United States. *Am J Psychiatry.* 2006 Jan;163(1):101-8.

Pinquart M, Sorensen S. How effective are psychotherapeutic and other psychosocial interventions with older adults? A meta-analysis. *J Ment Health Aging.* 2001;7(2):207-43.

Richards JB, Papaioannou A, Adachi JD, Joseph L, Whitson HE, Prior JC, et al. Effect of selective serotonin reuptake inhibitors on the risk of fracture. *Arch Intern Med.* 2007 Jan 22;167(2):188-94.

Robinson W, Geske J, Prest L, Barnacle R. Depression treatment in primary care. *J Am Board Fam Pract.* 2005;18:79-86.

Serfaty M, Wilkinson S, Freeman C, Mannix K, King M. The ToT Study: Helping with Touch or Talk (ToT): A pilot randomised controlled trial to examine the clinical effectiveness of aromatherapy massage versus cognitive behaviour therapy for emotional distress in patients in cancer/palliative care. *Psycho Oncol.* 2012;21(5):563-9.

Simon GE, Fleck M, Lucas R, Bushnell DM. Prevalence and predictors of depression treatment in an international primary care study. *Am J Psychiatry.* 2004;161(9):1626-34.

Simon GE, Savarino J, Operskalski B, Wang PS. Suicide risk during antidepressant treatment. *Am J Psychiatry.* 2006 Jan;163(1):41-7.

Sondergard L, Kvist K, Andersen PK, Kessing LV. Do antidepressants prevent suicide? *Int Clin Psychopharmacol.* 2006 Jul;21(4):211-8.

Thase ME, Entsuah AR, Rudolph RL. Remission rates during treatment with venlafaxine or selective serotonin reuptake inhibitors. *Br J Psychiatry.* 2001 Mar;178:234-41.

Trinh NH, Bedoya CA, Chang TE, Flaherty K, Fava M, Yeung A. A study of a culturally focused psychiatric consultation service for Asian American and Latino American primary care patients with depression. *BMC Psychiatry.* 2011;11:166.

Trivedi MH, Rush AJ, Wisniewski SR, Nierenberg AA, Warden D, Ritz L, et al. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: Implications for clinical practice. *Am J Psychiatry.* 2006 Jan;163(1):28-40.

Urizar GG, Jr., Munoz RF. Impact of a prenatal cognitive-behavioral stress management intervention on salivary cortisol levels in low-income mothers and their infants. *Psychoneuroendocrinol.* 2011 Nov;36(10):1480-94.

Wagner J, Allen NA, Swalley LM, Melkus GD, Whittemore R. Depression, depression treatment, and insulin sensitivity in adults at risk for type 2 diabetes. *Diabetes Res Clin Pract.* 2009 Nov;86(2):96-103.

Wells KB, Tang L, Carlson GA, Asarnow JR. Treatment of youth depression in primary care under usual practice conditions: Observational findings from youth partners in care. *J Child Adolesc Psychiatr Nurs.* 2012;22(1):80-90.

Zhou W, He G, Gao J, Yuan Q, Feng H, Zhang CK. The effects of group reminiscence therapy on depression, self-esteem, and affect balance of Chinese community-dwelling elderly. *Arch Gerontol Geriatr.* 2012;54(3):e440-e447.

Excluded: Modeling paper with limited applicability for our research question

Valenstein M, Vijan S, Zeber JE, Boehm K, and Buttar A. The cost-utility of screening for depression in primary care. *Ann Intern Med.* 2001; 134(5):345-60. PMID:11242495.