

## Adult Obesity KQ1 Prevention Excluded Studies List

### No outcomes for population of interest

The effects of nonpharmacologic interventions on blood pressure of persons with high normal levels. Results of the Trials of Hypertension Prevention, Phase I. *JAMA : the journal of the American Medical Association* 1992;267(9):1213-20.

Randomised controlled trial evaluating cardiovascular screening and intervention in general practice: principal results of British family heart study. Family Heart Study Group. *BMJ (Clinical research ed)* 1994;308(6924):313-20.

The Diabetes Prevention Program. Design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care* 1999;22(4):623.

Managing weight gain after breast cancer. *Journal of Supportive Oncology* 2006;4(2):69-70.

Nutrition, rehabilitation/quality of life, and cardiovascular special studies. *Am J Kidney Dis* 2012;59(1 SUPPL. 1):e267-e270.

Abildso CG, Zizzi SJ, Reger-Nash B. Evaluating an insurance-sponsored weight management program with the RE-AIM Model, West Virginia, 2004-2008. *Prev Chronic Dis* 2010 May;7(3):A46.

Abildso C, Zizzi S, Gilleland D, Thomas J, Bonner D. A mixed methods evaluation of a 12-week insurance-sponsored weight management program incorporating cognitive-behavioral counseling. *Journal of Mixed Methods Research* 2010 Oct;4(4):278-94.

Absetz P, Valve R, Oldenburg B, Heinonen H, Nissinen A, Fogelholm M, et al. Type 2 diabetes prevention in the "real world": one-year results of the GOAL Implementation Trial. *Diabetes Care* 2007;30(10):2465-70.

Absetz P, Oldenburg B, Hankonen N, Valve R, Heinonen H, Nissinen A, et al. Type 2 diabetes prevention in the real world: three-year results of the GOAL lifestyle implementation trial. *Diabetes Care* 2009;32(8):1418-20.

Abujudeh BA, Abu Al Rub RF, Al-Faouri IG, Gharaibeh MK. The impact of lifestyle modification in preventing or delaying the progression of type 2 diabetes mellitus among high-risk people in Jordan. *Journal of Research in Nursing* 2012;17(1):32-44.

Ackermann RT, Finch EA, Brizendine E, Zhou H, Marrero DG. Translating the Diabetes Prevention Program into the community. The DEPLOY Pilot Study. *Am J Prev Med* 2008;35(4):357-63.

Ackermann RT, Edelstein SL, Narayan KMV, Zhang P, Engelgau MM, Herman WH, et al. Changes in health state utilities with changes in body mass in the diabetes prevention program. *Obesity* 2009 Dec;17(12):2176-81.

Ackermann RT, Finch EA, Caffrey HM, Lipscomb ER, Hays LM, Saha C. Long-term effects of a community-based lifestyle intervention to prevent type 2 diabetes: the DEPLOY extension pilot study. *Chronic Illn* 2011;7(4):279-90.

Ades PA, Savage PD, Harvey-Berino J. The treatment of obesity in cardiac rehabilitation. [Review]. *Journal of Cardiopulmonary Rehabilitation & Prevention* 2010 Sep;30(5):289-98.

Agarwal N, Rice SPL, Bolusani H, Luzio SD, Dunseath G, Ludgate M, et al. Metformin reduces arterial stiffness and improves endothelial function in young women with polycystic ovary syndrome: A randomized, placebo-controlled, crossover trial. *J Clin Endocrinol Metab* 2010;95(2):722-30.

Aggarwal B. Predictors of physical activity at 1 year in a randomized controlled trial of family members of patients with cardiovascular disease. *The Journal of cardiovascular nursing* 2010;25(6):444.

Akcakoyun F. Changes in serum lipid profile following moderate exercise. *African Journal of Pharmacy and Pharmacology* 2010;4(11):829-33.

Al-Solaiman Y, Jesri A, Mountford WK, Lackland DT, Zhao Y, Egan BM. DASH lowers blood pressure in obese hypertensives beyond potassium, magnesium and fibre. *J Hum Hypertens* 2010 Apr;24(4):237-46.

Alahuhta MA, Korhakangas EE, Keranen AM, Kyngas HA, Laitinen JH. Using pictures as vignettes to assess stages of change in weight management. *SCAND J PUBLIC HEALTH* 2011 Jun;39(4):403-9.

Albarran NB, Ballesteros MN, Morales GG, Ortega MI. Dietary behavior and type 2 diabetes care. *Patient Education and Counseling* 2006;61(2):191-9.

Albu J, Gottlieb SH, August P, Nesto RW, Orchard TJ, Bypass AR, I. Modifications of coronary risk factors. *Am J Cardiol* 2006 Jun 19;97(12A):41G-52G.

Aldrich ND, Reicks MM, Sibley SD, Redmon JB, Thomas W, Raatz SK. Varying protein source and quantity do not significantly improve weight loss, fat loss, or satiety in reduced energy diets among midlife adults. *Nutr Res* 2011 Feb;31(2):104-12.

Allen N, Whittemore R, Melkus G. A continuous glucose monitoring and problem-solving intervention to change physical activity behavior in women with type 2 diabetes: a pilot study. *DIABETES TECHNOLOG THER* 2011 Nov;13(11):1091-9.

Allen P, Thompson JL, Herman CJ, Whyte AN, Wolfe VK, Qualls C, et al. Impact of periodic follow-up testing among urban American Indian women with impaired fasting glucose. *Prev Chronic Dis* 2008 Jul;5(3):A76.

Almeida FA, Shetterly S, Smith-Ray RL, Estabrooks PA. Reach and effectiveness of a weight loss intervention in patients with prediabetes in Colorado. *Prev Chronic Dis* 2010 Sep;7(5):A103.

Almendingen K, Trygg K, Pedersen JI. An assessment of the use of simple methods to predict individual energy intakes for intervention studies. *Eur J Clin Nutr* 1998 Jan;52(1):54-9.

Amundson HA, Butcher MK, Gohdes D, Hall TO, Harwell TS, Helgerson SD, et al. Translating the diabetes prevention program into practice in the general community: Findings from the Montana Cardiovascular Disease and Diabetes Prevention Program. *The Diabetes educator* 2009;35(2):209-23.

Andersen RE, Franckowiak SC, Snyder J, Bartlett SJ, Fontaine KR. Can inexpensive signs encourage the use of stairs? Results from a community intervention. *Ann Intern Med* 1998;129(5):363-9.

Andersen RE, Franckowiak SC, Zuzak KB, Cummings ES, Bartlett SJ, Crespo CJ. Effects of a culturally sensitive sign on the use of stairs in African American commuters. *Sozial- und Praventivmedizin* 2006;51(6):373-80.

Anderssen S, Holme I, Urdal P, Hjermann I. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press* 1995 Nov;4(6):343-9.

Andrews RC, Cooper AR, Montgomery AA, Norcross AJ, Peters TJ, Sharp DJ, et al. Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: The Early ACTID randomised controlled trial. *Lancet* 2011 Jul 9;378(9786):129-39.

Annesi JJ. Relation of perceived fitness with health changes and physiological changes in women exercisers. *PERCEPT MOTOR SKILLS* 2002 Aug;95(1):226.

Anton SD, Martin CK, Han H, Coulon S, Cefalu WT, Geiselman P, et al. Effects of stevia, aspartame, and sucrose on food intake, satiety, and postprandial glucose and insulin levels. *Appetite* 2010 Aug;55(1):37-43.

Apovian CM, Bergenstal RM, Cuddihy RM, Qu Y, Lenox S, Lewis MS, et al. Effects of exenatide combined with lifestyle modification in patients with type 2 diabetes. *Am J Med* 2010 May;123(5):468-17.

Archer SL, Greenlund KJ, Casper ML, Rith-Najarian S, Croft JB. Associations of community-based health education programs with food habits and cardiovascular disease risk factors among Native Americans with diabetes: the Inter-Tribal Heart Project, 1992 to 1994. *J AM DIET ASSOC* 2002 Aug;102(8):1132-5.

Ard JD, Kumanyika S, Stevens VJ, Vollmer WM, Samuel-Hodge C, Kennedy B, et al. Effect of group racial composition on weight loss in African Americans. *Obesity* 2008 Feb;16(2):306-10.

Arechavaleta R, Seck T, Chen Y, Krobot KJ, O'Neill EA, Duran L, et al. Efficacy and safety of treatment with sitagliptin or glimepiride in patients with type 2 diabetes inadequately controlled on metformin monotherapy: A randomized, double-blind, non-inferiority trial. *Diabetes Obes Metab* 2010;13(2):160-8.

Armitage CJ. Evidence that implementation intentions reduce dietary fat intake: A randomized trial. *Health Psychol* 2004;23(3):319-23.

Asadipooya K, Kalantar-Hormozi M, Nabipour I. Pioglitazone reduces central obesity in polycystic ovary syndrome women. *Gynecol Endocrinol* 2012;28(1):16-9.

Aschner P, Katzeff HL, Guo H, Sunga S, Williams-Herman D, Kaufman KD, et al. Efficacy and safety of monotherapy of sitagliptin compared with metformin in patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12(3):252-61.

Ashley JM, St Jeor ST, Schrage JP, Perumean-Chaney SE, Gilbertson MC, McCall NL, et al. Weight control in the physician's office. *Arch Intern Med* 2001;161(13):1599-604.

Atiomo W, Read A, Golding M, Silcocks P, Razali N, Sarkar S, et al. Local recruitment experience in a study comparing the effectiveness of a low glycaemic index diet with a low calorie healthy eating approach at achieving weight loss and reducing the risk of endometrial cancer in women with polycystic ovary syndrome (PCOS). *Contemp Clin Trials* 2009 Sep;30(5):451-6.

Atlantis E, Chow CM, Kirby A, Singh MF. An effective exercise-based intervention for improving mental health and quality of life measures: A randomized controlled trial. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 2004 Aug;39(2):424-34.

Axelrod L, Halter JB, Cooper DS. Hormone levels and fuel flow in patients with weight loss and lung cancer. Evidence for excessive metabolic expenditure and for an adaptive response mediated by a reduced level of 3,5,3'-triiodothyronine. *Metabolism* 1983;32(9):924-37.

Babamoto KS, Sey KA, Camilleri AJ, Karlan VJ, Catalasan J, Morisky DE. Improving diabetes care and health measures among hispanics using community health workers: results from a randomized controlled trial. *HEALTH EDUC BEHAV* 2009 Feb;36(1):113-26.

Bailey BW, Jacobsen DJ, Donnelly JE. Weight loss and maintenance outcomes using moderate and severe restriction in an outpatient setting. *Disease Management* 2008 Jun;11(3):176-80.

Bain SC, McKenna J. Clinical experience with liraglutide. *INT J CLIN PRACT* 2010;64(SUPPL. 167):44-8.

Baker RC, Kirschenbaum DS. Weight control during the holidays: Highly consistent self-monitoring as a potentially useful coping mechanism. *Health Psychol* 1998;17(4):367-70.

Banegas JR, Lopez-Garcia E, Dallongeville J, Guallar E, Halcox JP, Borghi C, et al. Achievement of treatment goals for primary prevention of cardiovascular disease in clinical practice across Europe: the EURIKA study. *Eur Heart J* 2011 Sep;32(17):2143-52.

Banks-Wallace J. Outcomes from Walk the Talk: a nursing intervention for Black women. *ABNF J* 2007;18(1):19-24.

Banzer JA, Maguire TE, Kennedy CM, O'Malley CJ, Balady GJ. Results of cardiac rehabilitation in patients with diabetes mellitus. *Am J Cardiol* 2004 Jan 1;93(1):81-4.

Barnard ND, Gloede L, Cohen J, Jenkins DJ, Turner-McGrievy G, Green AA, et al. A low-fat vegan diet elicits greater macronutrient changes, but is comparable in adherence and acceptability, compared with a more conventional diabetes diet among individuals with type 2 diabetes. *J AM DIET ASSOC* 2009 Feb;109(2):263-72.

Barnstuble JA, Klesges RC, Terbizan D. Predictors of weight loss in a behavioral treatment program. *BEHAV THER* 1986 Jun;17(3):288-94.

Baron P, Roy J, Ranger N. Stimulus control and reduction of weight as a function of degree of extroversion. [French]. *Revue de Modification du Comportement* 1984;14(1):29-34.

Barratt R, Frost G, Millward DJ, Truby H. A randomised controlled trial investigating the effect of an intensive lifestyle intervention v. standard care in adults with type 2 diabetes immediately after initiating insulin therapy. *Br J Nutr* 2008 May;99(5):1025-31.

Barte JCM, ter Bogt NCW, Beltman FW, van der Meer K, Bemelmans WJE. Process evaluation of a lifestyle intervention in primary care: implementation issues and the participants' satisfaction of the GOAL study. *Health education & behavior : the official publication of the Society for Public Health Education* 2012;39(5):564-73.

Bavikati VV, Sperling LS, Salmon RD, Faircloth GC, Gordon TL, Franklin BA, et al. Effect of comprehensive therapeutic lifestyle changes on prehypertension. *Am J Cardiol* 2008 Dec 15;102(12):1677-80.

Bays H. Phentermine, topiramate and their combination for the treatment of adiposopathy ('sick fat') and metabolic disease. [Review]. *Expert Rev Cardiovasc Ther* 2010 Dec;8(12):1777-801.

Bazzarre TL, Evans BW, Kinard JD, Truslow WW. The effects of attendance during a 12-week lifestyle management program on body weight, body fat, cardiovascular performance and blood lipids. *Journal of Obesity & Weight Regulation* 1985;4(4):274-81.

Beavers KM, Lyles MF, Davis CC, Wang X, Beavers DP, Nicklas BJ. Is lost lean mass from intentional weight loss recovered during weight regain in postmenopausal women? *Am J Clin Nutr* 2011 Sep;94(3):767-74.

Beck-Nielsen H, Pedersen O, Sorensen NS. Effects of dietary changes on cellular insulin binding and in vivo insulin sensitivity. *Metabolism* 1980;29(5):482-7.

Belski R, Mori TA, Puddey IB, Sipsas S, Woodman RJ, Ackland TR, et al. Effects of lupin-enriched foods on body composition and cardiovascular disease risk factors: a 12-month randomized controlled weight loss trial. *Int J Obes* 2011 Jun;35(6):810-9.

Benassi-Evans B, Clifton P, Noakes M, Fenech M. High-protein/high red meat and high-carbohydrate weight-loss diets do not differ in their effect on faecal water genotoxicity tested by use of the WIL2-NS cell line and with other biomarkers of bowel health. *Mutation Research* 2010 Dec 21;703(2):130-6.

Bendixen H, Madsen J, Bay-Hansen D, Boesen U, Ovesen LF, Bartels EM, et al. An observational study of slimming behavior in Denmark in 1992 and 1998. *Obes Res* 2002 Sep;10(9):911-22.

Benyamini Y, Raz O. "I can tell you if I'll really lose all that weight": Dispositional and situated optimism as predictors of weight loss following a group intervention. *Journal of Applied Social Psychology* 2007 Apr;37(4):844-61.

Bergenstal R, Lewin A, Bailey T, Chang D, Gylvin T, Roberts V, et al. Efficacy and safety of biphasic insulin aspart 70/30 versus exenatide in subjects with type 2 diabetes failing to achieve glycemic control with metformin and a sulfonylurea. *Curr Med Res Opin* 2009 Jan;25(1):65-75.

Best JH, Boye KS, Rubin RR, Cao D, Kim TH, Peyrot M. Improved treatment satisfaction and weight-related quality of life with exenatide once weekly or twice daily. *Diabet Med* 2009 Jul;26(7):722-8.

Best JH, Rubin RR, Peyrot M, Li Y, Yan P, Malloy J, et al. Weight-related quality of life, health utility, psychological well-being, and satisfaction with exenatide once weekly compared with sitagliptin or pioglitazone after 26 weeks of treatment. *Diabetes Care* 2011 Feb;34(2):314-9.

Bettens C, Heraief E, Burckhardt P. Short and long term results of a progressive reintroduction of carbohydrates (PRCH) after a protein-sparing modified fast (PSMF). *Int J Obes* 1989;13:Suppl-7.

Beyazit E, Mollaoglu M. Investigation of effect on glycosylated hemoglobin, blood pressure, and body mass index of diabetes intensive education program in patients with type 2 diabetes mellitus. *Am j men's health* 2011 Jul;5(4):351-7.

Binks M, Van MT. Utilization patterns and user characteristics of an ad libitum Internet weight loss program. *J Med Internet Res* 2010;12(1):e9.

Black DR, Friesen JG. Deposit money: A component in a self-directed minimal intervention program for weight control. *BEHAV THER* 1983 Mar;14(2):333-40.

Black DR, Lantz CE. Spouse involvement and a possible long-term follow-up trap in weight loss. *Behav Res Ther* 1984;22(5):557-62.

Blackburn GL, Wilson GT, Kanders BS, Stein LJ, Lavin PT, Adler J, et al. Weight cycling: the experience of human dieters. *Am J Clin Nutr* 1989 May;49(5:Suppl):Suppl-9.

Blair AJ, Lewis VJ, Booth DA. Response to leaflets about eating and shape by women concerned about their weight. *Behavioural Psychotherapy* 1992;20(3):279-86.

Blevins T, Han J, Nicewarner D, Chen S, Oliveira JHA, Aronoff S. Exenatide is non-inferior to insulin in reducing HbA1c: An integrated analysis of 1423 patients with type 2 diabetes. *POSTGRAD MED* 2010;122(3):118-28.

Blevins T, Pullman J, Malloy J, Yan P, Taylor K, Schulteis C, et al. DURATION-5: exenatide once weekly resulted in greater improvements in glycemic control compared with exenatide twice daily in patients with type 2 diabetes. *Journal of Clinical Endocrinology & Metabolism* 2011 May;96(5):1301-10.

Blickle JF, Hancu N, Piletic M, Profozic V, Shestakova M, Dain MP, et al. Insulin glargine provides greater improvements in glycaemic control vs. intensifying lifestyle management for people with type 2 diabetes treated with OADs and 7-8% A1c levels. The TULIP study. *Diabetes Obes Metab* 2009;11(4):379-86.

Blonde L, Dagogo-Jack S, Banerji MA, Pratley RE, Marcellari A, Braceras R, et al. Comparison of vildagliptin and thiazolidinedione as add-on therapy in patients inadequately controlled with metformin: results of the GALIANT trial--a primary care, type 2 diabetes study. *Diabetes Obes Metab* 2009 Oct;11(10):978-86.

Bluher M, Rudich A, Kloting N, Golan R, Henkin Y, Rubin E, et al. Two patterns of adipokine and other biomarker dynamics in a long-term weight loss intervention. *Diabetes Care* 2012;35(2):342-9.

Bo S, Ciccone G, Guidi S, Gambino R, Durazzo M, Gentile L, et al. Diet or exercise: what is more effective in preventing or reducing metabolic alterations? *EUR J ENDOCRINOL* 2008;159(6):685-91.

Bo S, Ciccone G, Baldi C, Benini L, Dusio F, Forastiere G, et al. Effectiveness of a lifestyle intervention on metabolic syndrome. A randomized controlled trial. *J Gen Intern Med* 2007;22(12):1695-703.

Bode BW, Testa MA, Magwire M, Hale PM, Hammer M, Blonde L, et al. Patient-reported outcomes following treatment with the human GLP-1 analogue liraglutide or glimepiride in monotherapy: Results from a randomized controlled trial in patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12(7):604-12.

Boeckner LS, Kohn H, Rockwell SK. A risk-reduction nutrition course for adults. *J AM DIET ASSOC* 1990 Feb;90(2):260-3.

Boeckner LS, McClelland JW, Britten P, Chapman-Novakofski K, Mustian D, Clark CD, et al. Evaluating diverse weight management programs with a standard evaluation questionnaire. *J NUTR EDUC* 1999 Sep;31(5):262-8.

Bolinder J, Ljunggren O, Kullberg J, Johansson L, Wilding J, Langkilde AM, et al. Effects of dapagliflozin on body weight, total fat mass, and regional adipose tissue distribution in patients with type 2 diabetes mellitus with inadequate glycemic control on metformin. *J Clin Endocrinol Metab* 2012;97(3):1020-31.

Bolli G, Dotta F, Colin L, Minic B, Goodman M. Comparison of vildagliptin and pioglitazone in patients with type 2 diabetes inadequately controlled with metformin. *Diabetes Obes Metab* 2009 Jun;11(6):589-95.

Bolocofsky DN, Spinler D, Coulthard-Morris L. Effectiveness of hypnosis as an adjunct to behavioral weight management. *Journal of Clinical Psychology* 1985 Jan;41(1):35-41.

Booth AO, Nowson CA, Huang N, Lombard C, Singleton KL. Evaluation of a brief pilot nutrition and exercise intervention for the prevention of weight gain in general practice patients. *Public Health Nutr* 2006 Dec;9(8):1055-61.

Borhani NO. Significance of physical activity for prevention and control of hypertension. *J Hum Hypertens* 1996;10(SUPPL. 2):S7-S11.

Boschi V, Muscariello E, Maresca I, Ricciardi Lo SF, Tranchese V, Bellini O, et al. Assessment of eating behaviour in young women requesting nutritional counselling and their mothers. *Eating & Weight Disorders: EWD* 2010 Mar;15(1-2):e60-e67.

Bosworth HB, Olsen MK, Neary A, Orr M, Grubber J, Svetkey L, et al. Take Control of Your Blood Pressure (TCYB) study: a multifactorial tailored behavioral and educational intervention for achieving blood pressure control. *PATIENT EDUC COUNS* 2008 Mar;70(3):338-47.

Boudhraa K, Jellouli MA, Amri M, Farhat M, Torkhani F, Gara MF. [Indication of metformin in the management of hormonal dysfunction secondary to polycystic ovarian syndrome: prospective comparative study about 63 cases] TO: La metformine dans la prise en charge de l'infertilité du SOPK hors FIV: étude prospective comparative a propos de 63 cas LA: *Fre. Tunisie Medicale* 2010;88(5):335-40.

Bowles SV, Picano J, Epperly T, Myer S. The LIFE program: a wellness approach to weight loss. *Mil Med* 2006 Nov;171(11):1089-94.

Bradford A, Tuffrey V, Polley M. Effect of auricular acupuncture on appetite: A pilot Study. *Med Acupunct* 2010;22(3):181-6.

Brady AJ, Perry C, Murdoch DL, McKay G. Sustained benefits of a health project for middle-aged football supporters, at Glasgow Celtic and Glasgow Rangers Football Clubs. *Eur Heart J* 2010 Dec;31(24):2696-8.

Bray GA, Smith SR, Dejonge L, De SR, Rood J, Champagne CM, et al. Effect of diet composition on energy expenditure during weight loss: The POUNDS LOST Study. *Int J Obes* 2012;36(3):448-55.

Brooking LA, Williams SM, Mann JI. Effects of macronutrient composition of the diet on body fat in indigenous people at high risk of type 2 diabetes. *Diabetes Res Clin Pract* 2012;96(1):40-6.

Browning JD, Baker JA, Rogers T, Davis J, Satapati S, Burgess SC. Short-term weight loss and hepatic triglyceride reduction: evidence of a metabolic advantage with dietary carbohydrate restriction. *Am J Clin Nutr* 2011 May;93(5):1048-52.

Bunck MC, Eliasson B, Corner A, Heine RJ, Shaginian RM, Taskinen MR, et al. Exenatide treatment did not affect bone mineral density despite body weight reduction in patients with type 2 diabetes. *Diabetes Obes Metab* 2011 Apr;13(4):374-7.

Burke LE, Styn MA, Steenkiste AR, Music E, Warziski M, Choo J. A randomized clinical trial testing treatment preference and two dietary options in behavioral weight management: preliminary results of the impact of diet at 6 months--PREFER study. *Obesity* 2006 Nov;14(11):2007-17.

Burke V, Mansour J, Beilin LJ, Mori TA. Long-term follow-up of participants in a health promotion program for treated hypertensives (ADAPT). *Nutrition Metabolism & Cardiovascular Diseases* 2008 Mar;18(3):198-206.

Burke V, Mansour J, Mori TA, Beilin LJ, Cutt HE, Wilson A. Changes in cognitive measures associated with a lifestyle program for treated hypertensives: a randomized controlled trial (ADAPT). *HEALTH EDUC RES* 2008 Apr;23(2):202-17.

Buse JB, Rosenstock J, Sesti G, Schmidt WE, Montanya E, Brett JH, et al. Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomised, parallel-group, multinational, open-label trial (LEAD-6). *Lancet* 2009 Jul;374(9683):39-47.

Buse JB, Drucker DJ, Taylor KL, Kim T, Walsh B, Hu H, et al. DURATION-1: Exenatide once weekly produces sustained glycemic control and weight loss over 52 weeks. *Diabetes Care* 2010;33(6):1255-61.

Butryn ML, Thomas JG, Lowe MR. Reductions in internal disinhibition during weight loss predict better weight loss maintenance. *Obesity* 2009 May;17(5):1101-3.

Byrne S, Barry D, Petry NM. Predictors of weight loss success. Exercise vs. dietary self-efficacy and treatment attendance. *Appetite* 2012 Apr;58(2):695-8.

Caan B, Ballard-Barbash R, Slattery ML, Pinsky JL, Iber FL, Mateski DJ, et al. Low energy reporting may increase in intervention participants enrolled in dietary intervention trials. *J AM DIET ASSOC* 2004;104(3):357-66.

Camhi SM, Stefanick ML, Ridker PM, Young DR. Changes in C-reactive protein from low-fat diet and/or physical activity in men and women with and without metabolic syndrome. *Metabolism* 2010 Jan;59(1):54-61.

Campbell MK, Tessaro I, DeVellis B, Benedict S, Kelsey K, Belton L, et al. Effects of a tailored health promotion program for female blue-collar workers: Health Works for Women. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 2002 Mar;34(3):313-23.

Carey MG, Al-Zaiti SS, Liao LM, Martin HN, Butler RA. A low-glycemic nutritional fitness program to reverse metabolic syndrome in professional firefighters: results of a pilot study. *The Journal of cardiovascular nursing* 2011;26(4):298-304.

Caruso-Davis MK, Guillot TS, Podichetty VK, Mashtalir N, Dhurandhar NV, Dubuisson O, et al. Efficacy of low-level laser therapy for body contouring and spot fat reduction. *Obes Surg* 2011 Jun;21(6):722-9.

Casas RS, Pettee Gabriel KK, Kriska AM, Kuller LH, Conroy MB. Association of leisure physical activity and sleep with cardiovascular risk factors in postmenopausal women. *Menopause* 2012;19(4):413-9.

Chakkalakal RJ, Camp AW, Magenheimer E, Savoye M, Lubsen J, Lucas G, et al. Preventing diabetes among Fair Haven families: a community-based approach to quality improvement. *J Health Care Poor Underserved* 2012;23(3 Suppl):247-54.

Champagne CM, Broyles ST, Moran LD, Cash KC, Levy EJ, Lin PH, et al. Dietary intakes associated with successful weight loss and maintenance during the Weight Loss Maintenance trial. *J AM DIET ASSOC* 2011 Dec;111(12):1826-35.

Chang P. Effects of interviewer questions and response type on compliance: An analogue study. *Journal of Counseling Psychology* 1994 Jan;41(1):74-82.

Cheng J, Brar PS, Lee AR, Green PHR. Body mass index in celiac disease: Beneficial effect of a gluten-free diet. *J Clin Gastroenterol* 2010;44(4):267-71.

Cheong SH, McCargar LJ, Paty BW, Tudor-Locke C, Bell RC. The First Step First Bite Program: guidance to increase physical activity and daily intake of low-glycemic index foods. *J AM DIET ASSOC* 2009 Aug;109(8):1411-6.

Church TS, Earnest CP, Thompson AM, Priest EL, Rodarte RQ, Saunders T, et al. Exercise without weight loss does not reduce C-reactive protein: the INFLAME study. *Med Sci Sports Exerc* 2010 Apr;42(4):708-16.

Clark D, Chrysler L, Perkins A, Keith NR, Willis DR, Abernathy G, et al. Screening, referral, and participation in a weight management program implemented in five CHCs. *J HEALTH CARE POOR UNDERSERV* 2010 May;21(2):617-28.



Clark DO, Keith N, Chrysler L, Perkins AJ, Willis D. Design and reach of a primary care weight management program. *J HEALTH CARE POOR UNDERSERV* 2008 Feb;19(1):171-9.

Clark M, Hampson SE, Avery L, Simpson R. Effects of a tailored lifestyle self-management intervention in patients with type 2 diabetes. *Br J Health Psychol* 2004 Sep;9(Pt:3):3-79.

Cobiac L, Vos T, Veerman L. Cost-effectiveness of Weight Watchers and the Lighten Up to a Healthy Lifestyle program. *Aust NZ J Public Health* 2010 Jun;34(3):240-7.

Cocco G, Pandolfi S. Physical exercise with weight reduction lowers blood pressure and improves abnormal left ventricular relaxation in pharmacologically treated hypertensive patients. *J Clin Hypertens* 2011 Jan;13(1):23-9.

Cockcroft A, Gooch C, Ellinghouse C, Johnston M, Michie S. Evaluation of a programme of health measurements and advice among hospital staff. *Occup Med* 1994;44(2):70-6.

Collins RL, Rothblum ED, Wilson GT. The comparative efficacy of cognitive and behavioral approaches to the treatment of obesity. *Cognitive Therapy and Research* 1986 Jun;10(3):299-317.

Colvin RH, Olson SB. Winners revisited: An 18-month follow-up of our successful weight losers. *Addict Behav* 1984;9(3):305-6.

Conlon KE, Ehrlinger J, Eibach RP, Crescioni AW, Alquist JL, Gerend MA, et al. Eyes on the prize: The longitudinal benefits of goal focus on progress toward a weight loss goal. *Journal of Experimental Social Psychology* 2011 Jul;47(4):853-5.

Copeland AL, Martin PD, Geiselman PJ, Rash CJ, Kendzor DE. Smoking cessation for weight-concerned women: Group vs. individually tailored, dietary, and weight-control follow-up sessions. *Addict Behav* 2006 Jan;31(1):115-27.

Coppell KJ, Kataoka M, Williams SM, Chisholm AW, Vorgers SM, Mann JI. Nutritional intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug treatment - Lifestyle over and above drugs in diabetes (LOADD) study: Randomised controlled trial. *BMJ (Online)* 2010;341(7766):237.

Coupar AM, Kennedy T. Running a weight control group: experiences of a psychologist and a general practitioner. *J R Coll Gen Pract* 1980;30(210):41-8.

Courtney MR, Conard SE, Dunn P, Scarborough K. The Game of Health: an innovative lifestyle change program implemented in a family practice. *J AM ACAD NURSE PRACT* 2011 Jun;23(6):289-97.

Craigie AM, Macleod M, Barton KL, Treweek S, Anderson AS, WeighWell team. Supporting postpartum weight loss in women living in deprived communities: design implications for a randomised control trial. *Eur J Clin Nutr* 2011 Aug;65(8):952-8.

Croswell J, Luger S. Screening for and management of obesity in adults. *Am Fam Physician* 2012;86(10):947-8.

Cullen KW, Lara SA, Thompson D, Watson KB, Reed D, Konzelmann K. Creating healthful home food environments: results of a study with participants in the expanded food and nutrition education program. *J NUTR EDUC BEHAV* 2009 Nov;41(6):380-8.

Cullen KW, Thompson DI, Scott AR, Lara-Smallling A, Watson KB, Konzelmann K. The impact of goal attainment on behavioral and mediating variables among low income women participating in an Expanded Food and Nutrition Education Program intervention study. *Appetite* 2010 Oct;55(2):305-10.

Dahn JR, Fitzpatrick SL, Llabre MM, Apterbach GS, Helms RL, Cugnetto ML, et al. Weight management for veterans: Examining change in weight before and after MOVE. *Obesity* 2011;19(5):977-81.

Daniel EL. A multi-intervention weight management program for low-income rural women. *J AM DIET ASSOC* 1989 Sep;89(9):1310-1.

Dasgupta K, Grover SA, Da CD, Lowensteyn I, Yale JF, Rahme E. Impact of modified glucose target and exercise interventions on vascular risk factors. *DIABETES RES CLIN PRACT* 2006 Apr;72(1):53-60.

Davenport MH, Giroux I, Sopper MM, Mottola MF. Postpartum exercise regardless of intensity improves chronic disease risk factors. *Med Sci Sports Exerc* 2011 Jun;43(6):951-8.

Davies MJ HSSTCMCMC. Effectiveness of the diabetes education and self management for ongoing and newly diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: cluster randomised controlled trial. *BMJ (Clinical research ed)* 2008;336(7642):491.

Davies MJ, Heller S, Skinner TC, Campbell MJ, Carey ME, Cradock S, et al. Effectiveness of the diabetes education and self management for ongoing and newly diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: cluster randomised controlled trial. *BMJ (Clinical research ed)* 2008 Mar;336(Clinical research ed):491-5.

Davis BS, McWhirter MF, Gordon DS. Where needs and demands diverge: Health promotion in primary care. *Public Health* 1996;110(2):95-101.

Davis NJ, Crandall JP, Gajavelli S, Berman JW, Tomuta N, Wylie-Rosett J, et al. Differential effects of low-carbohydrate and low-fat diets on inflammation and endothelial function in diabetes. *Journal of Diabetes & its Complications* 2011 Nov;25(6):371-6.

de Groot LCPG, van Staveren WA. Low-protein intakes and protein turnover in elderly women. *Nutr Rev* 1996;54(2 I):58-60.

de la Rocha O. The reorganization of arithmetic practice in the kitchen. *Anthropology & Education Quarterly* 1985;16(3):193-8.

de Souza RJ, Bray GA, Carey VJ, Hall KD, LeBoff MS, Loria CM, et al. Effects of 4 weight-loss diets differing in fat, protein, and carbohydrate on fat mass, lean mass, visceral adipose tissue, and hepatic fat: results from the POUNDS LOST trial. *Am J Clin Nutr* 2012 Mar;95(3):614-25.

Dedmon RE, Kubiak MK, Konkol PD, Gander JW, Paschke AC. Employees as health educators: a reality at Kimberly-Clark. *Occupational Health & Safety* 1980 Apr 21;49(4):18-9.

DeFronzo RA, Triplitt C, Qu Y, Lewis MS, Maggs D, Glass LC. Effects of exenatide plus rosiglitazone on beta-cell function and insulin sensitivity in subjects with type 2 diabetes on metformin. *Diabetes Care* 2010;33(5):951-7.

Del CP, Bryan DR, Garvey WT, Gower BA, Hunter GR. Dietary adherence during weight loss predicts weight regain. *Obesity* 2011 Jun;19(6):1177-81.

Dennis S, Williams A, Taggart J, Newall A, Denney-Wilson E, Zwar N, et al. Which providers can bridge the health literacy gap in lifestyle risk factor modification education: a systematic review and narrative synthesis. *BMC Fam Pract* 2012;13:44.

Denolle T, Sharareh A, Dib M, Agaesse MF, Auguste V, Boutier H, et al. [Effectiveness of a health network in secondary prevention among coronary patients]. [French]. *Archives des Maladies du Coeur et des Vaisseaux* 2007 Aug;100(8):625-9.

Derosa G, D'Angelo A, Fogari E, Salvadeo S, Gravina A, Ferrari I, et al. Nateglinide and glibenclamide metabolic effects in naive type 2 diabetic patients treated with metformin. *J Clin Pharm Ther* 2009 Feb;34(1):13-23.

Derosa G, Maffioli P, Salvadeo SAT, Ferrari I, Gravina A, Mereu R, et al. Sibutramine and L-carnitine compared to sibutramine alone on insulin resistance in diabetic patients. *Intern Med* 2010;49(16):1717-25.

Derosa G, Maffioli P, Ferrari I, Palumbo I, Randazzo S, D'Angelo A, et al. Effects of one year treatment of sibutramine on insulin resistance parameters in type 2 diabetic patients. *Journal of Pharmacy & Pharmaceutical Sciences* 2010;13(3):378-90.

Derosa G, Maffioli P, Ferrari I, Palumbo I, Randazzo S, D'Angelo A, et al. Variation of inflammatory parameters after sibutramine treatment compared to placebo in type 2 diabetic patients. *Journal of Clinical Pharmacy & Therapeutics* 2011 Oct;36(5):592-601.

Derosa G, Putignano P, Bossi AC, Bonaventura A, Querci F, Franzetti IG, et al. Exenatide or glimepiride added to metformin on metabolic control and on insulin resistance in type 2 diabetic patients. *Eur J Pharmacol* 2011 Sep;666(1-3):251-6.

Di NJ, Furst G, Park K, Byrd-Bredbenner C. Designing culturally sensitive dietary interventions for African Americans: Review and recommendations. *Nutr Rev* 2013;71(4):224-38.

Dodd KJ, Duffy S, Stewart JA, Impey J, Taylor N. A small group aerobic exercise programme that reduces body weight is feasible in adults with severe chronic schizophrenia: a pilot study. *Disability & Rehabilitation* 2011;33(13-14):1222-9.

Domel SB, Alford BB, Cattlett HN, Gench BE. Weight control for black women. *J AM DIET ASSOC* 1992 Mar;92(3):346-8.

Domel SB, Alford BB, Cattlett HN, Rodriguez ML, Gench BE. A pilot weight control program for Hispanic women. *J AM DIET ASSOC* 1992;92(10):1270-1.

Donner TW, Magder LS, Zarbalian K. Dietary supplementation with d-tagatose in subjects with type 2 diabetes leads to weight loss and raises high-density lipoprotein cholesterol. *Nutr Res* 2010;30(12):801-6.

Dornelas EA WR. The DIET study: long-term outcomes of a cognitive-behavioral weight-control intervention in independent-living elders. *Dietary Intervention: Evaluation of Technology. J AM DIET ASSOC* 1998;98(11):1276.

Dostalova I, Roubicek T, Bartlova M, Mraz M, Lacinova Z, Haluzikova D, et al. Increased serum concentrations of macrophage inhibitory cytokine-1 in patients with obesity and type 2 diabetes mellitus: the influence of very low calorie diet. *EUR J ENDOCRINOL* 2009 Sep;161(3):397-404.

Dowell AC, Ochera JJ, Hilton SR, Bland JM, Harris T, Jones DR, et al. Prevention in practice: Results of a 2-year follow-up of routine health promotion interventions in general practice. *Fam Pract* 1996;13(4):357-62.

Drieling RL, Gardner CD, Ma J, Ahn DK, Stafford RS. No beneficial effects of pine bark extract on cardiovascular disease risk factors. *Arch Intern Med* 2010 Sep 27;170(17):1541-7.

Eakin E R. Maintenance of physical activity and dietary change following a telephone-delivered intervention. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association* 2010;29(6):566.

Eakin EG, RMLSO. The Logan Healthy Living Program: a cluster randomized trial of a telephone-delivered physical activity and dietary behavior intervention for primary care patients with type 2 diabetes or hypertension from a socially disadvantaged community--rationale, design and recruitment. *Contemp Clin Trials* 2008;29(3):439.

Eakin EG, Reeves MM, Marshall AL, Dunstan DW, Graves N, Healy GN, et al. Living Well with Diabetes: a randomized controlled trial of a telephone-delivered intervention for maintenance of weight loss, physical activity and glycaemic control in adults with type 2 diabetes. *BMC Public Health* 2010;10:452.

Earnest CP, Lavie CJ, Blair SN, Church TS. Heart rate variability characteristics in sedentary postmenopausal women following six months of exercise training: the DREW study. *PLoS ONE* 2008;3(6):e2288, 2008.

Egger G, Fisher G, Piers S, Bedford K, Morseau G, Sabasio S, et al. Abdominal obesity reduction in indigenous men. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 1999 Jun;23(6):564-9.

Egger G, Pearson S, Pal S, Swinburn B. Dissecting obesogenic behaviours: the development and application of a test battery for targeting prescription for weight loss.[Erratum appears in *Obes Rev*. 2007 Nov;8(6):533]. *Obes Rev* 2007 Nov;8(6):481-6.

Eiben G, Lissner L. Health Hunters--an intervention to prevent overweight and obesity in young high-risk women. *Int J Obes* 2006 Apr;30(4):691-6.

Elbel R, Aldana S, Bloswick D, Lyon JL. A pilot study evaluating a peer led and professional led physical activity intervention with blue-collar employees. *Work* 2003;21(3):199-210.

Elder C, Gallison C, Lindberg NM, DeBar L, Funk K, Ritenbaugh C, et al. Randomized trial of Tapas Acupressure Technique for weight loss maintenance: rationale and study design. *J ALTERN COMPLEMENT MED* 2010 Jun;16(6):683-90.

Elder JP, Candelaria JI, Woodruff SI, Criqui MH, Talavera GA, Rupp JW. Results of language for health: cardiovascular disease nutrition education for Latino English-as-a-second-language students. *HEALTH EDUC BEHAV* 2000 Feb;27(1):50-63.

Elliot DL, Goldberg L, Kuehl KS, Moe EL, Breger RKR, Pickering MA. The PHLAME (Promoting Healthy Lifestyles: Alternative Models' Effects) firefighter study: Outcomes of two models of behavior change. *J Occup Environ Med* 2007;49(2):204-13.

Elm CM, Wallander ID, Endrizzi B, Zelickson BD. Efficacy of a multiple diode laser system for body contouring. *Lasers in Surgery & Medicine* 2011 Feb;43(2):114-21.

Engel SG, Kolotkin RL, Teixeira PJ, Sardinha LB, Vieira PN, Palmeira AL, et al. Psychometric and cross-national evaluation of a Portuguese version of the Impact of Weight on Quality of Life-Lite (IWQOL-Lite) questionnaire. *EUR EAT DISORD REV* 2005;13(2):133-43.

Englert HS, Diehl HA, Greenlaw RL, Willich SN, Aldana S. The effect of a community-based coronary risk reduction: the Rockford CHIP. *PREV MED* 2007 Jun;44(6):513-9.

Erez G, Tirosh A, Rudich A, Meiner V, Schwarzfuchs D, Sharon N, et al. Phenotypic and genetic variation in leptin as determinants of weight regain. *Int J Obes* 2011 Jun;35(6):785-92.

Eriksson MK, HL. Quality of life and cost-effectiveness of a 3-year trial of lifestyle intervention in primary health care. *Arch Intern Med* 2010;170(16):1470.

Eriksson KF, Lindgarde F. No excess 12-year mortality in men with impaired glucose tolerance who participated in the Malmo Preventive Trial with diet and exercise. *Diabetologia* 1998 Sep;41(9):1010-6.

Eslam M, Aparcero R, Gomez MR. Changes in insulin sensitivity and body weight during and after peginterferon and ribavirin therapy for hepatitis C. *Ann Hepatol* 2011;10(2):241-3.

Esposito K, Ida MM, Ciotola M, Di PC, Scognamiglio P, Gicchino M, et al. Effects of a mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: A randomized trial. *Obstet Gynecol Surv* 2010;65(6):379-80.

Estruch R. Anti-inflammatory effects of the Mediterranean diet: The experience of the PREDIMED study. *Proc Nutr Soc* 2010;69(3):333-40.

Fallahi KM, Mazaheri M, Tamizi Z, Khankh HR, Babaei RM, Ghazanfari N, et al. The effect of weight monitoring and recording on control of obesity and overweight. *Eating & Weight Disorders: EWD* 2011 Jun;16(2):e137-e141.

Faucher MA, Mobley J. A Community Intervention on Portion Control Aimed at Weight Loss in Low-Income Mexican American Women. *J Midwifery Women's Health* 2010;55(1):60-4.

Feenstra TL, van Baal PM, Jacobs-van der Bruggen MO, Hoogenveen RT, Kommer G-J, Baan CA. Targeted versus universal prevention. A resource allocation model to prioritize cardiovascular prevention. *Cost Eff Resour Allocat* 2011;9, 2011. Article Number: 14. Date of Publication: 06 Oct 2011.

Ferrannini E, Fonseca V, Zinman B, Matthews D, Ahren B, Byiers S, et al. Fifty-two-week efficacy and safety of vildagliptin vs. glimepiride in patients with type 2 diabetes mellitus inadequately controlled on metformin monotherapy. *Diabetes Obes Metab* 2009;11(2):157-66.

Ferrer-Garcia JC, Sanchez LP, Pablos-Abella C, Albalat-Galera R, Elvira-Macagno L, Sanchez-Juan C, et al. [Benefits of a home-based physical exercise program in elderly subjects with type 2 diabetes mellitus]. *Endocrinol nutr* 2011 Oct;58(8):387-94.

Finch EA, Linde JA, Jeffery RW, Rothman AJ, King CM, Levy RL. The effects of outcome expectations and satisfaction on weight loss and maintenance: correlational and experimental analyses--a randomized trial. *Health Psychol* 2005 Nov;24(6):608-16.

Fleming RM. The effect of high-, moderate-, and low-fat diets on weight loss and cardiovascular disease risk factors.[Erratum appears in *Prev Cardiol* 2002 Fall;5(4):203]. *Preventive Cardiology* 2002;5(3):110-8.

Flocke SA, Gordon LE, Pomiecko GL. Evaluation of a community health promotion resource for primary care practices. *Am J Prev Med* 2006 Mar;30(3):243-51.

Fogel J, Shlivko S. Weight problems and spam e-mail for weight loss products. *Southern Medical Journal* 2010;103(1):31-6.

Fogelfeld L, Dharmalingam M, Robling K, Jones C, Swanson D, Jacober S. A randomized, treat-to-target trial comparing insulin lispro protamine suspension and insulin detemir in insulin-naive patients with Type 2 diabetes. *Diabet Med* 2010 Feb;27(2):181-8.

Fortmann SP, Williams PT, Hulley SB, Maccoby N, Farquhar JW. Does dietary health education reach only the privileged? The Stanford Three Community Study. *Circulation* 1982 Jul;66(1):77-82.

Foster GD, Wyatt HR, Hill JO, Makris AP, Rosenbaum DL, Brill C, et al. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet. *Obstet Gynecol Surv* 2010;65(12):769-70.

Franciosi M, Lucisano G, Pellegrini F, Cantarello A, Consoli A, Cucco L, et al. ROSES: Role of self-monitoring of blood glucose and intensive education in patients with Type2 diabetes not receiving insulin. A pilot randomized clinical trial. *Diabet Med* 2011;28(7):789-96.

Friedl KE, DeWinne CM, Taylor RL. The use of the Durnin-Womersley generalized equations for body fat estimation and their impact on the Army Weight Control Program. *Mil Med* 1987 Mar;152(3):150-5.

Fu P-K, Hsu H-Y, Wang P-Y. Transient global amnesia after taking sibutramine: A case report. *Neurologist* 2010;16(2):129-31.

Fuchs FD, Fuchs SC, Moreira LB, Gus M, Nobrega AC, Poli-de-Figueiredo CE, et al. Prevention of hypertension in patients with pre-hypertension: Protocol for the PREVER-prevention trial. *Trials* 2011;12, 2011. Article Number: 65. Date of Publication: 05 Mar 2011.

Funk KL, Stevens VJ, Appel LJ, Bauck A, Brantley PJ, Champagne CM, et al. Associations of internet website use with weight change in a long-term weight loss maintenance program. *J Med Internet Res* 2010;12(3):e29.

Funk KL, Stevens VJ, Bauck A, Brantley PJ, Hornbrook M, Jerome GJ, et al. Development and implementation of a tailored self-assessment tool in an internet-based weight loss maintenance program. *Clin Pract Epidemiol Ment Health* 2011;7:67-73.

Gades MD, Stern JS. Chitosan supplementation does not affect fat absorption in healthy males fed a high-fat diet, a pilot study. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 2002 Jan;26(1):119-22.

Garber A, Henry RR, Ratner R, Hale P, Chang CT, Bode B. Liraglutide, a once-daily human glucagon-like peptide 1 analogue, provides sustained improvements in glycaemic control and weight for 2 years as monotherapy compared with glimepiride in patients with type 2 diabetes. *Diabetes Obes Metab* 2011;13(4):348-56.

Gardner CD. Low-carbohydrate ketogenic diet and the combination of orlistat with a low-fat diet lead to comparable improvements in weight and blood lipids, but LCKD more beneficial for blood pressure. *Evid -Based Med* 2010;15(3):91-2.

Garry JP, Whetstone LM, Morrissey SL. Progress toward Healthy People 2000. Physical activity and weight status in eastern North Carolina. *N C Med J* 2002 Jan;63(1):6-10.

Gerber BS, Stolley MR, Thompson AL, Sharp LK, Fitzgibbon ML. Mobile phone text messaging to promote healthy behaviors and weight loss maintenance: a feasibility study. *Health Informatics Journal* 2009 Mar;15(1):17-25.

Gifford EV, Lillis J. Avoidance and inflexibility as a common clinical pathway in obesity and smoking treatment. [References]. *J HEALTH PSYCHOL* 2009 Oct;14(7):992-6.

Gilbert J-A, Gasteyger C, Raben A, Meier DH, Astrup A, Sjodin A. The effect of tesofensine on appetite sensations. *Obesity* 2012;20(3):553-61.

Gill A, Hoogwerf BJ, Burger J, Bruce S, MacConnell L, Yan P, et al. Effect of exenatide on heart rate and blood pressure in subjects with type 2 diabetes mellitus: A double-blind, placebo-controlled,

randomized pilot study. *Cardiovasc diabetol* 2010;9 , 2010. Article Number: 6. Date of Publication: 28 Jan 2010.

Gill K, Overdorf V. Incentives for exercise in younger and older women. *Journal of Sport Behavior* 1994 Jun;17(2):87-97.

Gillen LJ, Tapsell LC, Patch CS, Owen A, Batterham M. Structured dietary advice incorporating walnuts achieves optimal fat and energy balance in patients with type 2 diabetes mellitus. *J AM DIET ASSOC* 2005 Jul;105(7):1087-96.

Gilstrap LG, Malhotra R, Peltier-Saxe D, Slicas D, Pineda E, Culhane-Hermann C, et al. Community-based primary prevention programs decrease the rate of metabolic syndrome among socioeconomically disadvantaged women. *J Women's Health* 2013;22(4):322-9.

Gittelsohn J, Dyckman W, Tan ML, Boggs MK, Frick KD, Alfred J, et al. Development and implementation of a food store-based intervention to improve diet in the Republic of the Marshall Islands. *HEALTH PROMOT PRACT* 2006 Oct;7(4):396-405.

Gittelsohn J, Song HJ, Suratkar S, Kumar MB, Henry EG, Sharma S, et al. An urban food store intervention positively affects food-related psychosocial variables and food behaviors. *HEALTH EDUC BEHAV* 2010 Jun;37(3):390-402.

Glanz K, Murphy S, Moylan J, Evensen D, Curb JD. Improving dietary self-monitoring and adherence with hand-held computers: A pilot study. [References]. *Am J Health Promot* 2006 Jan;20(3):165-70.

Glasgow RE KD. Twelve-month outcomes of an Internet-based diabetes self-management support program. *Patient Education and Counseling* 2012;87(1):81.

Godin G, Belanger-Gravel A, Amireault S, Gallani MC, Vohl MC, Perusse L. Effect of implementation intentions to change behaviour: moderation by intention stability. *Psychol Rep* 2010 Feb;106(1):147-59.

Goetzel RZ, Kahr TY, Aldana SG, Kenny GM. An evaluation of Duke University's live for life health promotion program and its impact on employee health. *Am J Health Promot* 1996;10(5):340-2.

Goke B, Hershon K, Kerr D, Calle PA, Schweizer A, Foley J, et al. Efficacy and safety of vildagliptin monotherapy during 2-year treatment of drug-naive patients with type 2 diabetes: comparison with metformin. *Horm Metab Res* 2008 Dec;40(12):892-und.

Gold DB, Anderson DR, Serxner SA. Impact of a telephone-based intervention on the reduction of health risks. *Am J Health Promot* 2000;15(2):97-106.

Gorin AA, Wing RR, Fava JL, Jakicic JM, Jeffery R, West DS, et al. Weight loss treatment influences untreated spouses and the home environment: evidence of a ripple effect. *Int J Obes (Lond)* 2008 Nov;32(2005):1678-84.

Goyer L, Dufour R, Janelle C, Blais C, L'Abbe C, Raymond E, et al. Randomized controlled trial on the long-term efficacy of a multifaceted, interdisciplinary lifestyle intervention in reducing cardiovascular risk and improving lifestyle in patients at risk of cardiovascular disease. *J Behav Med* 2013;36(2):212-24.

Gracey M, Bridge E, Martin D, Jones T, Spargo RM, Shephard M, et al. An Aboriginal-driven program to prevent, control and manage nutrition-related "lifestyle" diseases including diabetes. *Asia Pac J Clin Nutr* 2006;15(2):178-88.

Grammatikopoulou MG, Panayiotoglou A, Hassapidou M. Evaluation of commercial weight-loss programmes in Greece. [References]. *International Journal of Consumer Studies* 2008 Jan;32(1):59-64.

Greene GW, Schembre SM, White AA, Hoerr SL, Lohse B, Shoff S, et al. Identifying clusters of college students at elevated health risk based on eating and exercise behaviors and psychosocial determinants of body weight. *J AM DIET ASSOC* 2011 Mar;111(3):394-400.

Greene LF, Malpede CZ, Henson CS, Hubbert KA, Heimburger DC, Ard JD. Weight maintenance 2 years after participation in a weight loss program promoting low-energy density foods. *Obesity* 2006 Oct;14(10):1795-801.

Greenwood TC. Comparison of a low carbohydrate and calorie-restricted diet on weight loss, morphometric variables, and diet acceptability. *Dissertation Abstracts International: Section B: The Sciences and Engineering* 2009;69(8-B):2009-B2009.

Greer DM, Hill DC. Implementing an evidence-based metabolic syndrome prevention and treatment program utilizing group visits. *J AM ACAD NURSE PRACT* 2011;23(2):76-83.

Griffin HJ, Cheng HL, O'Connor HT, Rooney KB, Petocz P, Steinbeck KS. Higher protein diet for weight management in young overweight women: A 12-month randomized controlled trial. *Diabetes Obes Metab* 2013;15(6):572-5.

Grover SA LI. Patient knowledge of coronary risk profile improves the effectiveness of dyslipidemia therapy: the CHECK-UP study: a randomized controlled trial. *Arch Intern Med* 2007;167(21):2296.

Guimaraes ACA, Baptista F. Influence of habitual physical activity on the symptoms of climacterium/menopause and the quality of life of middle-aged women. *Int J Womens Health* 2011;3(1):319-28.

Gupta AK, Smith SR, Greenway FL, Bray GA. Pioglitazone treatment in type 2 diabetes mellitus when combined with portion control diet modifies the metabolic syndrome. *Diabetes Obes Metab* 2009;11(4):330-7.

Gupta AK, Bray GA, Greenway FL, Martin CK, Johnson WD, Smith SR. Pioglitazone, but not metformin, reduces liver fat in Type-2 diabetes mellitus independent of weight changes. *Journal of Diabetes & its Complications* 2010 Sep;24(5):289-96.

Gusi N, Reyes MC, Gonzalez-Guerrero JL, Herrera E, Garcia JM. Cost-utility of a walking programme for moderately depressed, obese, or overweight elderly women in primary care: a randomised controlled trial. *BMC Public Health* 2008;8:231.

Gutschall MD, Miller CK, Mitchell DC, Lawrence FR. A randomized behavioural trial targeting glycaemic index improves dietary, weight and metabolic outcomes in patients with type 2 diabetes. *Public Health Nutr* 2009;12(10):1846-54.

Haglin L. All-cause mortality of patients with dyslipidemia up to 19 years after a multidisciplinary lifestyle modification programme: a randomized trial. *Eur J Cardiovasc Prev Rehabil* 2011;18(1):79.

Hall PA, Fong GT, Cheng AY. Time perspective and weight management behaviors in newly diagnosed type 2 diabetes: A mediational analysis. *J Behav Med* 2012;35(6):569-80.

Halm MA, Denker J. Primary prevention programs to reduce heart disease risk in women. *Clin Nurse Spec* 2003;17(2):101.

Hamdy O, Carver C. The Why WAIT program: improving clinical outcomes through weight management in type 2 diabetes. *Curr Diab Rep* 2008 Oct;8(5):413-20.

Hamman RF, Wing RR, Edelstein SL, Lachin JM, Bray GA, Delahanty L, et al. Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care* 2006;29(9):2102-7.



Hankonen N, Vollmann M, Renner B, Absetz P. What is setting the stage for abdominal obesity reduction? A comparison between personality and health-related social cognitions. [References]. *J Behav Med* 2010 Oct;33(5):415-22.

Hardcastle S, Taylor A, Bailey M, Castle R. A randomised controlled trial on the effectiveness of a primary health care based counselling intervention on physical activity, diet and CHD risk factors. *PATIENT EDUC COUNS* 2008 Jan;70(1):31-9.

Hardcastle SJ, Taylor AH, Bailey MP, Harley RA, Hagggar MS. Effectiveness of a motivational interviewing intervention on weight loss, physical activity and cardiovascular disease risk factors: A randomised controlled trial with a 12-month post-intervention follow-up. *Int j behav nutr phys act* 2013;10.

Harman NL, Leeds AR, Griffin BA. Increased dietary cholesterol does not increase plasma low density lipoprotein when accompanied by an energy-restricted diet and weight loss.[Erratum appears in *Eur J Nutr*. 2008 Oct;47(7):408]. *EUR J NUTR* 2008 Sep;47(6):287-93.

Harris JK, French SA, Jeffery RW, McGovern PG, Wing RR. Dietary and physical activity correlates of long-term weight loss. *Obes Res* 1994 Jul;2(4):307-13.

Hartman TJ, Zhang Z, Albert PS, Bagshaw D, Mentor-Marcel R, Mitchell DC, et al. Reduced energy intake and weight loss on a legume-enriched diet lead to improvements in biomarkers related to chronic disease. *Top Clin Nutr* 2011;26(3):208-15.

Hawkins MR, Cooke AJ, Major JE. An evaluation of the weight control program at a U.S. Army installation. *Mil Med* 1986 Mar;151(3):185-6.

Heald AH, Knapman H, Nair S, Chambers T, Radford D, Rushton T, et al. A primary care register for impaired glucose handling (IGH): Impact on cardiometabolic profile. *Prim Care Diabetes* 2012;6(3):213-9.

Heath GW, Parra DC, Sarmiento OL, Andersen LB, Owen N, Goenka S, et al. Evidence-based intervention in physical activity: Lessons from around the world. *The Lancet* 2012;379(9838):272-81.

Heideman WH, Nierkens V, Stronks K, Middelkoop BJ, Twisk JW, Verhoeff AP, et al. DiAlert: a lifestyle education programme aimed at people with a positive family history of type 2 diabetes and overweight, study protocol of a randomised controlled trial. *BMC Public Health* 2011;11:751.

Heinberg LJ, Haythornthwaite JA, Rosofsky W, McCarron P, Clarke A. Body image and weight loss maintenance in elderly African American hypertensives. [References]. *Am J Health Behav* 2000 May;24(3):163-73.

Hendricks EJ, Greenway FL. A study of abrupt phentermine cessation in patients in a weight management program. *American Journal of Therapeutics* 2011 Jul;18(4):292-9.

Herder C, Peltonen M, Koenig W, Sutfels K, Lindstrom J, Martin S, et al. Anti-inflammatory effect of lifestyle changes in the Finnish Diabetes Prevention Study. *Diabetologia* 2009;52(3):433-42.

Higgins LC, Gray W. Changing the body image concern and eating behaviour of chronic dieters: The effects of a psychoeducational intervention. *PSYCHOL HEALTH* 1998 Nov;13(6):1045-60.

Hill DH, Madison R. A health education program for weight reduction in a hypertension clinic. *PUBLIC HEALTH REP* 1980 May;95(3):271-5.

Hill JO, Wyatt HR, Peters JC. Energy balance and obesity. *Circulation* 2012;126(1):126-32.

Hiramitsu S, Ishiguro Y, Matsuyama H, Yamada K, Kato K, Noba M, et al. The effects of ezetimibe on surrogate markers of cholesterol absorption and synthesis in Japanese patients with dyslipidemia. *J Atheroscler Thromb* 2010;17(1):106-14.

Hjermann I, Leren P, Norman N. Serum insulin response to oral glucose load during a dietary intervention trial in healthy coronary high risk men: The Oslo Study. *Scand J Clin Lab Invest* 1980;40(1):89-94.

Hochstenbach-Waelen A, Westerterp KR, Soenen S, Westerterp-Plantenga MS. No long-term weight maintenance effects of gelatin in a supra-sustained protein diet. *Physiol Behav* 2010 Sep 1;101(2):237-44.

Hochstenbach-Waelen A, Soenen S, Westerterp KR, Westerterp-Plantenga MS. Effects of a supra-sustained gelatin-milk protein diet compared with (supra-)sustained milk protein diets on body-weight loss. *Br J Nutr* 2011 May;105(9):1388-98.

Hollander P, Cooper J, Bregnhøj J, Pedersen CB. A 52-week, multinational, open-label, parallel-group, noninferiority, treat-to-target trial comparing insulin detemir with insulin glargine in a basal-bolus regimen with mealtime insulin aspart in patients with type 2 diabetes. *Clin Ther* 2008 Nov;30(11):1976-87.

Hollander PA, Amod A, Litwak LE, Chaudhari U, ARPEGGIO Study Group. Effect of rimonabant on glycemic control in insulin-treated type 2 diabetes: the ARPEGGIO trial. *Diabetes Care* 2010 Mar;33(3):605-7.

Holmes AL, Sanderson B, Maisiak R, Brown A, Bittner V. Dietitian services are associated with improved patient outcomes and the MEDFICTS dietary assessment questionnaire is a suitable outcome measure in cardiac rehabilitation. *J AM DIET ASSOC* 2005;105(10):1533-40.

Houlihan MM, Dickson-Parnell BE, Jackson J, Zeichner A. Appearance changes associated with participation in a behavioral weight control program. *Addict Behav* 1987;12(2):157-63.

Hovell MF, Mulvihill MM, Buono MJ, Liles S, Schade DH, Washington TA, et al. Culturally tailored aerobic exercise intervention for low-income Latinas. *Am J Health Promot* 2008 Jan;22(3):155-63.

Hoy MK, Winters BL, Chlebowski RT, Papoutsakis C, Shapiro A, Lubin MP, et al. Implementing a low-fat eating plan in the Women's Intervention Nutrition Study. *J AM DIET ASSOC* 2009 Apr;109(4):688-96.

Huffman KM, Redman LM, Landerman LR, Pieper CF, Stevens RD, Muehlbauer MJ, et al. Caloric restriction alters the metabolic response to a mixed-meal: Results from a randomized, controlled trial. *PLoS ONE* 2012;7(4).

Hunter GR, Brock DW, Byrne NM, Chandler-Laney PC, Del CP, Gower BA. Exercise training prevents regain of visceral fat for 1 year following weight loss. *Obesity* 2010 Apr;18(4):690-5.

Ibanez L, Lopez-Bermejo A, Diaz M, Enriquez G, Del RL, De ZF. Low-dose pioglitazone and low-dose flutamide added to metformin and oestro-progestagens for hyperinsulinaemic women with androgen excess: add-on benefits disclosed by a randomized double-placebo study over 24 months. *Clin Endocrinol (Oxf)* 2009 Sep;71(3):351-7.

Ilanne-Parikka P, Laaksonen DE, Eriksson JG, Lakka TA, Lindstrom J, Peltonen M, et al. Leisure-time physical activity and the metabolic syndrome in the Finnish diabetes prevention study. *Diabetes Care* 2010;33(7):1610-7.

Ippoliti F, Liguori A, Petti F, Canitano N, Rughini S. Leptin, ghrelin and TNF-alpha before and after hypo-caloric traditional Chinese diet and auricular acupuncture. *J Tradit Chin Med* 2008;28(1):24-29.

Irving BA, Nair KS, Srinivasan M. Effects of insulin sensitivity, body composition, and fitness on lipoprotein particle sizes and concentrations determined by nuclear magnetic resonance. *J Clin Endocrinol Metab* 2011;96(4):E713-E718.

Isaakidis P, Raguenaud M-E, Say C, De CH, Khim C, Pottier R, et al. Treatment of hypertension in rural Cambodia: Results from a 6-year programme. *J Hum Hypertens* 2011;25(4):241-9.

Izquierdo R, Laguna CT, Meyer S, Ploutz-Snyder RJ, Palmas W, Eimicke JP, et al. Telemedicine intervention effects on waist circumference and body mass index in the IDEATel project. *DIABETES TECHNOLOG THER* 2010 Mar;12(3):213-20.

Jackson EG. Eating order: a 13-week trust model class for dieting casualties. *J NUTR EDUC BEHAV* 2008 Jan;40(1):43-8.

Jacobs DR, Sluik D, Rokling-Andersen MH, Anderssen SA, Drevon CA. Association of 1-y changes in diet pattern with cardiovascular disease risk factors and adipokines: results from the 1-y randomized Oslo Diet and Exercise Study. *The American journal of clinical nutrition* 2009 Feb;89(2):509-17.

Jansink R, Braspenning J, Keizer E, van der Weijden T, Elwyn G, Grol R. No identifiable Hb1Ac or lifestyle change after a comprehensive diabetes programme including motivational interviewing: a cluster randomised trial. *SCAND J PRIM HEALTH CARE* 2013;31(2):119-27.

Jason LA, Greiner BJ, Naylor K, Johnson SP, Van EL. A large-scale, short-term, media-based weight loss program. *Am J Health Promot* 1991 Jul;5(6):432-7.

Jay M. Impact of VA weight management program for veterans. *J Clin Outcomes Manage* 2011;18(7):12-5.

Jeffery RW, Hellerstedt WL, Schmid TL. Correspondence programs for smoking cessation and weight control: a comparison of two strategies in the Minnesota Heart Health Program. *Health Psychol* 1990;9(5):585-98.

Jeffery RW, French SA. Preventing weight gain in adults: the pound of prevention study. *Am J Public Health* 1999 May;89(5):747-51.

Jeffery RW, Linde JA, Finch EA, Rothman AJ, King CM. A satisfaction enhancement intervention for long-term weight loss. *Obesity* 2006 May;14(5):863-9.

Jeffery RW. Financial incentives and weight control. *PREV MED* 2012;55(SUPPL.):S61-S67.

Jeffery RW. Behavioral treatment of obesity with monetary contracting: Two-year follow-up. *Addict Behav* 1984;9(3):311-3.

Jeffery RW, French SA, Schmid TL. Attributions for dietary failures: Problems reported by participants in the Hypertension Prevention Trial. [References]. *Health Psychol* 1990;9(3):315-29.

Jenkins DJ, Kendall CW, Faulkner DA, Kemp T, Marchie A, Nguyen TH, et al. Long-term effects of a plant-based dietary portfolio of cholesterol-lowering foods on blood pressure. *Eur J Clin Nutr* 2008 Jun;62(6):781-8.

Jenkins DJ, Chiavaroli L, Wong JM, Kendall C, Lewis GF, Vidgen E, et al. Adding monounsaturated fatty acids to a dietary portfolio of cholesterol-lowering foods in hypercholesterolemia. *CMAJ* :

Canadian Medical Association journal = journal de l'Association medicale canadienne 2010 Dec;182(18):1961-7.

Jenkins DJ, Srichaikul K, Wong JM, Kendall CW, Bashyam B, Vidgen E, et al. Supplemental barley protein and casein similarly affect serum lipids in hypercholesterolemic women and men. *J Nutr* 2010 Sep;140(9):1633-7.

Jenum AK, Lorentzen C, Anderssen SA, Birkeland KI, Holme I, Lund-Larsen PG, et al. Promoting physical activity in a multi-ethnic district - Methods and baseline results of a pseudo-experimental intervention study. *Eur J Cardiovasc Prev Rehabil* 2003;10(5):387-96.

Jilcott SB KTS-HCR. Linking clinical care to community resources for cardiovascular disease prevention: the North Carolina Enhanced WISEWOMAN project. *J Womens Health (Larchmt)* 2006;15(5):569.

Johansen KS, Bjorge B, Hjellset VT, Holmboe-Ottesen G, Raberg M, Wandel M. Changes in food habits and motivation for healthy eating among Pakistani women living in Norway: results from the InnvaDiab-DEPLAN study. *Public Health Nutr* 2010 Jun;13(6):858-67.

John LK, Loewenstein G, Troxel AB, Norton L, Fassbender JE, Volpp KG. Financial incentives for extended weight loss: a randomized, controlled trial. *J Gen Intern Med* 2011 Jun;26(6):621-6.

Johnson F, Wardle J. The association between weight loss and engagement with a web-based food and exercise diary in a commercial weight loss programme: A retrospective analysis. *International Journal of Behavioral Nutrition and Physical Activity* 2011;8, 2011. Article Number: 83. Date of Publication: 02 Aug 2011.

Johnson HM, Turke TL, Grossklaus M, Dall T, Carimi S, Koenig LM, et al. Effects of an office-based carotid ultrasound screening intervention. *Journal of the American Society of Echocardiography* 2011;24(7):738-47.

Johnston CS, Trier CM, Fleming KR. The effect of peanut and grain bar preloads on postmeal satiety, glycemia, and weight loss in healthy individuals: An acute and a chronic randomized intervention trial. *Nutr J* 2013;12(1).

Jones A, Davies DH, Dove JR, Collinson MA, Brown PMR. Identification and treatment of risk factors for coronary heart disease in general practice: A possible screening model. *Br Med J* 1988;296(6638):1711-4.

Jorgensen T. A randomized non-pharmacological intervention study for prevention of ischaemic heart disease: baseline results Inter99. *Eur J Cardiovasc Prev Rehabil* 2003;10(5):377.

Juarascio A, Forman E, Timko CA, Butryn M, Goodwin C. The development and validation of the food craving acceptance and action questionnaire (FAAQ). *Eat behav* 2011;12(3):182-7.

Jupp JJ, Collins JK, McCabe MP, Walker WL. Hypnotic susceptibility and depth: Predictors of outcome in a weight control therapy. *Australian Journal of Clinical & Experimental Hypnosis* 1986 May;14(1):31-40.

Kabbani TA, Goldberg A, Kelly CP, Pallav K, Tariq S, Peer A, et al. Body mass index and the risk of obesity in coeliac disease treated with the gluten-free diet. *Aliment Pharmacol Ther* 2012;35(6):723-9.

Kahn SE, Haffner SM, Viberti G, Herman WH, Lachin JM, Kravitz BG, et al. Rosiglitazone decreases C-reactive protein to a greater extent relative to glyburide and metformin over 4 years despite greater weight gain: Observations from A Diabetes Outcome Progression Trial (ADOPT). *Diabetes Care* 2010;33(1):177-83.

Kattelman KK, Conti K, Ren C. The medicine wheel nutrition intervention: a diabetes education study with the Cheyenne River Sioux Tribe.[Reprint in J Am Diet Assoc. 2010 May;110(5 Suppl):S44-51; PMID: 20399298]. J AM DIET ASSOC 2009 Sep;109(9):1532-9.

Keele-Smith R, Leon T. Evaluation of Individually Tailored Interventions on Exercise Adherence. [References]. WEST J NURS RES 2003 Oct;25(6):623-40.

Kelders SM, van Gemert-Pijnen JEW, Werkman A, Seydel ER. Evaluation of a web-based lifestyle coach designed to maintain a healthy bodyweight. [References]. J Telemed Telecare 2010;16(1):3-7.

Keller C, Trevino RP. Effects of two frequencies of walking on cardiovascular risk factor reduction in Mexican American women. RES NURS HEALTH 2001 Oct;24(5):390-401.

Keller C, Records K, Ainsworth B, Belyea M, Permana P, Coonrod D, et al. Madres para la Salud: design of a theory-based intervention for postpartum Latinas. Contemp Clin Trials 2011 May;32(3):418-27.

Keller CS, Cantue A. Camina por Salud: walking in Mexican-American women. Appl Nurs Res 2008 May;21(2):110-3.

Kellow N. Evaluation of a rural community pharmacy-based Waist Management Project: bringing the program to the people. Aust J Prim Health 2011;17(1):16-22.

Kempf K, Kruse J, Martin S. ROSSO-in-praxi: A self-monitoring of blood glucose-structured 12-week lifestyle intervention significantly improves glucometabolic control of patients with type 2 diabetes mellitus. Diabetes Technol Ther 2010;12(7):547-53.

Kennedy BM, Champagne CM, Ryan DH, Newton R, Jr., Conish BK, Harsha DW, et al. The "Rolling Store:" an economical and environmental approach to the prevention of weight gain in African American women. Ethn Dis 2009;19(1):7-12.

Kenya S, He Q, Fullilove R, Kotler DP. Developing an objective evaluation method to estimate diabetes risk in community-based settings. Diabetes Technol Ther 2011;13(5):557-61.

Ketola E, Sipila R, Makela M. Effectiveness of individual lifestyle interventions in reducing cardiovascular disease and risk factors. Ann Med 2000;32(4):239-51.

King AC, Kiernan M, Oman RF, Kraemer HC, Hull M, Ahn D. Can we identify who will adhere to long-term physical activity? Signal detection methodology as a potential aid to clinical decision making. [References]. Health Psychol 1997 Jul;16(4):380-9.

Kirk CC, Griffey DC. The effects of imagery and language cognitive strategies on dietary intake, weight loss, and perception of food. Imagination, Cognition and Personality 1995;15(2):145-57.

Koelewijn-van Loon MS vdW. Involving patients in cardiovascular risk management with nurse-led clinics: a cluster randomized controlled trial. CMAJ: Canadian Medical Association Journal 2009;181(12):E267.

Kohler B, Andreen I. Body weight maintenance and levels of mutans streptococci and lactobacilli in a group of Swedish women seven years after completion of a Weight Watchers' diet. Swedish Dental Journal 2011;35(3):101-9.

Kohut ML, McCann DA, Russell DW, Konopka DN, Cunnick JE, Franke WD, et al. Aerobic exercise, but not flexibility/resistance exercise, reduces serum IL-18, CRP, and IL-6 independent of beta-blockers, BMI, and psychosocial factors in older adults. [References]. Brain Behav Immun 2006 May;20(3):201-9.

Koioy E, Tziomalos K, Dinas K, Katsikis I, Kalaitzakis E, Delkos D, et al. The effect of weight loss and treatment with metformin on serum vaspin levels in women with polycystic ovary syndrome. *Endocr J* 2011;58(4):237-46.

Koioy E, Tziomalos K, Katsikis I, Kandaraki EA, Kalaitzakis E, Delkos D, et al. Weight loss significantly reduces serum lipocalin-2 levels in overweight and obese women with polycystic ovary syndrome. *Gynecol Endocrinol* 2012;28(1):20-4.

Koutsari C, Ali AH, Nair KS, Rizza RA, O'Brien P, Khosla S, et al. Fatty acid metabolism in the elderly: Effects of dehydroepiandrosterone and testosterone replacement in hormonally deficient men and women. *J Clin Endocrinol Metab* 2009;94(9):3414-23.

Krakoff J, Clark JM, Crandall JP, Wilson C, Molitch ME, Brancati FL, et al. Effects of metformin and weight loss on serum alanine aminotransferase activity in the diabetes prevention program. *Obesity* 2010 Sep;18(9):1762-7.

Kramer MK, McWilliams JR, Chen HY, Siminerio LM. A community-based diabetes prevention program: evaluation of the group lifestyle balance program delivered by diabetes educators. *DIABETES EDUC* 2011 Sep;37(5):659-68.

Kramer MK, Kriska AM, Venditti EM, Semler LN, Miller RG, McDonald T, et al. A novel approach to diabetes prevention: evaluation of the Group Lifestyle Balance program delivered via DVD. *Diabetes Res Clin Pract* 2010;90(3):e60-e63.

Krousel-Wood MA, Berger L, Jiang X, Blonde L, Myers L, Webber L. Does home-based exercise improve body mass index in patients with type 2 diabetes? Results of a feasibility trial. *Diabetes Res Clin Pract* 2008 Feb;79(2):230-6.

Kruskall LJ, Campbell WW, Evans WJ. The Yale Physical Activity Survey for older adults: predictions in the energy expenditure due to physical activity. *J AM DIET ASSOC* 2004 Aug;104(8):1251-7.

Kuller LH, Pettee Gabriel KK, Kinzel LS, Underwood DA, Conroy MB, Chang Y, et al. The Women on the Move through Activity and Nutrition (WOMAN) study: Final 48-month results. *Obesity* 2012;20(3):636-43.

Kuller LH, Kinzel LS, Pettee KK, Kriska AM, Simkin-Silverman LR, Conroy MB, et al. Lifestyle intervention and coronary heart disease risk factor changes over 18 months in postmenopausal women: the Women On the Move through Activity and Nutrition (WOMAN study) clinical trial. *J Womens Health (Larchmt)* 2006;15(8):962-74.

Kumanyika SK, Shults J, Fassbender J, Whitt MC, Brake V, Kallan MJ, et al. Outpatient weight management in African-Americans: The Healthy Eating and Lifestyle Program (HELP) study. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 2005;41(2):488-502.

Kurukulasuriya LR, Sowers JR. Therapies for type 2 diabetes: Lowering HbA1c and associated cardiovascular risk factors. *Cardiovasc diabetol* 2010;9, 2010. Article Number: 45. Date of Publication: 30 Aug 2010.

La Londe MA, Graffagnino CL, Falko JM, Snow RJ, Spencer K, Caulin-Glaser T. Effect of a weight management program on the determinants and prevalence of metabolic syndrome. *Obesity* 2008 Mar;16(3):637-42.

Laaksonen DE, Lindstrom J, Lakka TA, Eriksson JG, Niskanen L, Wikstrom K, et al. Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study. *Diabetes* 2005;54(1):158-65.

Laatikainen T, Dunbar JA, Chapman A, Kilkkinen A, Vartiainen E, Heistaro S, et al. Prevention of type 2 diabetes by lifestyle intervention in an Australian primary health care setting: Greater Green Triangle (GGT) Diabetes Prevention Project. *BMC Public Health* 2007;7:249.

Laatikainen T, Philpot B, Hankonen N, Sippola R, Dunbar JA, Absetz P, et al. Predicting changes in lifestyle and clinical outcomes in preventing diabetes: the Greater Green Triangle Diabetes Prevention Project. *PREV MED* 2012;54(2):157-61.

Ladson G, Dodson WC, Sweet SD, Archibong AE, Kunselman AR, Demers LM, et al. The effects of metformin with lifestyle therapy in polycystic ovary syndrome: A randomized double-blind study. *Fertil Steril* 2011;95(3):1059-66.

Lambers S, Van LC, Van AK, Calders P. Influence of combined exercise training on indices of obesity, diabetes and cardiovascular risk in type 2 diabetes patients. *Clin Rehabil* 2008 Jun;22(6):483-92.

Lampman RM, Schteingart DE. Effects of exercise training on glucose control, lipid metabolism, and insulin sensitivity in hypertriglyceridemia and non-insulin dependent diabetes mellitus. *Med Sci Sports Exerc* 1991;23(6):703-12.

Lane W, Weinrib S, Rappaport J. The effect of liraglutide added to U-500 insulin in patients with type 2 diabetes and high insulin requirements. *DIABETES TECHNOL THER* 2011 May;13(5):592-5.

Larson DE, Ferraro RT, Robertson DS, Ravussin E. Energy metabolism in weight-stable postobese individuals. *Am J Clin Nutr* 1995 Oct;62(4):735-9.

Latimer L, Walker L, Sunghun K, Pasch K, Sterling B. Self-efficacy Scale for Weight Loss among Multi-ethnic Women of Lower Income: A Psychometric Evaluation. *J NUTR EDUC BEHAV* 2011 Jul;43(4):279-83.

Latka RN, Alvarez-Reeves M, Cadmus L, Irwin ML. Adherence to a randomized controlled trial of aerobic exercise in breast cancer survivors: the Yale exercise and survivorship study. *J cancer surviv* 2009 Sep;3(3):148-57.

Lau C, Vistisen D, Toft U, Tetens I, Glumer C, Pedersen O, et al. The effects of adding group-based lifestyle counselling to individual counselling on changes in plasma glucose levels in a randomized controlled trial: the Inter99 study. *Diabetes Metab* 2011 Dec;37(6):546-52.

Lee IS, Shin G, Choue R. A 12-week regimen of caloric restriction improves levels of adipokines and pro-inflammatory cytokines in Korean women with BMIs greater than 23 kg/m<sup>2</sup>. *Inflamm Res* 2010;59(5):399-405.

Lee RE, Medina AV, Mama SK, Reese-Smith JY, O'Connor DP, Brosnan M, et al. Health is Power: an ecological, theory-based health intervention for women of color. [Review]. *Contemp Clin Trials* 2011 Nov;32(6):916-23.

Lefevre M, Redman LM, Heilbronn LK, Smith JV, Martin CK, Rood JC, et al. Caloric restriction alone and with exercise improves CVD risk in healthy non-obese individuals. *Atherosclerosis* 2009 Mar;203(1):206-13.

Leijon ME, Bendtsen P, Nilsen P, Ekberg K, Stahle A. Physical activity referrals in Swedish primary health care - prescriber and patient characteristics, reasons for prescriptions, and prescribed activities. *BMC Health Serv Res* 2008;8:201.

Lenz TL, Petersen K, Monaghan MS. Counseling patients about lifestyle modification. *U S Pharm* 2008;33(1):38-45.

Leslie WS, Lean ME, Baillie HM, Hankey CR. Weight management: a comparison of existing dietary approaches in a work-site setting. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 2002 Nov;26(11):1469-75.

Levit S, Toledano Y, Wainstein J. Improved glycaemic control with reduced hypoglycaemic episodes and without weight gain using long-term modern premixed insulins in type 2 diabetes. *INT J CLIN PRACT* 2011;65(2):165-71.

Levitsky DA, Pacanowski C. Losing weight without dieting. Use of commercial foods as meal replacements for lunch produces an extended energy deficit. *Appetite* 2011 Oct;57(2):311-7.

Lewis AL, Eves FF. Specific effects of a calorie-based intervention on stair climbing in overweight commuters. *Ann Behav Med* 2011 Oct;42(2):257-61.

Li G. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* 2008;371(9626):1783.

Li SC, Liu YH, Liu JF, Chang WH, Chen CM, Chen CY. Almond consumption improved glycemic control and lipid profiles in patients with type 2 diabetes mellitus. *Metabolism* 2011 Apr;60(4):474-9.

Liao L, Tian YJ, Zhao JJ, Xin Y, Xing HY, Dong JJ. Metformin versus metformin plus rosiglitazone in women with polycystic ovary syndrome. *Chin Med J* 2011 Mar;124(5):714-8.

Lin P-H, Wang Y, Grambow SC, Goggins W, Almirall D. Dietary saturated fat intake is negatively associated with weight maintenance among the PREMIER participants. *Obesity* 2012;20(3):571-5.

Lind M, Jendle J, Torffvit O, Lager I. Glucagon-like peptide 1 (GLP-1) analogue combined with insulin reduces HbA1c and weight with low risk of hypoglycemia and high treatment satisfaction. *Prim Care Diabetes* 2012;6(1):41-6.

Linde JA, Jeffery RW. Testing a brief self-directed behavioral weight control program. *Behav Med* 2011 Apr;37(2):47-53.

Lindenmayer JP, Khan A, Wance D, Maccabee N, Kaushik S, Kaushik S. Outcome evaluation of a structured educational wellness program in patients with severe mental illness. *J Clin Psychiatry* 2009 Oct;70(10):1385-96.

Lindstrom J, Peltonen M, Tuomilehto J. Lifestyle strategies for weight control: Experience from the Finnish Diabetes Prevention Study. *Proc Nutr Soc* 2005;64(1):81-8.

Lindstrom J, Ilanne-Parikka P, Peltonen M, Aunola S, Eriksson JG, Hemio K, et al. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet* 2006;368(9548):1673-9.

Linkov F, Maxwell GL, Felix AS, Lin Y, Lenzner D, Bovbjerg DH, et al. Longitudinal evaluation of cancer-associated biomarkers before and after weight loss in RENEW study participants: Implications for cancer risk reduction. *Gynecol Oncol* 2012;125(1):114-9.

Linmans JJ, Spigt MG, Deneer L, Lucas AEM, de Bakker M, Gidding LG, et al. Effect of lifestyle intervention for people with diabetes or prediabetes in real-world primary care: propensity score analysis. *BMC Fam Pract* 2011;12:95.

Lisspers J, Sundin O, Hofman-Bang C, Nordlander R, Nygren A, Ryden L, et al. Behavioral effects of a comprehensive, multifactorial program for lifestyle change after percutaneous transluminal coronary angioplasty: a prospective, randomized controlled study. *J Psychosom Res* 1999;46(2):143-54.



Lisspers J. Long-term effects of lifestyle behavior change in coronary artery disease: effects on recurrent coronary events after percutaneous coronary intervention. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association* 2005;24(1):41.

List JF, Woo V, Morales E, Tang W, Fiedorek FT. Sodium-glucose cotransport inhibition with dapagliflozin in type 2 diabetes. *Diabetes Care* 2009 Apr;32(4):650-7.

Liu X, Miller YD, Burton NW, Chang JH, Brown WJ. Qi-gong mind-body therapy and diabetes control. A randomized controlled trial. *Am J Prev Med* 2011 Aug;41(2):152-8.

Liutkus J, Rosas GJ, Norwood P, Pop L, Northrup J, Cao D, et al. A placebo-controlled trial of exenatide twice-daily added to thiazolidinediones alone or in combination with metformin. *Diabetes Obes Metab* 2010;12(12):1058-65.

Lloyd A, Khan R. Evaluation of Healthy Choices: a commercial weight loss programme commissioned by the NHS. *PERSPECT PUBLIC HEALTH* 2011 Jul;131(4):177-83.

Lloyd KB, Thrower MR, Walters NB, Krueger KP, Stamm PL, Evans RL. Implementation of a weight management pharmaceutical care service. *Ann Pharmacother* 2007 Feb;41(2):185-92.

Long DA, Perry TL. An application of survival analysis to population health management program evaluation. *Am J Health Promot* 2007 Jul;21(6):529-33.

Long SJ, Hart K, Morgan LM. The ability of habitual exercise to influence appetite and food intake in response to high- and low-energy preloads in man. *Br J Nutr* 2002;87(5):517-23.

Lovelady C. Balancing exercise and food intake with lactation to promote post-partum weight loss. *Proc Nutr Soc* 2011;70(2):181-4.

Lowndes J, Zoeller RF, Caplan JD, Kyriazis GA, Moyna NM, Seip RL, et al. Leptin responses to long-term cardiorespiratory exercise training without concomitant weight loss: a prospective study. *J SPORTS MED PHYS FITNESS* 2008 Sep;48(3):391-7.

Lund SS, Tarnow L, Stehouwer CD, Schalkwijk CG, Teerlink T, Gram J, et al. Impact of metformin versus repaglinide on non-glycaemic cardiovascular risk markers related to inflammation and endothelial dysfunction in non-obese patients with type 2 diabetes. *EUR J ENDOCRINOL* 2008 May;158(5):631-41.

Lundholm K, Gunnebo L, Korner U, Iresjo B-M, Engstrom C, Hyltander A, et al. Effects by daily long term provision of ghrelin to unselected weight-losing cancer patients: A randomized double-blind study. *Cancer* 2010;116(8):2044-52.

Lupton BS, Fonnebo V, Sogaard AJ, Langfeldt E. The Finnmark Intervention Study. Better health for the fishery population in an Arctic village in North Norway. *SCAND J PRIM HEALTH CARE* 2002;20(4):213-8.

Makoundou V, Habicht F, Bobbioni-Harsch E, Pataky Z, Golay A. [Long term weight loss maintenance]. [French]. *Rev Med Suisse* 2010 Mar 31;6(242):682-4.

Makrilakis K, Liatis S, Grammatikou S, Perrea D, Katsilambros N. Implementation and effectiveness of the first community lifestyle intervention programme to prevent Type 2 diabetes in Greece. the DE-PLAN study. *Diabet Med* 2010;27(4):459-65.

Manglik S, Cobanov B, Flores G, Nadjafi R, Tayek JA. Serum insulin but not leptin is associated with spontaneous and growth hormone (GH)-releasing hormone-stimulated GH secretion in

normal volunteers with and without weight loss. *Metabolism: Clinical & Experimental* 1998 Sep;47(9):1127-33.

Mann DM, Lin JJ. Increasing efficacy of primary care-based counseling for diabetes prevention: rationale and design of the ADAPT (Avoiding Diabetes Thru Action Plan Targeting) trial. *Implementation Science* 2012;7:6.

Manore MM. Dietary supplements for improving body composition and reducing body weight: Where is the evidence? *International Journal of Sport Nutrition and Exercise Metabolism* 2012;22(2):139-54.

Marston AR, Criss J. Further evidence for the maintenance of successful weight reduction. *Int J Eating Disord* 1986 May;5(4):765-9.

Martin CK, Heilbronn LK, de JL, DeLany JP, Volaufova J, Anton SD, et al. Effect of calorie restriction on resting metabolic rate and spontaneous physical activity. *Obesity* 2007 Dec;15(12):2964-73.

Martin CK, Anton SD, Han H, York-Crowe E, Redman LM, Ravussin E, et al. Examination of cognitive function during six months of calorie restriction: results of a randomized controlled trial. *Rejuvenation Research* 2007 Jun;10(2):179-90.

Martin CK, Talamini L, Johnson A, Hymel AM, Khavjou O. Weight loss and retention in a commercial weight-loss program and the effect of corporate partnership. *Int J Obes* 2010 Apr;34(4):742-50.

Martin CK, Das SK, Lindblad L, Racette SB, McCrory MA, Weiss EP, et al. Effect of calorie restriction on the free-living physical activity levels of nonobese humans: results of three randomized trials. *J Appl Physiol* 2011 Apr;110(4):956-63.

Martin CK, O'Neil PM, Pawlow L. Changes in Food Cravings during Low-Calorie and Very-Low-Calorie Diets. [References]. *Obes Res* 2006 Jan;14(1):115-21.

Maruthur NM, Wang NY, Appel LJ. Lifestyle interventions reduce coronary heart disease risk: results from the PREMIER Trial. *Circulation* 2009 Apr;119(15):2026-31.

Mathieu M-E, Brochu M, Beliveau L. DiabetAction: Changes in physical activity practice, fitness, and metabolic syndrome in type 2 diabetic and at-risk individuals. *Clin J Sport Med* 2008;18(1):70-5.

Matthews DR, Dejager S, Ahren B, Fonseca V, Ferrannini E, Couturier A, et al. Vildagliptin add-on to metformin produces similar efficacy and reduced hypoglycaemic risk compared with glimepiride, with no weight gain: results from a 2-year study. *Diabetes Obes Metab* 2010 Sep;12(9):780-9.

Mattila E, Parkka J, Hermersdorf M, Kaasinen J, Vainio J, Samposalo K, et al. Mobile diary for wellness management--results on usage and usability in two user studies. *IEEE Transactions on Information Technology in Biomedicine* 2008 Jul;12(4):501-12.

Mattila E, Lappalainen R, Parkka J, Salminen J, Korhonen I. Use of a mobile phone diary for observing weight management and related behaviours. *Journal of Telemedicine & Telecare* 2010 Jul;16(5):260-4.

Mayer-Davis EJ, D'antonio A, Martin M, Wandersman A, Parra-Medina D, Schulz R. Pilot study of strategies for effective weight management in type 2 diabetes: Pounds Off with Empowerment (POWER). *FAM COMMUNITY HEALTH* 2001 Jul;24(2):27-35.

Mcadam-Marx C, Bouchard J, Aagren M, Nelson R, Brixner D. Analysis of glycaemic control and weight change in patients initiated with human or analog insulin in an US ambulatory care setting. *Diabetes Obes Metab* 2010;12(1):54-64.

McBride PE, Einerson JA, Grant H, Sargent C, Underbakke G, Vitcenda M, et al. Putting the Diabetes Prevention Program into practice: a program for weight loss and cardiovascular risk reduction for patients with metabolic syndrome or type 2 diabetes mellitus. *Journal of Nutrition, Health & Aging* 2008 Dec;12(10):745S-9S.

McCall AL. Is there a magic diet? Studying the balance of macronutrients needed for best weight loss. *Curr Diab Rep* 2010;10(3):165-9.

McCauley KM. Modifying women's risk for cardiovascular disease. *J Obstet Gynecol Neonatal Nurs* 2007;36(2):116-24.

McGuffin LE, Wallace JMW, McCrorie TA, Price RK, Pourshahidi LK, Livingstone MBE. Family eating out-of-home: A review of nutrition and health policies. *Proc Nutr Soc* 2013;72(1):126-39.

McMahon A. Feasibility of a men's health promotion programme in Irish primary care. *Ir J Med Sci* 2002;171(1):20.

McTigue KM, Conroy MB, Hess R, Bryce CL, Fiorillo AB, Fischer GS, et al. Using the internet to translate an evidence-based lifestyle intervention into practice. *Telemed J E Health* 2009;15(9):851-8.

McTigue KM, Conroy MB, Bigi L, Murphy C, McNeil M. Weight loss through living well: translating an effective lifestyle intervention into clinical practice. *The Diabetes educator* 2009;35(2):199-208.

Mensink M, Feskens EJM, Saris WHM, de Bruin TWA, Blaak EE. Study on Lifestyle Intervention and Impaired Glucose Tolerance Maastricht (SLIM): Preliminary results after one year. [References]. *Int J Obes* 2003 Mar;27(3):377-84.

Merriam PA TTRMOBM. Methodology of a diabetes prevention translational research project utilizing a community-academic partnership for implementation in an underserved Latino community. *BMC Med Res Methodol* 2009;9:20.

Merrill RM, Massey MT, Aldana SG, Greenlaw RL, Diehl HA, Salberg A. C-reactive protein levels according to physical activity and body weight for participants in the coronary health improvement project. *PREV MED* 2008 May;46(5):425-30.

Miller CK, Gutschall MD, Holloman C. Self-monitoring predicts change in fiber intake and weight loss in adults with diabetes following an intervention regarding the glycemic index. [References]. *Patient Education and Counseling* 2009 Aug;76(2):213-9.

Milne RM, Mann JI, Chisholm AW, Williams SM. Long-term comparison of three dietary prescriptions in the treatment of NIDDM. *Diabetes Care* 1994 Jan;17(1):74-80.

Mitchel NS, Dickinson LM, Kempe A, Tsai AG. Determining the effectiveness of take off pounds sensibly (TOPS), a nationally available nonprofit weight loss program. [References]. *Obesity* 2011 Mar;19(3):568-73.

Mitjavila MT, Fandos M, Salas-Salvado J, Covas M-I, Borrego S, Estruch R, et al. The Mediterranean diet improves the systemic lipid and DNA oxidative damage in metabolic syndrome individuals. A randomized, controlled, trial. *Clin Nutr* 2013;32(2):172-8.

Mitsui T, Shimaoka K, Tsuzuku S, Kajioaka T, Sakakibara H. Gentle exercise of 40 minutes with dietary counseling is effective in treating metabolic syndrome. *Tohoku J Exp Med* 2008 Aug;215(4):355-61.

Monnikhof EM, Peeters PH, Schuit AJ. Design of the sex hormones and physical exercise (SHAPE) study. *BMC Public Health* 2007;7:232.

Moon KT. Long-term effects of weight loss diets on overall mood. *Am Fam Physician* 2010;81(8):1029-30.

Moreira MM, Souza HP, Schwingel PA, Sa CK, Zoppi CC. Effects of aerobic and anaerobic exercise on cardiac risk variables in overweight adults. *Arq Bras Cardiol* 2008;91(4):200-6.

Moreno-Navarrete JM, Manco M, Ibaez J, Garcia-Fuentes E, Ortega F, Gorostiaga E, et al. Metabolic endotoxemia and saturated fat contribute to circulating NGAL concentrations in subjects with insulin resistance. *Int J Obes* 2010;34(2):240-9.

Morgan LM, Griffin BA, Millward DJ, deLooy A, Fox KR, Baic S, et al. Comparison of the effects of four commercially available weight-loss programmes on lipid-based cardiovascular risk factors. *Public Health Nutr* 2009 Jun;12(6):799-807.

Morrison F, Shubina M, Turchin A. Lifestyle counseling in routine care and long-term glucose, blood pressure, and cholesterol control in patients with diabetes. *Diabetes Care* 2012;35(2):334-41.

Muckle S. An evaluation of a primary care-based weight management initiative. *Community Pract* 2007 Jul;80(7):20-3.

Mueller C, Masri B, Hogg J, Mastrogiacomo M, Chiu YL. Carbohydrate- vs fat-controlled diet effect on weight loss and coronary artery disease risk: a pilot feeding study. *NUTR CLIN PRACT* 2010 Oct;25(5):542-7.

Muller-Ehmsen J, Braun D, Schneider T, Pfister R, Worm N, Wielckens K, et al. Decreased number of circulating progenitor cells in obesity: beneficial effects of weight reduction. *Eur Heart J* 2008 Jun;29(12):1560-8.

Muller JE, Strater-Muller D, Marks H-J, Glasner M, Knepe P, Clemens-Harmening B, et al. Carbohydrate restricted diet in conjunction with metformin and liraglutide is an effective treatment in patients with deteriorated type 2 diabetes mellitus: Proof-of-concept study. *Nutr Metab* 2011;8 , 2011. Article Number: 92. Date of Publication: 2011.

Muniyappa R. The Diabetes Prevention Program. *Curr Diab Rep* 2003;3(3):221.

Nauck M, Marre M. Adding liraglutide to oral antidiabetic drug monotherapy: efficacy and weight benefits. *POSTGRAD MED* 2009 May;121(3):5-15.

Nauck MA, Ratner RE, Kapitza C, Berria R, Boldrin M, Balena R. Treatment with the human once-weekly glucagon-like peptide-1 analog taspoglutide in combination with metformin improves glycemic control and lowers body weight in patients with type 2 diabetes inadequately controlled with metformin alone: a double-blind placebo-controlled study. *Diabetes Care* 2009 Jul;32(7):1237-43.

Nauck MA, Vardarli I. Comparative evaluation of incretin-based antidiabetic medications and alternative therapies to be added to metformin in the case of monotherapy failure. *J Diabetes Invest* 2010;1(1-2):24-36.

Neumann JK. Self-management vs. therapist-management of weight reduction treatment. *Bulletin of the Society of Psychologists in Addictive Behaviors* 1985;4(1):41-8.

Neve MJ, Morgan PJ, Collins CE. Participant characteristics and reach of a commercial web-based weight loss program. *Nutrition & Dietetics* 2010 Dec;67(4):267-74.

Neve MJ, Collins CE, Morgan PJ. Dropout, nonusage attrition, and pretreatment predictors of nonusage attrition in a commercial Web-based weight loss program. *J Med Internet Res* 2010;12(4):e69.

Neve M, Morgan PJ, Collins CE. Weight change in a commercial web-based weight loss program and its association with website use: Cohort study. [References]. *J Med Internet Res* 2011 Oct;13(4):110-9.

Neyestani TR, Alipour-Birgani R, Siassi F, Rajayi M, Djalali M, Mohamadi M. Glycemic optimization may reduce lipid peroxidation independent of weight and blood lipid changes in Type 2 diabetes mellitus. *Diabetes, Nutrition & Metabolism - Clinical & Experimental* 2004 Oct;17(5):275-9.

Nicklas BJ, Gaukstern JE, Legault C, Leng I, Rejeski WJ. Intervening on spontaneous physical activity to prevent weight regain in older adults: Design of a randomized, clinical trial. *Contemp Clin Trials* 2012;33(2):450-5.

Nowalk MP, Wing RR. Changes in nutrient intake of hypertensives during a behavioral weight-control program. *Addict Behav* 1985;10(4):357-63.

Nyer PU, Dellande S. Public commitment as a motivator for weight loss. [References]. *Psychology & Marketing* 2010 Jan;27(1):1-12.

Obarzanek E, Vollmer WM, Lin PH, Cooper LS, Young DR, Ard JD, et al. Effects of individual components of multiple behavior changes: The PREMIER trial. [References]. *Am J Health Behav* 2007 Sep;31(5):545-60.

Olendzki BC, Ma Y, Schneider KL, Merriam P, Culver AL, Ockene IS, et al. A simple dietary message to improve dietary quality: Results from a pilot investigation. *Nutrition (Burbank, Los Angeles County, Calif)* 2009;25(7-8):736-44.

Orchard TJ, Temprosa M, Goldberg R, Haffner S, Ratner R, Marcovina S, et al. The effect of metformin and intensive lifestyle intervention on the metabolic syndrome: the Diabetes Prevention Program randomized trial. *Ann Intern Med* 2005;142(8):611-9.

Ouwehand C, Papias EK. Eat it or beat it. The differential effects of food temptations on overweight and normal-weight restrained eaters. *Appetite* 2010;55(1):56-60.

Owens DR, Luzio SD, Sert-Langeron C, Riddle MC. Effects of initiation and titration of a single preprandial dose of insulin glulisine while continuing titrated insulin glargine in type 2 diabetes: a 6-month 'proof-of-concept' study. *Diabetes Obes Metab* 2011 Nov;13(11):1020-7.

Paillard T, Lafont C, Costes-Salon MC, Dupui P. Comparison between three strength development methods on body composition in healthy elderly women. [References]. *Journal of Nutrition, Health & Aging* 2003;7(2):117-9.

Pallister C, Avery A, Stubbs J, Lavin J. Influence of Slimming World's lifestyle programme on diet, activity behaviour and health of participants and their families. *J HUM NUTR DIET* 2009 Aug;22(4):351-8.

Parker VG, Coles C, Logan BN, Davis L. The LIFE Project: A community-based weight loss intervention program for rural African American women. [References]. *Family & Community Health: The Journal of Health Promotion & Maintenance* 2010 Apr;33(2):133-43.

Parra-Medina D, Wilcox S, Wilson DK, Addy CL, Felton G, Poston MB. Heart Healthy and Ethnically Relevant (HHER) Lifestyle trial for improving diet and physical activity in underserved African American women. *Contemp Clin Trials* 2010;31(1):92-104.

Parra-Medina D. Results of the Heart Healthy and Ethnically Relevant Lifestyle trial: a cardiovascular risk reduction intervention for African American women attending community health centers. *Am J Public Health* 2011;101(10):1914.

Patel JV, Hughes EA, Lip GYH, Gill PS. Diabetes Health, Residence & Metabolism in Asians: The DHRMA study, research into foods from the Indian subcontinent - a blinded, randomised, placebo controlled trial. *BMC Cardiovasc Disord* 2011;11, 2011. Article Number: 70. Date of Publication: 02 Dec 2011.

Pendlebury J, Bushe CJ, Wildgust HJ, Holt RI. Long-term maintenance of weight loss in patients with severe mental illness through a behavioural treatment programme in the UK. *Acta Psychiatr Scand* 2007 Apr;115(4):286-94.

Peng W. Design and evaluation of a computer game to promote a healthy diet for young adults. *HEALTH COMMUN* 2009 Mar;24(2):115-27.

Perreault L, Kahn SE, Christophi CA, Knowler WC, Hamman RF, Diabetes Prevention Program Research Group. Regression from pre-diabetes to normal glucose regulation in the diabetes prevention program. *Diabetes Care* 2009 Sep;32(9):1583-8.

Peterson KE, Sorensen G, Pearson M, Hebert JR, Gottlieb BR, McCormick MC. Design of an intervention addressing multiple levels of influence on dietary and activity patterns of low-income, postpartum women. *HEALTH EDUC RES* 2002;17(5):531-40.

Petrella RJ AK. Efficacy of a family practice-based lifestyle intervention program to increase physical activity and reduce clinical and physiological markers of vascular health in patients with high normal blood pressure and/or high normal blood glucose (SNAC): study protocol for a randomized controlled trial. *Trials* 2011;12:45.

Petrofsky J, Batt J, Berk L, Schweichler C, Arai D, Conas C, et al. The effect of an exercise and diet program on fitness, posture and self image in women. *J APPL RES* 2010;10(1):40-51.

Phelan S, Roberts M, Lang W, Wing RR. Empirical evaluation of physical activity recommendations for weight control in women. *Med Sci Sports Exerc* 2007 Oct;39(10):1832-6.

Phelan S, Wyatt H, Nassery S, Dibello J, Fava JL, Hill JO, et al. Three-year weight change in successful weight losers who lost weight on a low-carbohydrate diet. [References]. *Obesity* 2007 Oct;15(10):2470-7.

Philippou E, McGowan BM, Brynes AE, Dornhorst A, Leeds AR, Frost GS. The effect of a 12-week low glycaemic index diet on heart disease risk factors and 24 h glycaemic response in healthy middle-aged volunteers at risk of heart disease: a pilot study. *Eur J Clin Nutr* 2008 Jan;62(1):145-9.

Piatt GA ARBMS. 3-year follow-up of clinical and behavioral improvements following a multifaceted diabetes care intervention: results of a randomized controlled trial. *The Diabetes educator* 2010;36(2):301.

Pieper C, Redman L, Racette S, Roberts S, Bhapkar M, Rochon J, et al. Development of adherence metrics for caloric restriction interventions. *Clinical Trials* 2011;8(2):155-64.

Pinto BM, Maruyama NC, Clark MM, Cruess DG, Park E, Roberts M. Motivation to modify lifestyle risk behaviors in women treated for breast cancer. *Mayo Clin Proc* 2002 Feb;77(2):122-9.

Pittaway JK, Ahuja KD, Cehun M, Chronopoulos A, Robertson IK, Nestel PJ, et al. Dietary supplementation with chickpeas for at least 5 weeks results in small but significant reductions in serum total and low-density lipoprotein cholesterol in adult women and men. *Ann Nutr Metab* 2006;50(6):512-8.

Plasqui G, Soenen S, Westerterp-Plantenga MS, Westerterp KR. Measurement of longitudinal changes in body composition during weight loss and maintenance in overweight and obese subjects

using air-displacement plethysmography in comparison with the deuterium dilution technique. *Int J Obes* 2011 Aug;35(8):1124-30.

Pollet PE, Leathwood PD, Richardson DP. Weight control using the Lean Plan diet: short and long term effects. *Int J Obes* 1989;13(Suppl 2):193-5.

Porterfield DS, Hinnant LW, Kane H, Horne J, McAleer K, Roussel A. Linkages between clinical practices and community organizations for prevention: A literature review and environmental scan. *Am J Prev Med* 2012;42(6 SUPPL. 2):S163-S171.

Poskiparta M, Kasila K, Kiuru P. Dietary and physical activity counselling on type 2 diabetes and impaired glucose tolerance by physicians and nurses in primary healthcare in Finland. *SCAND J PRIM HEALTH CARE* 2006 Dec;24(4):206-10.

Pratley R, Nauck M, Bailey T, Montanya E, Cuddihy R, Filetti S, et al. One year of liraglutide treatment offers sustained and more effective glycaemic control and weight reduction compared with sitagliptin, both in combination with metformin, in patients with type 2 diabetes: a randomised, parallel-group, open-label trial. *INT J CLIN PRACT* 2011 Apr;65(4):397-407.

Pratt DS, Jandzio M, Tomlinson D, Kang X, Smith E. The 5-10-25 challenge: an observational study of a web-based wellness intervention for a global workforce. *Disease Management* 2006 Oct;9(5):284-90.

Pritchard DA, Hyndman J, Taba F. Nutritional counselling in general practice: a cost effective analysis. *J EPIDEMIOLOG COMMUNITY HEALTH* 1999 May;53(5):311-6.

Pronk NP, Boucher JL, Gehling E, Boyle RG, Jeffery RW. A platform for population-based weight management: description of a health plan-based integrated systems approach. *Am J Managed Care* 2002 Oct;8(10):847-57.

Proper KI, van der Beek AJ, Hildebrandt VH, Twisk JW, van MW. Short term effect of feedback on fitness and health measurements on self reported appraisal of the stage of change. *Br J Sports Med* 2003 Dec;37(6):529-34.

Racette SB, Weiss EP, Schechtman KB, Steger-May K, Villareal DT, Obert KA, et al. Influence of weekend lifestyle patterns on body weight. [References]. *Obesity* 2008 Aug;16(8):1826-30.

Racette SB, Das SK, Bhapkar M, Hadley EC, Roberts SB, Ravussin E, et al. Approaches for quantifying energy intake and %calorie restriction during calorie restriction interventions in humans: the multicenter CALERIE study. *American Journal of Physiology - Endocrinology & Metabolism* 2012 Feb;302(4):E441-E448.

Ramezankhany A, Ali PN, Hedayati M. Comparing effects of aerobics, pilates exercises and low calorie diet on leptin levels and lipid profiles in sedentary women. *Iran J Basic Med Sci* 2011;14(3):256-63.

Rankins J, Wortham J, Brown LL. Modifying soul food for the Dietary Approaches to Stop Hypertension diet (DASH) plan: Implications for metabolic syndrome (DASH of Soul). *Ethn Dis* 2007;17(3):S4.

Rankovic G, Milicic B, Savic T, Dindic B, Mancev Z, Pesic G. Effects of physical exercise on inflammatory parameters and risk for repeated acute coronary syndrome in patients with ischemic heart disease. *Vojnosanitetski pregled Military-medical and pharmaceutical review* 2009 Jan;66(1):44-medical.

Ratliff J, Mutungi G, Puglisi MJ, Volek JS, Fernandez ML. Carbohydrate restriction (with or without additional dietary cholesterol provided by eggs) reduces insulin resistance and plasma leptin without modifying appetite hormones in adult men. *Nutr Res* 2009;29(4):262-8.

Ratner R, Wynne A, Nakhle S, Brusco O, Vlainic A, Rendell M. Influence of preprandial vs. postprandial insulin glulisine on weight and glycaemic control in patients initiating basal-bolus regimen for type 2 diabetes: a multicenter, randomized, parallel, open-label study (NCT00135096). *Diabetes Obes Metab* 2011 Dec;13(12):1142-8.

Rautio N, Jokelainen J, Oksa H, Saaristo T, Peltonen M, Puolijoki H, et al. Family history of diabetes and effectiveness of lifestyle counselling on the cardio-metabolic risk profile in individuals at high risk of Type 2 diabetes: 1-year follow-up of the FIN-D2D project. *Diabet Med* 2012;29(2):207-11.

Rautio N, Jokelainen J, Oksa H, Saaristo T, Peltonen M, Niskanen L, et al. Socioeconomic position and effectiveness of lifestyle intervention in prevention of type 2 diabetes: one-year follow-up of the FIN-D2D project. *SCAND J PUBLIC HEALTH* 2011;39(6):561-70.

Raz I, Hoekstra J, Fonseca V, Boldrin M, Kipnes M, Balena R, et al. Efficacy and safety of taspoglutide monotherapy in drug-naive type 2 diabetic patients after 24 weeks of treatment: Results of a randomized, double-blind, placebo-controlled phase 3 study (T-emerge 1). *Diabetes Care* 2012;35(3):485-7.

Razquin C, Martinez JA, Martinez-Gonzalez MA, Mitjavila MT, Estruch R, Marti A. A 3 years follow-up of a Mediterranean diet rich in virgin olive oil is associated with high plasma antioxidant capacity and reduced body weight gain. *Eur J Clin Nutr* 2009 Dec;63(12):1387-93.

Reames B, Burnett MF. Weight management practices of black paraprofessional women. *J AM DIET ASSOC* 1991 Jul;91(7):841-3.

Reasner C, Olansky L, Seck TL, Williams-Herman DE, Chen M, Terranella L, et al. The effect of initial therapy with the fixed-dose combination of sitagliptin and metformin compared with metformin monotherapy in patients with type 2 diabetes mellitus. *Diabetes Obes Metab* 2011 Jul;13(7):644-52.

Redenius R, Murphy C, O'Neill E, Al-Hamwi M, Zallek SN. Does CPAP lead to change in BMI? *J Clin Sleep Med* 2008 Jun 15;4(3):205-9.

Redman LM, Elkind-Hirsch K, Ravussin E. Aerobic exercise in women with polycystic ovary syndrome improves ovarian morphology independent of changes in body composition. *Fertil Steril* 2011;95(8):2696-9.

Rejeski WJ, Mihalko SL, Ambrosius WT, Bearon LB, McClelland JW. Weight loss and self-regulatory eating efficacy in older adults: the cooperative lifestyle intervention program. *Journals of Gerontology Series B-Psychological Sciences & Social Sciences* 2011 May;66(3):279-86.

Relton C, Strong M, Li J. The 'Pounds for Pounds' weight loss financial incentive scheme: an evaluation of a pilot in NHS Eastern and Coastal Kent. *J PUBLIC HEALTH* 2011 Dec;33(4):536-42.

Ricanati EH, Golubi M, Yang D, Saager L, Mascha EJ, Roizen MF. Mitigating preventable chronic disease: Progress report of the Cleveland Clinic's Lifestyle 180 program. *Nutr Metab* 2011;8, 2011. Article Number: 83. Date of Publication: 2011.

Richard C, Couillard C, Royer M-M, Desroches S, Couture P, Lamarche B. Impact of the Mediterranean diet with and without weight loss on plasma cell adhesion molecule concentrations in men with the metabolic syndrome. *Mediterr J Nutr Metab* 2011;4(1):33-9.



Richardson CR, Buis LR, Janney AW, Goodrich DE, Sen A, Hess ML, et al. An online community improves adherence in an internet-mediated walking program. Part 1: results of a randomized controlled trial. *J Med Internet Res* 2010;12(4):e71.

Richman RM, Loughnan GT, Droulers AM, Steinbeck KS, Caterson ID. Self-efficacy in relation to eating behaviour among obese and non-obese women. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 2001 Jun;25(6):907-13.

Rickman AD, Williamson DA, Martin CK, Gilhooly CH, Stein RI, Bales CW, et al. The CALERIE Study: design and methods of an innovative 25% caloric restriction intervention. [Review]. *Contemp Clin Trials* 2011 Nov;32(6):874-81.

Riddle M, Pencek R, Charenkavanich S, Lutz K, Wilhelm K, Porter L. Randomized comparison of pramlintide or mealtime insulin added to basal insulin treatment for patients with type 2 diabetes. *Diabetes Care* 2009 Sep;32(9):1577-82.

Riedt CS, Buckley BT, Brolin RE, Ambia-Sobhan H, Rhoads GG, Shapses SA. Blood lead levels and bone turnover with weight reduction in women. *Journal of Exposure Science & Environmental Epidemiology* 2009 Jan;19(1):90-6.

Rimmer JH, Braunschweig C, Silverman K, Riley B, Creviston T, Nicola T. Effects of a short-term health promotion intervention for a predominantly African-American group of stroke survivors. *Am J Prev Med* 2000;18(4):332-8.

Rittig K, Hieronimus A, Thamer C, Machann J, Peter A, Stock J, et al. Reducing visceral adipose tissue mass is essential for improving endothelial function in type 2 diabetes prone individuals. *Atherosclerosis* 2010;212(2):575-9.

Roach JB, Yadrick MK, Johnson JT, Boudreaux LJ, Forsythe WA, III, Billon W. Using self-efficacy to predict weight loss among young adults. *J AM DIET ASSOC* 2003 Oct;103(10):1357-9.

Rocha-Goldberg MP, Corsino L, Batch B, Voils CI, Thorpe CT, Bosworth HB, et al. Hypertension Improvement Project (HIP) Latino: results of a pilot study of lifestyle intervention for lowering blood pressure in Latino adults. *ETHNICITY HEALTH* 2010 Jun;15(3):269-82.

Rodriguez A, Reviriego J, Karamanos V, del Canizo FJ, Vlachogiannis N, Drossinos V. Management of cardiovascular risk factors with pioglitazone combination therapies in type 2 diabetes: An observational cohort study. *Cardiovasc diabetol* 2011;10, 2011. Article Number: 18. Date of Publication: 11 Feb 2011.

Rolls BJ, Roe LS, Meengs JS. Salad and satiety: energy density and portion size of a first-course salad affect energy intake at lunch. *J AM DIET ASSOC* 2004 Oct;104(10):1570-6.

Rose SB, Lawton BA, Elley CR, Dowell AC, Fenton AJ. The 'Women's Lifestyle Study', 2-year randomized controlled trial of physical activity counselling in primary health care: rationale and study design. *BMC Public Health* 2007;7:166.

Rosenberger PH, Ruser C, Kashaf S. MOVE! multidisciplinary programs: Challenges and resources for weight management treatment in VHA. *Transl Behav Med* 2011;1(4):629-34.

Rosenstock J, Davies M, Home PD, Larsen J, Koenen C, Scherthaner G. A randomised, 52-week, treat-to-target trial comparing insulin detemir with insulin glargine when administered as add-on to glucose-lowering drugs in insulin-naive people with type 2 diabetes. *Diabetologia* 2008 Mar;51(3):408-16.

Rosenstock J, Hollander P, Chevalier S, Iranmanesh A, SERENADE Study Group. SERENADE: the Study Evaluating Rimonabant Efficacy in Drug-naive Diabetic Patients: effects of monotherapy with rimonabant, the first selective CB1 receptor antagonist, on glycemic control, body weight, and lipid profile in drug-naive type 2 diabetes. *Diabetes Care* 2008 Nov;31(11):2169-76.

Rosenstock J, Cefalu WT, Hollander PA, Belanger A, Eliaschewitz FG, Gross JL, et al. Two-year pulmonary safety and efficacy of inhaled human insulin (Exubera) in adult patients with type 2 diabetes. *Diabetes Care* 2008 Sep;31(9):1723-8.

Rosenstock J, Rendell MS, Gross JL, Fleck PR, Wilson CA, Mekki Q. Alogliptin added to insulin therapy in patients with type 2 diabetes reduces HbA(1C) without causing weight gain or increased hypoglycaemia. *Diabetes Obes Metab* 2009 Dec;11(12):1145-52.

Rotenberg KJ, Carte L, Speirs A. The effects of modeling dietary restraint on food consumption: do restrained models promote restrained eating? *Eat behav* 2005 Jan;6(1):75-84.

Roumen C, Feskens EJ, Corpeleijn E, Mensink M, Saris WH, Blaak EE. Predictors of lifestyle intervention outcome and dropout: the SLIM study. *Eur J Clin Nutr* 2011 Oct;65(10):1141-7.

Rowley KG, Daniel M, Skinner K, Skinner M, White GA, O'Dea K. Effectiveness of a community-directed 'healthy lifestyle' program in a remote Australian aboriginal community. *Aust NZ J Public Health* 2000 Apr;24(2):136-44.

Rubio-Aliaga I, Marvin-Guy LF, Wang P, Wagniere S, Mansourian R, Fuerholz A, et al. Mechanisms of weight maintenance under high- and low-protein, low-glycaemic index diets. *Mol Nutr Food Res* 2011 Nov;55(11):1603-12.

Rush EC, Chandu V, Plank LD. Reduction of abdominal fat and chronic disease factors by lifestyle change in migrant Asian Indians older than 50 years. *Asia Pac J Clin Nutr* 2007;16(4):671-6.

Saaristo T, Moilanen L, Korpi-Hyovalti E, Vanhala M, Saltevo J, Niskanen L, et al. Lifestyle intervention for prevention of type 2 diabetes in primary health care: one-year follow-up of the Finnish National Diabetes Prevention Program (FIN-D2D). *Diabetes Care* 2010;33(10):2146-51.

Salas-Salvado J, Fernandez-Ballart J, Ros E, Martinez-Gonzalez M-A, Fito M, Estruch R, et al. Effect of a Mediterranean diet supplemented with nuts on metabolic syndrome status: One-year results of the PREDIMED randomized trial. *Arch Intern Med* 2008;168(22):2449-58.

Saltychev M, Laimi K, El-Metwally A, Oksanen T, Pentti J, Virtanen M, et al. Effectiveness of multidisciplinary early rehabilitation in reducing behaviour-related risk factors. *J Rehabil Med* 2012;44(4):370-7.

Sangster J, Furber S, Allman-Farinelli M, Haas M, Phongsavan P, Mark A, et al. A population-based lifestyle intervention to promote healthy weight and physical activity in people with cardiac disease: the PANACHE (Physical Activity, Nutrition And Cardiac HEalth) study protocol. *BMC Cardiovasc Disord* 2010;10:17.

Sarkadi A, Rosenqvist U. Field test of a group education program for type 2 diabetes: measures and predictors of success on individual and group levels. *PATIENT EDUC COUNS* 2001 Aug;44(2):129-39.

Sartore G, Reitano R, Barison A, Magnanini P, Cosma C, Burlina S, et al. The effects of psyllium on lipoproteins in type II diabetic patients. *Eur J Clin Nutr* 2009 Oct;63(10):1269-71.

Sartorelli DS SEFLCM. Beneficial effects of short-term nutritional counselling at the primary health-care level among Brazilian adults. *Public Health Nutr* 2005;8(7):820.

Scala D, D'Avino M, Cozzolino S, Mancini A, Andria B, Caruso G, et al. Promotion of behavioural change in people with hypertension: an intervention study. *Pharm World Sci* 2008;30(6):834-9.

Schafer S, Kantartzis K, Machann J, Venter C, Niess A, Schick F, et al. Lifestyle intervention in individuals with normal versus impaired glucose tolerance. *Eur J Clin Invest* 2007;37(7):535-43.

Schattner MA, Willis HJ, Raykher A, Brown P, Quesada O, Scott B, et al. Long-term enteral nutrition facilitates optimization of body weight. *Jpen: Journal of Parenteral & Enteral Nutrition* 2005 May;29(3):198-203.

Schautz B, Later W, Heller M, Muller MJ, Bosy-Westphal A. Associations between breast adipose tissue, body fat distribution and cardiometabolic risk in women: cross-sectional data and weight-loss intervention. *Eur J Clin Nutr* 2011 Jul;65(7):784-90.

Scherbaum WA, Schweizer A, Mari A, Nilsson PM, Lalanne G, Jauffret S, et al. Efficacy and tolerability of vildagliptin in drug-naïve patients with type 2 diabetes and mild hyperglycaemia\*. *Diabetes Obes Metab* 2008;10(8):675-82.

Scheuermann W, Ploch M, Morgenstern W, Nussel E. Changes in cholesterol and body weight at midpoint of the German Cardiovascular Prevention Study: Attempts at an interpretation. *Ann Epidemiol* 1993;3(5 SUPPL.):S28-S35.

Schoenthaler A, Luerassi L, Teresi JA, Silver S, Kong J, Odedosu T, et al. A practice-based trial of blood pressure control in African Americans (TLC-Clinic): study protocol for a randomized controlled trial. *Trials* 2011;12, 2011. Article Number: 265. Date of Publication: 22 Dec 2011.

Schweiger U, Laessle RG, Pirke KM. Macronutrient intake and mood during weight-reducing diets. *Annals of the New York Academy of Sciences* 1987;499:335-7.

Seck TL, Engel SS, Williams-Herman DE, Sisk CM, Golm GT, Wang H, et al. Sitagliptin more effectively achieves a composite endpoint for A1C reduction, lack of hypoglycemia and no body weight gain compared with glipizide. *DIABETES RES CLIN PRACT* 2011 Jul;93(1):e15-e17.

Shahar DR, Schwarzfuchs D, Fraser D, Vardi H, Thiery J, Fiedler GM, et al. Dairy calcium intake, serum vitamin D, and successful weight loss. *Am J Clin Nutr* 2010 Nov;92(5):1017-22.

Shea MK, Nicklas BJ, Houston DK, Miller ME, Davis CC, Kitzman DW, et al. The effect of intentional weight loss on all-cause mortality in older adults: results of a randomized controlled weight-loss trial. *Am J Clin Nutr* 2011 Sep;94(3):839-46.

Sherwood NE, Morton N, Jeffery RW, French SA, Neumark-Sztainer D, Falkner NH. Consumer preferences in format and type of community-based weight control programs. *Am J Health Promot* 1998 Sep;13(1):12-8.

Sherwood NE, Crain AL, Martinson BC, Hayes MG, Anderson JD, Clausen JM, et al. Keep it off: a phone-based intervention for long-term weight-loss maintenance. *Contemp Clin Trials* 2011 Jul;32(4):551-60.

Shintani T, Beckham S, O'Connor HK, Hughes C, Sato A. The Waiānae Diet Program: a culturally sensitive, community-based obesity and clinical intervention program for the Native Hawaiian population. *Hawaii Med J* 1994;53(5):136-41.

Shintani TT, Beckham S, Brown AC, O'Connor HK. The Hawaii Diet: ad libitum high carbohydrate, low fat multi-cultural diet for the reduction of chronic disease risk factors: obesity, hypertension, hypercholesterolemia, and hyperglycemia. *Hawaii Med J* 2001 Mar;60(3):69-73.

Shrestha P, Ghimire L. A review about the effect of life style modification on diabetes and quality of life. *Glob J Health Sci* 2012;4(6):185-90.

Simmons D, Rush E, Crook N, Te W. Development and piloting of a community health worker-based intervention for the prevention of diabetes among New Zealand Maori in Te Wai o Rona: Diabetes Prevention Strategy. *Public Health Nutr* 2008 Dec;11(12):1318-25.

Slinde F, Gronberg AM, Engstrom CR, Rossander-Hulthen L, Larsson S. Individual dietary intervention in patients with COPD during multidisciplinary rehabilitation. *Respir Med* 2002 May;96(5):330-6.

Smith M, Politzer N, Macgarvie D, McAndrews MP, Del CM. Efficacy and tolerability of the modified Atkins diet in adults with pharmaco-resistant epilepsy: a prospective observational study. *Epilepsia* 2011 Apr;52(4):775-80.

Smith PJ, Blumenthal JA, Babyak MA, Georgiades A, Hinderliter A, Sherwood A. Effects of exercise and weight loss on depressive symptoms among men and women with hypertension. *J Psychosom Res* 2007 Nov;63(5):463-9.

Smolander J, Blair SN, Kohl HW. Work ability, physical activity, and cardiorespiratory fitness: 2-year results from Project Active. *J Occup Environ Med* 2000;42(9):906-10.

Sola E, Vaya A, Villa P, Espana F, Estelles A, Garcia K, et al. Obesity and activated protein C resistance. *Pathophysiology of Haemostasis & Thrombosis* 2008;36(2):64-8.

Sonntag U, Henkel J, Renneberg B, Bockelbrink A, Braun V, Heintze C. Counseling overweight patients: analysis of preventive encounters in primary care. *International Journal for Quality in Health Care* 2010 Dec;22(6):486-92.

Sopko G, Jacobs J, Jeffery R. Effects on blood lipids and body weight in high risk men of a practical exercise program. *Atherosclerosis* 1983;49(3):219-29.

Sorensen LB, Soe M, Halkier KH, Stigsby B, Astrup A. Effects of increased dietary protein-to-carbohydrate ratios in women with polycystic ovary syndrome. *Am J Clin Nutr* 2012 Jan;95(1):39-48.

Sowers JR, Whitfield LA, Beck FWJ. Role of enhanced sympathetic nervous system activity and reduced Na<sup>+</sup>,K<sup>+</sup>-dependent adenosine triphosphatase activity in maintenance of elevated blood pressure in obesity: Effects of weight loss. *Clin Sci* 1982;63(Suppl. 8):121S-45.

Staffileno BA, Minnick A, Coke LA, Hollenberg SM. Blood pressure responses to lifestyle physical activity among young, hypertension-prone African-American women. [References]. *J CARDIOVASC NURS* 2007 Mar;22(2):107-17.

Stamler R, Stamler J, Gosch FC, McDonald AM. Primary prevention of hypertension--a randomized controlled trial. *Annals of Clinical Research* 1984;16:Suppl-42.

Steffen PR, Sherwood A, Gullette EC, Georgiades A, Hinderliter A, Blumenthal JA. Effects of exercise and weight loss on blood pressure during daily life. *Med Sci Sports Exerc* 2001 Oct;33(10):1635-40.

Steinhardt MA, Bezner JR, Adams TB. Outcomes of a traditional weight control program and a nondiet alternative: A one-year comparison. *Journal of Psychology: Interdisciplinary and Applied* 1999 Sep;133(5):495-513.

Stensvold D, Tjonna AE, Skaug E-A, Aspenes S, Stolen T, Wisloff U, et al. Strength training versus aerobic interval training to modify risk factors of metabolic syndrome. *J Appl Physiol* 2010;108(4):804-10.

Step toe A, Kerry S, Rink E, Hilton S. The impact of behavioral counseling on stage of change in fat intake, physical activity, and cigarette smoking in adults at increased risk of coronary heart disease. *Am J Public Health* 2001 Feb;91(2):265-9.

Stewart LK, Earnest CP, Blair SN, Church TS. Effects of different doses of physical activity on C-reactive protein among women. *Med Sci Sports Exerc* 2010 Apr;42(4):701-7.

Stice E, Chase A, Stormer S, Appel A. A randomized trial of a dissonance-based eating disorder prevention program. *Int J Eating Disord* 2001 Apr;29(3):247-62.

Stice E, Ragan J. A preliminary controlled evaluation of an eating disturbance psychoeducational intervention for college students. *Int J Eating Disord* 2002 Mar;31(2):159-71.

Stice E, Sysko R, Roberto CA, Allison S. Are dietary restraint scales valid measures of dietary restriction? Additional objective behavioral and biological data suggest not. *Appetite* 2010;54(2):331-9.

Stoddard AM, Palombo R, Troped PJ, Sorensen G, Will JC. Cardiovascular disease risk reduction: the Massachusetts WISEWOMAN project. *J Women's Health* 2004 Jun;13(5):539-46.

Stoernell CK, Tangney CC, Rockway SW. Short-term changes in lipoprotein subclasses and C-reactive protein levels of hypertriglyceridemic adults on low-carbohydrate and low-fat diets. *Nutrition research (New York, NY)* 2008;28(7):443-9.

Stolley MR, Fitzgibbon ML, Wells A, Martinovich Z. Addressing multiple breast cancer risk factors in African-American women. *J Natl Med Assoc* 2004 Jan;96(1):76-86.

Stotland S, Zuroff DC. Relations between multiple measures of dieting self-efficacy and weight change in a behavioral weight control program. *BEHAV THER* 1991;22(1):47-59.

Stubbs RJ, Pallister C, Whybrow S, Avery A, Lavin J. Weight outcomes audit for 34,271 adults referred to a primary care/commercial weight management partnership scheme. *Obes Facts* 2011;4(2):113-20.

Sturt J, Ali S, Robertson W, Metcalfe D, Grove A, Bourne C, et al. Neurolinguistic programming: A systematic review of the effects on health outcomes. *Br J Gen Pract* 2012;62(604):e757-e764.

Sun AS, Ostadal O, Ryznar V, Dulik I, Dusek J, Vaclavik A, et al. Phase I/II study of stage III and IV non-small cell lung cancer patients taking a specific dietary supplement. *NUTR CANCER* 1999;34(1):62-9.

Svetkey LP, Pollak KI, Yancy WS, Dolor RJ, Batch BC, Samsa G, et al. Hypertension improvement project: randomized trial of quality improvement for physicians and lifestyle modification for patients. *Hypertension* 2009 Dec;54(6):1226-33.

Svetkey LP, Ard JD, Stevens VJ, Loria CM, Young DY, Hollis JF, et al. Predictors of long-term weight loss in adults with modest initial weight loss, by sex and race. *Obesity (Silver Spring)* 2012;20(9):1820-8.

Tangney DJ, Brown DB, Brown GJ, Prescott GJ. Cardiovascular risk reduction in men: a nine-year cohort study. *Br J Gen Pract* 2002 Sep;52(482):743-5.

Tapper K, Shaw C, Ilesley J, Hill AJ, Bond FW, Moore L. Exploratory randomised controlled trial of a mindfulness-based weight loss intervention for women. *Appetite* 2009 Apr;52(2):396-404.

Tarkun I, Dikmen E, Cetinarslan B, Canturk Z. Impact of treatment with metformin on adipokines in patients with polycystic ovary syndrome. *European Cytokine Network* 2010 Dec 1;21(4):272-7.

Tate DF. A series of studies examining Internet treatment of obesity to inform Internet interventions for substance use and misuse. *Substance Use & Misuse* 2011;46(1):57-65.

Taylor J, Cottrell C, Chatterton H, Hill J, Hughes R, Wohlgemuth C, et al. Identifying risk and preventing progression to Type 2 diabetes in vulnerable and disadvantaged adults: a pragmatic review. *Diabet Med* 2013;30(1):16-25.

Teas J, Baldeon ME, Chiriboga DE, Davis JR, Sarries AJ, Braverman LE. Could dietary seaweed reverse the metabolic syndrome? *Asia Pac J Clin Nutr* 2009;18(2):145-54.

Teixeira AM, Martins RA, Neves AP, Coelho-Silva MJ, Verissimo MT. The effect of aerobic versus strength-based training on high-sensitivity C-reactive protein in older adults. *Eur J Appl Physiol* 2010;110(1):161-9.

Tetlie T, Eik-Nes N, Palmstierna T, Callaghan P, Nottestad JA. The effect of exercise on psychological & physical health outcomes: preliminary results from a Norwegian forensic hospital. *Journal of Psychosocial Nursing & Mental Health Services* 2008 Jul;46(7):38-43.

Thomas D, Vydellingum V, Lawrence J. E-mail contact as an effective strategy in the maintenance of weight loss in adults. *J HUM NUTR DIET* 2011 Feb;24(1):32-8.

Thomas GR, Kerr MP. Longitudinal follow-up of weight change in the context of a community-based health promotion programme for adults with an intellectual disability. [References]. *Journal of Applied Research in Intellectual Disabilities* 2011 Jul;24(4):381-7.

Thoolen BJ, de RD, Bensing J, Gorter K, Rutten G. Beyond good intentions: The role of proactive coping in achieving sustained behavioural change in the context of diabetes management. *PSYCHOL HEALTH* 2009 Mar;24(3):237-54.

Togami T. Interventions in local communities and work sites through Physical Activity and Nutrition Programme. *Obes Rev* 2008 Mar;9:Suppl-9.

Toll BA, Leary V, Wu R, Salovey P, Meandzija B, O'Malley SS. A preliminary investigation of naltrexone augmentation of bupropion to stop smoking with less weight gain. [References]. *Addict Behav* 2008 Jan;33(1):173-9.

Topol EJ, Bousser MG, Fox KA, Creager MA, Despres JP, Easton JD, et al. Rimonabant for prevention of cardiovascular events (CRESCENDO): a randomised, multicentre, placebo-controlled trial. *Lancet* 2010 Aug 14;376(9740):517-23.

Trevisan MC, Souza JM, Marucci MF. Influence of soy protein intake and weight training on the resting energy expenditure of postmenopausal women. *Rev Assoc Med Bras* 2010 Sep;56(5):572-8.

Trueman P, Haynes SM, Felicity LG, Louise ME, McQuigg MS, Mongia S, et al. Long-term cost-effectiveness of weight management in primary care. *INT J CLIN PRACT* 2010 May;64(6):775-83.

Tsilchorozidou T, Mohamed-Ali V, Conway GS. Determinants of interleukin-6 and C-reactive protein vary in polycystic ovary syndrome, as do effects of short- and long-term metformin therapy. *Hormone Research* 2009;71(3):148-54.

Tucker LA, Mortell R. Comparison of the effects of walking and weight training programs on body image in middle-aged women: an experimental study. *Am J Health Promot* 1993 Sep;8(1):34-42.

Turk MW, Sereika SM, Yang K, Ewing LJ, Hravnak M, Burke LE. Psychosocial correlates of weight maintenance among black & white adults. *Am J Health Behav* 2012 May;36(3):395-407.

Turner-McGrievy GM, Jenkins DJA, Barnard ND, Cohen J, Gloede L, Green AA. Decreases in dietary glycemic index are related to weight loss among individuals following therapeutic diets for type 2 diabetes. *J Nutr* 2011;141(8):1469-74.

Turner SL, Thomas AM, Wagner PJ, Moseley GC. A collaborative approach to wellness: diet, exercise, and education to impact behavior change. *J AM ACAD NURSE PRACT* 2008 Jun;20(6):339-44.

Tussing-Humphreys LM, Fitzgibbon ML, Kong A, Odoms-Young A. Weight loss maintenance in African American women: A systematic review of the behavioral lifestyle intervention literature. *J Obes* 2013;2013.

Twiner EM, Liu Z, Gimble J, Yu Y, Greenway F. Pharmacokinetic pilot study of the antiangiogenic activity of standardized platycodi radix. *Adv Ther* 2011 Oct;28(10):857-65.

Unalacak M, Kara IH, Baltaci D, Erdem O, Bucaktepe PGE. Effects of ramadan fasting on biochemical and hematological parameters and cytokines in healthy and obese individuals. *Metab syndr relat disord* 2011;9(2):157-61.

Uusitupa M. The Finnish Diabetes Prevention Study. *The British journal of nutrition* 2000;83 Suppl 1:S137.

Uusitupa MIJ. Early lifestyle intervention in patients with non-insulin-dependent diabetes mellitus and impaired glucose tolerance. *Ann Med* 1996;28(5):445-9.

Uusitupa M, Lindi V, Louheranta A, Salopuro T, Lindstrom J, Tuomilehto J, et al. Long-term improvement in insulin sensitivity by changing lifestyles of people with impaired glucose tolerance: 4-year results from the Finnish Diabetes Prevention Study. *Diabetes* 2003;52(10):2532-8.

Vadheim LM, Brewer KA, Kassner DR, Vanderwood KK, Hall TO, Butcher MK, et al. Effectiveness of a lifestyle intervention program among persons at high risk for cardiovascular disease and diabetes in a rural community. *J RURAL HEALTH* 2010;26(3):266-72.

van Sluijs EM, van Poppel MN, Twisk JW, Chin APM, Calfas KJ, van MW. Effect of a tailored physical activity intervention delivered in general practice settings: results of a randomized controlled trial. *Am J Public Health* 2005 Oct;95(10):1825-31.

van Strien T, van de Laar FA, van Leeuwe JFJ, Lucassen PLBJ, van den Hoogen HJM, Rutten GEHM, et al. The dieting dilemma in patients with newly diagnosed type 2 diabetes: Does dietary restraint predict weight gain 4 years after diagnosis? [References]. *Health Psychol* 2007 Jan;26(1):105-12.

van DS, Czernichow S, Chalmers J, Kengne AP, de Galan BE, Poulter N, et al. Weight changes and their predictors amongst 11 140 patients with type 2 diabetes in the ADVANCE trial. *Diabetes Obes Metab* 2012;14(5):464-9.

Vandelanotte C, De B, I, Sallis JF, Spittaels H, Brug J. Efficacy of sequential or simultaneous interactive computer-tailored interventions for increasing physical activity and decreasing fat intake. *Ann Behav Med* 2005 Apr;29(2):138-46.

Vardhan A, Adhikari Prabha MR, Kotian SM, Saxena N, Gupta S, Tripathy A. The value of the Indian diabetes risk score as a tool for reducing the risk of diabetes among Indian medical students. *J Clin Diagn Res* 2011;5(4):718-20.

Veerman JL, Barendregt JJ, Mackenbach JP, Brug J. Using epidemiological models to estimate the health effects of diet behaviour change: the example of tailored fruit and vegetable promotion. *Public Health Nutr* 2006 Jun;9(4):415-20.

Venables MC, Hulston CJ, Cox HR, Jeukendrup AE. Green tea extract ingestion, fat oxidation, and glucose tolerance in healthy humans. *Am J Clin Nutr* 2008 Mar;87(3):778-84.

Venditti EM, Bray GA, Carrion-Petersen ML, Delahanty LM, Edelstein SL, Hamman RF, et al. First versus repeat treatment with a lifestyle intervention program: Attendance and weight loss outcomes. [References]. *Int J Obes* 2008 Oct;32(10):1537-44.

Verheijden MW, Jans MP, Hildebrandt VH, Hopman-Rock M. Rates and determinants of repeated participation in a web-based behavior change program for healthy body weight and healthy lifestyle. *J Med Internet Res* 2007;9(1):e1-7.

Verweij LM, Proper KI, Leffelaar ER, Weel ANH, Nauta AP, Hulshof CTJ, et al. Barriers and facilitators to implementation of an occupational health guideline aimed at preventing weight gain among employees in the Netherlands. *J Occup Environ Med* 2012;54(8):954-60.

Vestfold Heartcare Study Group. Influence on lifestyle measures and five-year coronary risk by a comprehensive lifestyle intervention programme in patients with coronary heart disease. *Eur J Cardiovasc Prev Rehabil* 2003;10(6):429.

Villablanca AC, Arline S, Lewis J, Raju S, Sanders S, Carrow S. Outcomes of national community organization cardiovascular prevention programs for high-risk women. *J Cardiovasc Transl Res* 2009 Sep;2(3):306-20.

Villalon KL, Gozansky WS, Van Pelt RE, Wolfe P, Jankowski CM, Schwartz RS, et al. A losing battle: Weight regain does not restore weight loss-induced bone loss in postmenopausal women - Duplicate of Ref ID 9365. *Obesity* 2011;19(12):2345-50.

Vincent D. Culturally tailored education to promote lifestyle change in Mexican Americans with type 2 diabetes. *J AM ACAD NURSE PRACT* 2009 Sep;21(9):520-7.

Vissers D, Verrijken A, Mertens I, Van GC, Van de Sompel A, Truijen S, et al. Effect of long-term whole body vibration training on visceral adipose tissue: a preliminary report. *Obes Facts* 2010;3(2):93-100.

Volek JS, Phinney SD, Forsythe CE, Quann EE, Wood RJ, Puglisi MJ, et al. Carbohydrate restriction has a more favorable impact on the metabolic syndrome than a low fat diet. *Lipids* 2009 Apr;44(4):297-309.

von Huth SL. A randomized multifactorial intervention study for prevention of ischaemic heart disease (Inter99): the long-term effect on physical activity. *SCAND J PUBLIC HEALTH* 2008;36(4):380.

Wadden TA, West DS, Neiberg RH, Wing RR, Ryan DH, Johnson KC, et al. One-year weight losses in the Look AHEAD study: factors associated with success. *Obesity (Silver Spring, Md)* 2009;17(4):713-22.

Walters KL LJ. Project h[Latin small letter schwa]li?dx(w)/Healthy Hearts Across Generations: development and evaluation design of a tribally based cardiovascular disease prevention intervention for American Indian families. *J Prim Prev* 2012;33(4):197.

Wang P, Holst C, Andersen MR, Astrup A, Bouwman FG, van OS, et al. Blood profile of proteins and steroid hormones predicts weight change after weight loss with interactions of dietary protein level and glycemic index.[Erratum appears in *PLoS One*. 2011;6(7). doi:10.1371/annotation/2526dde5-cfaa-4f00-b554-25768bda2935]. *PLoS ONE [Electronic Resource]* 2011;6(2):e16773.

Wang P, Holst C, Astrup A, Bouwman FG, van OS, Wodzig WK, et al. Blood profiling of proteins and steroids during weight maintenance with manipulation of dietary protein level and glycaemic index. *Br J Nutr* 2012 Jan;107(1):106-19.



Wassertheil-Smoller S, Langford HG, Blaufox MD. Effective dietary intervention in hypertensives: Sodium restriction and weight reduction. *J AM DIET ASSOC* 1985;85(4):423-30.

Waters LA RMFBEE. Characteristics of control group participants who increased their physical activity in a cluster-randomized lifestyle intervention trial. *BMC Public Health* 2011;11:27.

Waters SB, Topp BG, Siler SQ, Alexander CM. Treatment with sitagliptin or metformin does not increase body weight despite predicted reductions in urinary glucose excretion. *Journal of Diabetes Science & Technology* 2009 Jan;3(1):68-82.

Webber KH, Tate DF, Quintiliani LM. Motivational interviewing in internet groups: a pilot study for weight loss. *J AM DIET ASSOC* 2008 Jun;108(6):1029-32.

Weekes CE, Emery PW, Elia M. Dietary counselling and food fortification in stable COPD: a randomised trial. *Thorax* 2009 Apr;64(4):326-31.

Weinberg RS. Effects of preexisting and manipulated self-efficacy on weight loss in a self-control program. *Journal of Research in Personality* 1984 Sep;18(3):352-8.

Weinstock RS, Brooks G, Palmas W, Morin PC, Teresi JA, Eimicke JP, et al. Lessened decline in physical activity and impairment of older adults with diabetes with telemedicine and pedometer use: results from the IDEATel study. *Age & Ageing* 2011 Jan;40(1):98-105.

Werkman A, Schuit AJ, Kwak L, Kremers SP, Visscher TL, Kok FJ, et al. Study protocol of a cluster randomised controlled trial investigating the effectiveness of a tailored energy balance programme for recent retirees. *BMC Public Health* 2006;6:293.

White DB, Bursac Z, DiLillo V, West DS. Weight loss goals among African-American women with type 2 diabetes in a behavioral weight control program. *Obesity* 2011 Nov;19(11):2283-5.

Wierenga ME. Life-style modification for weight control to improve diabetes health status. *Patient Education and Counseling* 1994 Apr;23(1):33-40.

Wikstrom K, Peltonen M, Eriksson JG, Aunola S, Ilanne-Parikka P, Keinanen-Kiukaanniemi S, et al. Educational attainment and effectiveness of lifestyle intervention in the Finnish Diabetes Prevention Study. *Diabetes Res Clin Pract* 2009;86(1):e1-e5.

Will CA, Ehrlinger J, Alquist JL, Conlon KE, Baumeister RF, Schatschneider C, et al. High trait self-control predicts positive health behaviors and success in weight loss. *J HEALTH PSYCHOL* 2011 Jul;16(5):750-9.

Williams K, Prevost AT, Griffin S, Hardeman W, Hollingworth W, Spiegelhalter D, et al. The ProActive trial protocol - a randomised controlled trial of the efficacy of a family-based, domiciliary intervention programme to increase physical activity among individuals at high risk of diabetes [ISRCTN61323766]. *BMC Public Health* 2004 Oct 18;4:48.

Williamson DA, Anton SD, Han H, Champagne CM, Allen R, Leblanc E, et al. Early behavioral adherence predicts short and long-term weight loss in the POUNDS LOST study. *J Behav Med* 2010 Aug;33(4):305-14.

Willis FB, Smith FM, Willis AP. Frequency of exercise for body fat loss: a controlled, cohort study. *J Strength Cond Res* 2009 Nov;23(8):2377-80.

Wilson K, Clark H, Hotz S, Dent R. Impact of smoking status on weight loss and cardiovascular risk factors. *J EPIDEMIOL COMMUNITY HEALTH* 2001 Mar;55(3):213-4.

Wilson MG, Goetzel RZ, Ozminkowski RJ, DeJoy DM, Della L, Roemer EC, et al. Using formative research to develop environmental and ecological interventions to address overweight and obesity. *Obesity* 2007 Nov;15:Suppl-47S.

Wing RR, Epstein LH. A community approach to weight control: the American Cancer Society Weight-A-Thon. *PREV MED* 1982 Mar;11(2):245-50.

Wing RR, Tate DF, Gorin AA, Raynor HA, Fava JL, Machan J. STOP regain: are there negative effects of daily weighing? [Erratum appears in *J Consult Clin Psychol*. 2007 Oct;75(5):715]. *Journal of Consulting & Clinical Psychology* 2007 Aug;75(4):652-6.

Wing RR, Crane MM, Thomas JG, Kumar R, Weinberg B. Improving weight loss outcomes of community interventions by incorporating behavioral strategies. *Am J Public Health* 2010 Dec;100(12):2513-9.

Wing RR, Tate DF, Gorin AA, Raynor HA, Fava JL, Machan J. "STOP regain": Are there negative effects of daily weighing? [References]. *J Consult Clin Psychol* 2007 Aug;75(4):652-6.

Winnick JJ, Gaillard T, Schuster DP. Resistance training differentially affects weight loss and glucose metabolism of White and African American patients with type 2 diabetes mellitus. *Ethn Dis* 2008;18(2):152-6.

Wise TN, Mann LS, Cooper JN, Rustigi EB. Patient characteristics and treatment results in a protein-modified fasting diet. *Int J Eating Disord* 1986 Sep;5(6):1089-94.

Wollner S, Blackburn D, Spellman K, Khaodhiar L, Blackburn GL. Weight-loss programs in convenient care clinics: A prospective cohort study. [References]. *Am J Health Promot* 2010 Sep;25(1):26-9.

Woollard J. Effects of a general practice-based intervention on diet, body mass index and blood lipids in patients at cardiovascular risk. *J CARDIOVASC RISK* 2003;10(1):31.

Wu J-D, Xu X-H, Zhu J, Ding B, Du T-X, Gao G, et al. Effect of exenatide on inflammatory and oxidative stress markers in patients with type 2 diabetes mellitus. *Diabetes Technol Ther* 2011;13(2):143-8.

Wu J, Zhu Y, Jiang Y, Cao Y. Effects of metformin and ethinyl estradiol-cyproterone acetate on clinical, endocrine and metabolic factors in women with polycystic ovary syndrome. *Gynecol Endocrinol* 2008 Jul;24(7):392-8.

Wyld B, Harrison A, Noakes M. The CSIRO Total Wellbeing Diet Book 1: sociodemographic differences and impact on weight loss and well-being in Australia. *Public Health Nutr* 2010 Dec;13(12):2105-10.

Wysham C, Bergenstal R, Malloy J, Yan P, Walsh B, Malone J, et al. DURATION-2: Efficacy and safety of switching from maximum daily sitagliptin or pioglitazone to once-weekly exenatide. *Diabet Med* 2011;28(6):705-14.

Yancey A, McCarthy WJ, Leslie J. Recruiting African-American women to community-based health promotion research. *Am J Health Promot* 1998 May;12(5):335-8.

Yancey AK, Jordan A, Bradford J, Voas J, Eller TJ, Buzzard M, et al. Engaging high-risk populations in community-level fitness promotion: ROCK! Richmond. *HEALTH PROMOT PRACT* 2003 Apr;4(2):180-8.

Yankura DJ, Conroy MB, Hess R, Pettee KK, Kuller LH, Kriska AM. Weight regain and health-related quality of life in postmenopausal women. [References]. *Obesity* 2008 Oct;16(10):2259-65.

Yatsuya H, Jeffery RW, Erickson DJ, Welsh EM, Flood AP, Jaeb MA, et al. Sex-specific HDL cholesterol changes with weight loss and their association with anthropometric variables: The LIFE study. *Obesity* 2011;19(2):429-35.

Yeh HC, Clark JM, Emmons KE, Moore RH, Bennett GG, Warner ET, et al. Independent but coordinated trials: insights from the practice-based Opportunities for Weight Reduction Trials Collaborative Research Group. *Clinical Trials* 2010 Aug;7(4):322-32.

Yoshinaga Y, Takahashi H, Yamamura Y, Hoshi K, Saito N, Someya K. Obesity and life style of middle aged Japanese women: an experience of a health promotion seminar at a public health center. *DIABETES RES CLIN PRACT* 1990;10:Suppl-82.

Young DR, Vollmer WM, King AC, Brown AJ, Stevens VJ, Elmer PJ, et al. Can individuals meet multiple physical activity and dietary behavior goals? *Am J Health Behav* 2009 May;33(3):277-86.

Young DR, Coughlin J, Jerome GJ, Myers V, Chae SE, Brantley PJ. Effects of the PREMIER interventions on health-related quality of life. *Ann Behav Med* 2010 Dec;40(3):302-12.

Zandstra EH, den HW, van der Meer N, van der Maas A. Improving compliance to meal-replacement food regimens. Forming implementation intentions (conscious IF-THEN plans) increases compliance. *Appetite* 2010 Dec;55(3):666-70.

## No intervention of interest

Eight-year follow-up results from the Rome Project of Coronary Heart Disease Prevention. Research Group of the Rome Project of Coronary Heart Disease Prevention. *PREV MED* 1986;15(2):176.

Albaugh VL, Singareddy R, Mauger D, Lynch CJ. A double blind, placebo-controlled, randomized crossover study of the acute metabolic effects of olanzapine in healthy volunteers. *PLoS ONE [Electronic Resource]* 2011;6(8):e22662.

Bajerska J, Jeszka J, Tarnowska AK, Czapka-Matyasik M. The effect of green and Oolong tea extracts supplementation on body composition in wrestlers. *Pak J Nutr* 2010;9(7):696-702.

Bartley PC, Bogoev M, Larsen J, Philotheou A. Long-term efficacy and safety of insulin detemir compared to Neutral Protamine Hagedorn insulin in patients with Type 1 diabetes using a treat-to-target basal-bolus regimen with insulin aspart at meals: a 2-year, randomized, controlled trial. *Diabet Med* 2008 Apr;25(4):442-9.

Beck AM, Damkjaer K, Sorbye LW. Physical and social functional abilities seem to be maintained by a multifaceted randomized controlled nutritional intervention among old (>65 years) Danish nursing home residents. *Arch Gerontol Geriatr* 2010;50(3):351-5.

Bell ZW, Canale RE, Bloomer RJ. A dual investigation of the effect of dietary supplementation with licorice flavonoid oil on anthropometric and biochemical markers of health and adiposity. *Lipids in Health & Disease* 2011;10:29.

Bigal ME, Lipton RB, Biondi DM, Xiang J, Hulihan J. Weight change and clinical markers of cardiovascular disease risk during preventive treatment of migraine. *Cephalalgia* 2009 Nov;29(11):1188-96.

Bize R, Willi C, Chiolerio A, Stoianov R, Payot S, Locatelli I, et al. Participation in a population-based physical activity programme as an aid for smoking cessation: a randomised trial. *Tobacco control* 2010 Dec;19(6):488-94.

Borrelli B, Spring B, Niaura R, Kristeller J, Ockene JK, Keuthen NJ. Weight suppression and weight rebound in ex-smokers treated with fluoxetine. *J Consult Clin Psychol* 1999 Feb;67(1):124-31.

Bouchard C, Tremblay A, Despres JP, Theriault G, Nadeau A, Lupien PJ, et al. The response to exercise with constant energy intake in identical twins. *Obes Res* 1994 Sep;2(5):400-10.

Bowden RG, Lanning BA, Doyle EI, Slonaker B, Johnston HM, Scanes G. The effects of weight loss attempts, exercise initiation, and dietary practices on health related quality of life. *Applied Research in Quality of Life* 2008 Jun;3(2):149-60.

Bravis V, Hui E, Salih S, Mehar S, Hassanein M, Devendra D. Ramadan Education and Awareness in Diabetes (READ) programme for Muslims with type 2 diabetes who fast during Ramadan. *Diabet Med* 2010;27(3):327-31.

Campbell WW, Crim MC, Young VR, Evans WJ. Increased energy requirements and changes in body composition with resistance training in older adults. *Am J Clin Nutr* 1994 Aug;60(2):167-75.

Catenacci VA, Grunwald GK, Ingebrigtsen JP, Jakicic JM, McDermott MD, Phelan S, et al. Physical activity patterns using accelerometry in the National Weight Control Registry. *Obesity* 2011 Jun;19(6):1163-70.

Cederberg H, Rajala U, Koivisto VM, Jokelainen J, Surcel HM, Keinänen-Kiukaanniemi S, et al. Unacylated ghrelin is associated with changes in body composition and body fat distribution during long-term exercise intervention. *European Journal of Endocrinology* 2011 Aug;165(2):243-8.

Cederberg H, Mikkola I, Jokelainen J, Laakso M, Harkonen P, Ikaheimo T, et al. Exercise during military training improves cardiovascular risk factors in young men. *Atherosclerosis* 2011 Jun;216(2):489-95.

Claesson AL, Holm G, Ernerson A, Lindstrom T, Nystrom FH. Two weeks of overfeeding with candy, but not peanuts, increases insulin levels and body weight. *Scandinavian Journal of Clinical & Laboratory Investigation* 2009;69(5):598-605.

DeVahl J, King R, Williamson JW. Academic incentives for students can increase participation in and effectiveness of a physical activity program. *J AM COLL HEALTH* 2005 May;53(6):295-8.

Dewey KG, Lovelady CA, Nommsen-Rivers LA, McCrory MA, Lonnerdal B. A randomized study of the effects of aerobic exercise by lactating women on breast-milk volume and composition. *New Engl J Med* 1994 Feb 17;330(7):449-53.

Diabetes Prevention Program Research Group, Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study.[Erratum appears in *Lancet*. 2009 Dec 19;374(9707):2054]. *Lancet* 2009 Nov 14;374(9702):1677-86.

Doherty K, Militello FS, Kinnunen T, Garvey AJ. Nicotine gum dose and weight gain after smoking cessation. *J Consult Clin Psychol* 1996 Aug;64(4):799-807.

Doyle J, Severance-Fonte T, Morandi-Matricaria E, Wogen J, Frech-Tamas F. Improved blood pressure control among School bus drivers with hypertension. *Popul Health Manag* 2010;13(2):97-103.

Du H, van der AD, Boshuizen HC, Forouhi NG, Wareham NJ, Halkjaer J, et al. Dietary fiber and subsequent changes in body weight and waist circumference in European men and women. *Am J Clin Nutr* 2010 Feb;91(2):329-36.

Espeland MA, Stefanick ML, Kritz-Silverstein D, Fineberg SE, Waclawiw MA, James MK, et al. Effect of postmenopausal hormone therapy on body weight and waist and hip girths. *J Clin Endocrinol Metab* 1997;82(5):1549-56.

Frankel AJ, Birkimer JC, Brown JH, Cunningham GK. A behavioral diet on network television. *Behavioral Counseling & Community Interventions* 1983;3(1):91-101.

Garthe I, Raastad T, Refsnes PE, Koivisto A, Sundgot-Borgen J. Effect of two different weight-loss rates on body composition and strength and power-related performance in elite athletes. *INT J SPORT NUTR EXERC METAB* 2011 Apr;21(2):97-104.

Garthe I, Raastad T, Sundgot-Borgen J. Long-term effect of weight loss on body composition and performance in elite athletes. *INT J SPORT NUTR EXERC METAB* 2011 Oct;21(5):426-35.

Golan R, Schwarzfuchs D, Stampfer MJ, Shai I. Halo effect of a weight-loss trial on spouses: the DIRECT-Spouse study. *Public Health Nutr* 2010;13(4):544-9.

Groeneveld IF, Proper KI, van der Beek AJ, Hildebrandt VH, van MW. Short and long term effects of a lifestyle intervention for construction workers at risk for cardiovascular disease: a randomized controlled trial. *BMC Public Health* 2011;11:836.

Grossman LD, Parlan G, Bailey AL, Yee G, Yu M, Chan JY. Tolerability outcomes of a multicenter, observational, open-label, drug-surveillance study in patients with type 2 diabetes mellitus treated with pioglitazone for 2 years. *Clin Ther* 2009 Jan;31(1):74-88.

Hall SM, Tunstall CD, Vila KL, Duffy J. Weight gain prevention and smoking cessation: cautionary findings. *Am J Public Health* 1992 Jun;82(6):799-803.

Hallas J, Bjerrum L, Stovring H, Andersen M. Use of a prescribed ephedrine/caffeine combination and the risk of serious cardiovascular events: a registry-based case-crossover study. *Am J Epidemiol* 2008 Oct 15;168(8):966-73.

Harrell JS, Johnston LF, Griggs TR, Schaefer P, Carr EG, Jr., McMurray RG, et al. An occupation based physical activity intervention program: improving fitness and decreasing obesity. *AAOHN J* 1996 Aug;44(8):377-84.

Hemo B, Endevelt R, Porath A, Stampfer MJ, Shai I. Adherence to weight loss medications; post-marketing study from HMO pharmacy data of one million individuals. *DIABETES RES CLIN PRACT* 2011 Nov;94(2):269-75.

Hensley RD, Jones AK, Williams AG, Willsher LB, Cain PP. One-year clinical outcomes for Louisiana residents diagnosed with type 2 diabetes and hypertension. *J AM ACAD NURSE PRACT* 2005 Sep;17(9):363-9.

Hofer T, Fontana L, Anton SD, Weiss EP, Villareal D, Malayappan B, et al. Long-term effects of caloric restriction or exercise on DNA and RNA oxidation levels in white blood cells and urine in humans. *Rejuvenation Research* 2008 Aug;11(4):793-9.

Iwamoto Y, Taniguchi T, Nonaka K, Okamoto T, Okuyama K, Arjona Ferreira JC, et al. Dose-ranging efficacy of sitagliptin, a dipeptidyl peptidase-4 inhibitor, in Japanese patients with type 2 diabetes mellitus. *Endocr J* 2010;57(5):383-94.

Jacobi C, Morris L, Beckers C, Bronisch-Holtze J, Winter J, Winzelberg AJ, et al. Maintenance of internet-based prevention: A randomized controlled trial. *Int J Eating Disord* 2007;40(2):114-9.

Jenkins DJ, Kendall CW, Marchie A, Josse AR, Nguyen TH, Faulkner DA, et al. Almonds reduce biomarkers of lipid peroxidation in older hyperlipidemic subjects. *J Nutr* 2008 May;138(5):908-13.

Jorde R, Sneve M, Emaus N, Figenschau Y, Grimnes G. Cross-sectional and longitudinal relation between serum 25-hydroxyvitamin D and body mass index: The Tromso study. *EUR J NUTR* 2010;49(7):401-7.

Kang JY, Cho SW, Sung SH, Park YK, Paek YM, Choi TI. Effect of a continuous diabetes lifestyle intervention program on male workers in Korea. *Diabetes Res Clin Pract* 2010;90(1):26-33.

Kasai M, Nosaka N, Maki H, Negishi S, Aoyama T, Nakamura M, et al. Effect of dietary medium- and long-chain triacylglycerols (MLCT) on accumulation of body fat in healthy humans. *Asia Pac J Clin Nutr* 2003;12(2):151-60.

Kawachi I, Troisi RJ, Rotnitzky AG, Coakley EH, Colditz GA. Can physical activity minimize weight gain in women after smoking cessation? *Am J Public Health* 1996 Jul;86(7):999-1004.

Killen JD, Fortmann SP, Newman B. Weight change among participants in a large sample minimal contact smoking relapse prevention trial. *Addict Behav* 1990;15(4):323-32.

Klem ML, Viteri JE, Wing RR. Primary prevention of weight gain for women aged 25-34: the acceptability of treatment formats. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 2000 Feb;24(2):219-25.

Koenigstorfer J, Schmidt WF. Effects of exercise training and a hypocaloric diet on female monozygotic twins in free-living conditions. *Physiol Behav* 2011 Oct 24;104(5):838-44.

Koszewski WM, Kuo M. Factors that influence the food consumption behavior and nutritional adequacy of college women. *J AM DIET ASSOC* 1996;96(12):1286-8.

Koulouri AA, Tigbe WW, Lean ME. The effect of advice to walk 2000 extra steps daily on food intake. *J HUM NUTR DIET* 2006 Aug;19(4):263-6.

Kutlubay Z. Evaluation of mesotherapeutic injections of three different combinations of lipolytic agents for body contouring. *Journal of Cosmetic & Laser Therapy* 2011 Aug;13(4):142-53.

Kwak L, Kremers SPJ, Candel MJJM, Visscher TLS, Brug J, van Baak MA. Changes in skinfold thickness and waist circumference after 12 and 24 months resulting from the NHF-NRG In Balance-project. *Int j behav nutr phys act* 2010;7.

Lee JW, Lee HR, Shim JY, Im JA, Lee DC. Abdominal visceral fat reduction is associated with favorable changes of serum retinol binding protein-4 in nondiabetic subjects. *Endocr J* 2008 Oct;55(5):811-8.

Levine MD, Perkins KA, Kalarchian MA, Cheng Y, Houck PR, Slane JD, et al. Bupropion and cognitive behavioral therapy for weight-concerned women smokers. *Arch Intern Med* 2010 Mar 22;170(6):543-50.

Levitsky DA, Garay J, Nausbaum M, Neighbors L, Dellavalle DM. Monitoring weight daily blocks the freshman weight gain: a model for combating the epidemic of obesity. *Int J Obes* 2006 Jun;30(6):1003-10.

Levitsky DA, DeRosimo L. One day of food restriction does not result in an increase in subsequent daily food intake in humans. *Physiol Behav* 2010;99(4):495-9.

Lingfors H, Lindstrom K, Persson LG, Bengtsson C, Lissner L. Evaluation of "Live for Life", a health promotion programme in the County of Skaraborg, Sweden. *J Epidemiol Community Health* 2001;55(4):277-82.

Lissner L, Heitmann BL, Bengtsson C. Low-fat diets may prevent weight gain in sedentary women: prospective observations from the population study of women in Gothenburg, Sweden. *Obes Res* 1997 Jan;5(1):43-8.

Liu TC, Liu YY, Lee SD, Huang CY, Chien KY, Cheng IS, et al. Effects of short-term detraining on measures of obesity and glucose tolerance in elite athletes. *J SPORTS SCI* 2008 Jul;26(9):919-25.

Lombard C, Deeks A, Jolley D, Ball K, Teede H. A low intensity, community based lifestyle programme to prevent weight gain in women with young children: cluster randomised controlled trial. *BMJ* 2010;341:c3215.

Luepker RV, Murray DM, Jacobs J, Mittelmark MB, Bracht N, Carlaw R, et al. Community education for cardiovascular disease prevention: Risk factor changes in the Minnesota Heart Health Program. *Am J Public Health* 1994;84(9):1383-93.

Maddison R, Roberts V, Bullen C, McRobbie H, Jiang Y, Prapavessis H, et al. Design and conduct of a pragmatic randomized controlled trial to enhance smoking-cessation outcomes with exercise: The Fit2Quit study. *Ment Health Phys Act* 2010;3(2):92-101.

Manini TM, Newman AB, Fielding R, Blair SN, Perri MG, Anton SD, et al. Effects of exercise on mobility in obese and nonobese older adults. *Obesity* 2010 Jun;18(6):1168-75.

Martin CK, Han H, Anton SD, Greenway FL, Smith SR. Effect of valproic acid on body weight, food intake, physical activity and hormones: results of a randomized controlled trial. *J Psychopharmacol* 2009 Sep;23(7):814-25.

McHugh F, Lindsay GM, Hanlon P, Hutton I, Brown MR, Morrison C, et al. Nurse led shared care for patients on the waiting list for coronary artery bypass surgery: a randomised controlled trial. *Heart* 2001 Sep;86(3):317-23.

Merriam PA, Tellez TL, Rosal MC, Olendzki BC, Ma Y, Pagoto SL, et al. Methodology of a diabetes prevention translational research project utilizing a community-academic partnership for implementation in an underserved Latino community. *BMC Med Res Methodol* 2009;Vol9, pp20, 2009:20.

Merrill RM, Bowden DE, Aldana SG. Factors associated with attrition and success in a worksite wellness telephonic health coaching program. *Education for Health* 2010 Nov;23(3):385.

Milanese C, Piscitelli F, Simoni C, Pugliarello R, Zancanaro C. Effects of whole-body vibration with or without localized radiofrequency on anthropometry, body composition, and motor performance in young nonobese women. *J Altern Complement Med* 2012;18(1):69-75.

Miles A, Rapoport L, Wardle J, Afuape T, Duman M. Using the mass-media to target obesity: an analysis of the characteristics and reported behaviour change of participants in the BBC's 'Fighting Fat, Fighting Fit' campaign. *HEALTH EDUC RES* 2001 Jun;16(3):357-72.

Morgan CL, Jenkins-Jones S, Evans M, Barnett AH, Poole CD, Currie CJ. Weight change in people with type 2 diabetes: Secular trends and the impact of alternative antihyperglycaemic drugs. *Diabetes Obes Metab* 2012;14(5):424-32.

Mori M, Aizawa T, Tokoro M, Miki T, Yamori Y. Soy isoflavone tablets reduce osteoporosis risk factors and obesity in middle-aged Japanese women. *Clinical & Experimental Pharmacology & Physiology* 2004 Dec;31:Suppl-41.

Newton RL, Jr., Han H, Stewart TM, Ryan DH, Williamson DA. Efficacy of a pilot Internet-based weight management program (H.E.A.L.T.H.) and longitudinal physical fitness data in Army Reserve soldiers. *Journal of Diabetes Science & Technology* 2011 Sep;5(5):1255-62.

Nicolino JC, Martz DM, Curtin L. Evaluation of a cognitive-behavioral therapy intervention to improve body image and decrease dieting in college women. [References]. *Eat behav* 2001;2(4):353-62.

Nides M, Rand C, Dolce J, Murray R, O'Hara P, Voelker H, et al. Weight gain as a function of smoking cessation and 2-mg nicotine gum use among middle-aged smokers with mild lung impairment in the first 2 years of the Lung Health Study. [References]. *Health Psychol* 1994 Jul;13(4):354-61.

Nilsson PM KEN. Life-style intervention at the worksite--reduction of cardiovascular risk factors in a randomized study. *Scand J Work Environ Health* 2001;27(1):57.

O'Brien KM, LeBow MD. Reducing maladaptive weight management practices: developing a psychoeducational intervention program. *Eat behav* 2007 Apr;8(2):195-210.

Pajak A, Szafraniec K, Janion M, Szpak A, Wizner B, Wolfshaut-Wolak R, et al. The impact of the Polish national Programme of Cardiovascular Disease Prevention on the quality of primary cardiovascular disease prevention in clinical practice. *Kardiol Pol* 2010;68(12):1332-41.



Perkins KA, Marcus MD, Levine MD, D'Amico D, Miller A, Broge M, et al. Cognitive-behavioral therapy to reduce weight concerns improves smoking cessation outcome in weight-concerned women. [References]. *J Consult Clin Psychol* 2001 Aug;69(4):604-13.

Persky I, Spring B, Vander Wal JS, Pagoto S, Hedeker D. Adherence across behavioral domains in treatment promoting smoking cessation plus weight control. *Health Psychol* 2005 Mar;24(2):153-60.

Petersen R, Sill S, Lu C, Young J, Edington DW. Effectiveness of employee internet-based weight management program. *Journal of Occupational & Environmental Medicine* 2008 Feb;50(2):163-71.

Peterson JM, Montgomery S, Haddad E, Kearney L, Tonstad S. Effect of consumption of dried California mission figs on lipid concentrations. *Ann Nutr Metab* 2011;58(3):232-8.

Presnell K, Stice E. An experimental test of the effect of weight-loss dieting on bulimic pathology: Tipping the scales in a different direction. [References]. *Journal of Abnormal Psychology* 2003 Feb;112(1):166-70.

Presnell K, Stice E, Tristan J. Experimental investigation of the effects of naturalistic dieting on bulimic symptoms: Moderating effects of depressive symptoms. [References]. *Appetite* 2008 Jan;50(1):91-101.

Quinn TJ, Jenkins M. Nutritional profiles of selected college females in a 15-week exercise and weight-control class. *Health Values: Health Behavior, Education & Promotion* 1991 May;15(3):34-41.

Rivera FI, Lieberman LS, Rivadeneyra GM, Sallas AM. Using a social marketing framework to transform an education program: Lessons from the Hispanic Obesity Prevention and Education (PESO) Program. [References]. *Social Marketing Quarterly* 2010 Jun;16(2):2-17.

Rohrer JE, Naessens JM, Liesinger J, Litchy W. Comparing diverse health promotion programs using overall self-rated health as a common metric. *Popul Health Manag* 2010;13(2):91-5.

Sacks G, Veerman JL, Moodie M, Swinburn B. 'Traffic-light' nutrition labelling and 'junk-food' tax: a modelled comparison of cost-effectiveness for obesity prevention. *Int J Obes* 2011 Jul;35(7):1001-9.

Sallit J, Ciccazzo M, Dixon Z. A cognitive-behavioral weight control program improves eating and smoking behaviors in weight-concerned female smokers. *J AM DIET ASSOC* 2009 Aug;109(8):1398-405.

Salopuro TM, Saaristo T, Oksa H, Puolijoki H, Vanhala M, Ebeling T, et al. Population-level effects of the national diabetes prevention programme (FIN-D2D) on the body weight, the waist circumference, and the prevalence of obesity. *BMC Public Health* 2011;11:350.

Schneider KL, Spring B, Pagoto SL. Affective benefits of exercise while quitting smoking: Influence of smoking-specific weight concern. [References]. *Psychology of Addictive Behaviors* 2007 Jun;21(2):255-60.

Schutt M, Kern W, Zimmermann A, Busch P, Kerner W, Voll A, et al. Association of antidiabetic therapies to glycemic control and to body weight in type 2 diabetes: A German multicenter analysis on 9.294 patients. *Exp Clin Endocrinol Diabetes* 2010;118(8):490-5.

Sharpe PA, Burroughs EL, Granner ML, Wilcox S, Hutto BE, Bryant CA, et al. Impact of a community-based prevention marketing intervention to promote physical activity among middle-aged women. *HEALTH EDUC BEHAV* 2010 Jun;37(3):403-23.

Simpson M, Earles J, Folen R, Trammel R, James L. The Tripler Army Medical Center's LE3AN program: a six-month retrospective analysis of program effectiveness for African-American and European-American females. *J Natl Med Assoc* 2004 Oct;96(10):1332-6.

Sneddon AA, Tsofliou F, Fyfe CL, Matheson I, Jackson DM, Horgan G, et al. Effect of a conjugated linoleic acid and omega-3 fatty acid mixture on body composition and adiponectin. *Obesity (Silver Spring, Md)* 2008;16(5):1019-24.

Sol BG, van der Graaf Y, van der Bijl JJ, Goessens BM, Visseren FL. The role of self-efficacy in vascular risk factor management: a randomized controlled trial. *PATIENT EDUC COUNS* 2008 May;71(2):191-7.

Sood A, Ebbert JO, Clark MM, Croghan IT, Schroeder DR, Hays JT. Sibutramine for weight gain attenuation during smoking cessation with varenicline: a pilot study. *Nicotine & Tobacco Research* 2009 Dec;11(12):1479-84.

Sorensen G, Stoddard A, Quintiliani L, Ebbeling C, Nagler E, Yang M, et al. Tobacco use cessation and weight management among motor freight workers: results of the gear up for health study. *Cancer Causes & Control* 2010 Dec;21(12):2113-22.

Spring B, Wurtman J, Wurtman R, el-Khoury A, Goldberg H, McDermott J, et al. Efficacies of dexfenfluramine and fluoxetine in preventing weight gain after smoking cessation. *Am J Clin Nutr* 1995 Dec;62(6):1181-7.

Spring B, Pagoto S, Pingitore R, Doran N, Schneider K, Hedeker D. Randomized controlled trial for behavioral smoking and weight control treatment: effect of concurrent versus sequential intervention. *Journal of Consulting & Clinical Psychology* 2004 Oct;72(5):785-96.

Suzuki T, Kohro T, Hayashi D, Yamazaki T, Nagai R. Frequency and impact of lifestyle modification in patients with coronary artery disease: The Japanese Coronary Artery Disease (JCAD) Study. *Am Heart J* 2012;163(2):268-73.

Suzuki T, Oba K, Igari Y, Watanabe K, Matsumura N, Futami-Suda S, et al. Effects of bile-acid-binding resin (colestimide) on blood glucose and visceral fat in Japanese patients with type 2 diabetes mellitus and hypercholesterolemia: An open-label, randomized, case-control, crossover study. *J Diabetes Complications* 2012;26(1):34-9.

Tayebi SM, Niaki AG, Hanachi P, Ghaziani FGA. The effect of Ramadan fasting and weight-lifting training on plasma volume, glucose and lipids profile of male weight-lifters. *Iran J Basic Med Sci* 2010;13(2 SPRING):57-62.

Taylor CB, Fortmann SP, Flora J, Kayman S, Barrett DC, Jatulis D, et al. Effect of long-term community health education on body mass index. The Stanford Five-City Project. *Am J Epidemiol* 1991 Aug 1;134(3):235-49.

Thomas EL, Makwana A, Newbould R, Rao AW, Gambarota G, Frost G, et al. Pragmatic study of orlistat 60[THIN SPACE]mg on abdominal obesity. *Eur J Clin Nutr* 2011 Nov;65(11):1256-62.

Toll BA, White M, Wu R, Meandzija B, Jatlow P, Makuch R, et al. Low-dose naltrexone augmentation of nicotine replacement for smoking cessation with reduced weight gain: a randomized trial. *DRUG ALCOHOL DEPENDENCE* 2010 Oct 1;111(3):200-6.

Tomiyama AJ, Mann T, Vinas D, Hunger JM, Dejager J, Taylor SE. Low calorie dieting increases cortisol. *Psychosom Med* 2010;72(4):357-64.

Tonstad S, Pometta D, Erkelens DW, Ose L, Moccetti T, Schouten JA, et al. The effect of the gastrointestinal lipase inhibitor, orlistat, on serum lipids and lipoproteins in patients with primary hyperlipidaemia. *Eur J Clin Pharmacol* 1994;46(5):405-10.

Udagawa K, Miyoshi M, Yoshiike N. Mid-term evaluation of "Health Japan 21": focus area for the nutrition and diet. *Asia Pac J Clin Nutr* 2008;17:Suppl-52.

Ussher M, West R, McEwen A, Taylor A, Steptoe A. Efficacy of exercise counselling as an aid for smoking cessation: A randomized controlled trial. [References]. *Addiction* 2003 Apr;98(4):523-32.

Verheijden MW, van Dommelen P, van Empelen P, Crone MR, Werkman AM, van Kesteren NMC. Changes in self-reported energy balance behaviours and body mass index during a mass media campaign. [References]. *Fam Pract* 2012;29(Suppl 1):i75-81.

Wammes B, Oenema A, Brug J. The evaluation of a mass media campaign aimed at weight gain prevention among young Dutch adults. *Obesity* 2007 Nov;15(11):2780-9.

Webb JB, Hardin AS. A preliminary evaluation of BMI status in moderating changes in body composition and eating behavior in ethnically-diverse first-year college women. *Eat behav* 2012;13(4):402-5.

Williams L, Germov J, Young A. Preventing weight gain: a population cohort study of the nature and effectiveness of mid-age women's weight control practices. *Int J Obes* 2007 Jun;31(6):978-86.

Williams NI, Reed JL, Leidy HJ, Legro RS, De Souza MJ. Estrogen and progesterone exposure is reduced in response to energy deficiency in women aged 25-40 years. *Hum Reprod* 2010 Sep;25(9):2328-39.

Wilson DK, Trumpeter NN, St George SM, Coulon SM, Griffin S, Lee Van HM, et al. An overview of the "Positive Action for Today's Health" (PATH) trial for increasing walking in low income, ethnic minority communities. *Contemp Clin Trials* 2010 Nov;31(6):624-33.

Ybarra J, Blanco-Vaca F, Fernandez S, Castellvi A, Bonet R, Palomer X, et al. The effects of liposuction removal of subcutaneous abdominal fat on lipid metabolism are independent of insulin sensitivity in normal-overweight individuals. *Obes Surg* 2008 Apr;18(4):408-14.

Yeh CH, Yeh SC. Effects of ear points' pressing on parameters related to obesity in non-obese healthy and obese volunteers. *Journal of alternative and complementary medicine (New York, NY)* 2008;14(3):309-14.

Yoshida H, Yanai H, Ito K, Tomono Y, Koikeda T, Tsukahara H, et al. Administration of natural astaxanthin increases serum HDL-cholesterol and adiponectin in subjects with mild hyperlipidemia. *Atherosclerosis* 2010 Apr;209(2):520-3.

Zhang Z, Lanza E, Kris-Etherton PM, Colburn NH, Bagshaw D, Rovine MJ, et al. A high legume low glycemic index diet improves serum lipid profiles in men. *Lipids* 2010 Sep;45(9):765-75.

Zhang Z, Lanza E, Ross AC, Albert PS, Colburn NH, Rovine MJ, et al. A high-legume low-glycemic index diet reduces fasting plasma leptin in middle-aged insulin-resistant and -sensitive men. *Eur J Clin Nutr* 2011 Mar;65(3):415-8.

Zhu Y, Zhang Y, Choi HK. The serum urate-lowering impact of weight loss among men with a high cardiovascular risk profile: The Multiple Risk Factor Intervention Trial. *Rheumatology (UK)* 2010;49(12):2391-9.

### **<30 participants per arm**

Abraira C, de BM, Myscowski JW. Comparison of unmeasured versus exchange diabetic diets in lean adults. Body weight and feeding patterns in a 2-year prospective pilot study. *Am J Clin Nutr* 1980 May;33(5):1064-70.

Aldred S, Rohalu M. A moderate intensity exercise program did not increase the oxidative stress in older adults. *Arch Gerontol Geriatr* 2011;53(3):350-3.

Alonso A, Zozaya C, Vazquez Z, Alfredo MJ, Martinez-Gonzalez MA. The effect of low-fat versus whole-fat dairy product intake on blood pressure and weight in young normotensive adults. *J HUM NUTR DIET* 2009 Aug;22(4):336-42.

Ando D, Hosaka Y, Suzuki K, Yamagata Z. Effects of exercise training on circulating high molecular weight adiponectin and adiponectin oligomer composition: A randomized controlled trial. *J Atheroscler Thromb* 2009;16(6):733-9.

Bahadori B, McCarty MF, Barroso-Aranda J, Gustin JC, Contreras F. A "mini-fast with exercise" protocol for fat loss. *Med Hypotheses* 2009 Oct;73(4):619-22.

Barclay C, Procter KL, Glendenning R, Marsh P, Freeman J, Mathers N. Can type 2 diabetes be prevented in UK general practice? A lifestyle-change feasibility study (ISAIAH). *Br J Gen Pract* 2008 Aug;58(553):541-7.

Becker DJ, Gordon RY, Halbert SC, French B, Morris PB, Rader DJ. Red yeast rice for dyslipidemia in statin-intolerant patients: a randomized trial. *Ann Intern Med* 2009;150(12):830-9.

Biesenbach G, Bodlaj G, Sedlak M, Pieringer H, Kiesling G. Exercise program for older patients with insulin-treated type 2 diabetes: long-term effects on metabolic control and BMI. *Zeitschrift fur Gerontologie und Geriatrie* 2009 Dec;42(6):465-9.

Black DR, Coe WC, Friesen JG, Wurzmans AG. Minimal interventions for weight control: A cost-effective alternative. *Addict Behav* 1984;9(3):279-85.

Brandon LJ, Elliott-Lloyd MB. Walking, body composition, and blood pressure dose-response in African American and white women. *Ethn Dis* 2006;16(3):675-81.

Bray GA, Smith SR, de JL, Xie H, Rood J, Martin CK, et al. Effect of dietary protein content on weight gain, energy expenditure, and body composition during overeating: a randomized controlled trial. *JAMA* 2012 Jan 4;307(1):47-55.

Brown WJ, Lee C, Oyomopito R. Effectiveness of a bilingual heart health program for Greek-Australian women. *HEALTH PROMOT INT* 1996;11(2):117-25.

Bunyard LB, Katzell LI, Busby-Whitehead MJ, Wu Z, Goldberg AP. Energy requirements of middle-aged men are modifiable by physical activity. *Am J Clin Nutr* 1998 Nov;68(5):1136-42.

Burani J, Longo PJ. Low-glycemic index carbohydrates: an effective behavioral change for glycemic control and weight management in patients with type 1 and 2 diabetes. *DIABETES EDUC* 2006 Jan;32(1):78-88.

Caplan GA, Colagiuri R, Lord SR, Colagiuri S. Exercise in older people with type II diabetes maintains bone density despite weight loss. *Australasian Journal on Ageing* 1995 May;14(2):71-5.

Cho JK, Lee SH, Lee JY, Kang HS. Randomized controlled trial of training intensity in adiposity. *INT J SPORTS MED* 2011;32(6):468-75.

Craigie AM, Barton KL, Macleod M, Williams B, van TE, Belch JJF, et al. A feasibility study of a personalised lifestyle programme (HealthForce) for individuals who have participated in cardiovascular risk screening. *PREV MED* 2011;52(5):387-9.

Damsgaard CT, Frokiaer H, Andersen AD, Lauritzen L. Fish oil in combination with high or low intakes of linoleic acid lowers plasma triacylglycerols but does not affect other cardiovascular risk markers in healthy men. *J Nutr* 2008 Jun;138(6):1061-6.

de Luis DA, Izaola O, Aller R, Cuellar L, Terroba MC, Martin T, et al. A randomized clinical trial with two enteral diabetes-specific supplements in patients with diabetes mellitus type 2: metabolic effects. *Eur Rev Med Pharmacol Sci* 2008 Jul;12(4):261-6.

De GK. Increasing physical activity in Belgian type 2 diabetes patients: a three-arm randomized controlled trial. *Int J Behav Med* 2011;18(3):188.

Dickson-Parnell BE, Zeichner A. Effects of a short-term exercise program on caloric consumption. *Health Psychol* 1985;4(5):437-48.

Diedrich A, Munroe DJ, Romano M. Promoting physical activity for persons with diabetes. *DIABETES EDUC* 2010 Jan;36(1):132-40.

Dixon AN, Valsamakis G, Hanif MW, Field A, Boutsiadis A, Harte A, et al. Effect of the orlistat on serum endotoxin lipopolysaccharide and adipocytokines in South Asian individuals with impaired glucose tolerance. *INT J CLIN PRACT* 2008 Jul;62(7):1124-9.

Djuric Z, Poore KM, Depper JB, Uhley VE, Lababidi S, Covington C, et al. Methods to increase fruit and vegetable intake with and without a decrease in fat intake: compliance and effects on body weight in the nutrition and breast health study. *NUTR CANCER* 2002;43(2):141-51.

Donges CE, Duffield R, Drinkwater EJ. Effects of resistance or aerobic exercise training on interleukin-6, C-reactive protein, and body composition. *Med Sci Sports Exerc* 2010;42(2):304-13.

Drummond S, Kirk T. The effect of different types of dietary advice on body composition in a group of Scottish men. *J Hum Nutr Diet* 1998;11(6):473-85.

Fitschen PJ, Rolfhus KR, Winfrey MR, Allen BK, Manzy M, Maher MA. Cardiovascular effects of consumption of black versus English walnuts. *J med food* 2011 Sep;14(9):890-8.

Gannon MC, Hoover H, Nuttall FQ. Further decrease in glycated hemoglobin following ingestion of a LoBAG 30diet for 10 weeks compared to 5 weeks in people with untreated type 2 diabetes. *Nutr Metab* 2010;7, 2010. Article Number: 64. Date of Publication: 2010.

Goodwin GM, Fairburn CG, Cowen PJ. The effects of dieting and weight loss on neuroendocrine responses to tryptophan, clonidine, and apomorphine in volunteers: Important implications for neuroendocrine investigations in depression. *Arch Gen Psychiatry* 1987 Nov;44(11):952-7.

Gow RW, Trace SE, Mazzeo SE. Preventing weight gain in first year college students: an online intervention to prevent the "freshman fifteen". *Eat behav* 2010 Jan;11(1):33-9.

Guo W, Kawano H, Piao L, Itoh N, Node K, Sato T. Effects of aerobic exercise on lipid profiles and high molecular weight adiponectin in Japanese workers. *Intern Med* 2011;50(5):389-95.

Hankey CR, Leslie WS, Currall JE, Matthews D, Lean ME. Weight change after myocardial infarction: statistical perspectives for future study. *J HUM NUTR DIET* 2002 Dec;15(6):439-44.

Hawkes AL, Patrao TA, Green A, Aitken JF. CanPrevent: A telephone-delivered intervention to reduce multiple behavioural risk factors for colorectal cancer. *BMC Cancer* 2012;12.

Heitzmann CA, Kaplan RM, Wilson DK, Sandler J. Sex differences in weight loss among adults with Type II diabetes mellitus. *J Behav Med* 1987 Apr;10(2):197-211.

Hodson L, Harnden KE, Roberts R, Dennis AL, Frayn KN. Does the DASH diet lower blood pressure by altering peripheral vascular function? *J Hum Hypertens* 2010 May;24(5):312-9.

Johnson SW. The impact of calculated ambulation on health promotion. *ABNF J* 2005 Mar;16(2):33-8.

Josse AR, Tang JE, Tarnopolsky MA, Phillips SM. Body composition and strength changes in women with milk and resistance exercise. *Med Sci Sports Exerc* 2010;42(6):1122-30.

Kaddissy G, Lattouf N. Daily integration of the regular aquatic rhythmic activity in women's life during menopause and its role in the prevention of the metabolic syndrome. [French]. *Kinesitherapie* 2011;11(118):48-53.

Kamioka H, Nakamura Y, Okada S, Kitayuguchi J, Kamada M, Honda T, et al. Effectiveness of comprehensive health education combining lifestyle education and hot spa bathing for male white-collar employees: a randomized controlled trial with 1-year follow-up. *J Epidemiol* 2009;19(5):219-30.

Kar P, Laight D, Rooprai HK, Shaw KM, Cummings M. Effects of grape seed extract in Type 2 diabetic subjects at high cardiovascular risk: a double blind randomized placebo controlled trial examining metabolic markers, vascular tone, inflammation, oxidative stress and insulin sensitivity. *Diabet Med* 2009 May;26(5):526-31.

Kato T, Inagaki K, Sawai Y, Kanayama H, Katada N, Itoh M. Comparison of efficacy of pitavastatin and colestimide in Japanese patients with diabetes mellitus complicated by hyperlipidemia and metabolic syndrome. *Experimental & Clinical Endocrinology & Diabetes* 2011 Oct;119(9):554-8.

King S, David S, Newton H, Hevey D, Rafferty F, Horgan JH. The effect of dietary modification on the training outcome and body composition in patients undergoing a cardiac rehabilitation programme. *Coronary Health Care* 2000;4(2):76-81.

Kohrt WM, Obert KA, Holloszy JO. Exercise training improves fat distribution patterns in 60- to 70-year-old men and women. *Journals of Gerontology* 1992;47(4):M99-M105.

Ku YH, Han KA, Ahn H, Kwon H, Koo BK, Kim HC, et al. Resistance exercise did not alter intramuscular adipose tissue but reduced retinol-binding protein-4 concentration in individuals with type 2 diabetes mellitus. *J Int Med Res* 2010 May;38(3):782-91.

Leermakers EA, Anglin K, Wing RR. Reducing postpartum weight retention through a correspondence intervention. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 1998 Nov;22(11):1103-9.

Leermakers EA, Jakicic JM, Viteri J, Wing RR. Clinic-based vs. home-based interventions for preventing weight gain in men. *Obes Res* 1998 Sep;6(5):346-52.

Leidy HJ, Dougherty KA, Frye BR, Duke KM, Williams NI. Twenty-four-hour ghrelin is elevated after calorie restriction and exercise training in non-obese women. [References]. *Obesity* 2007 Feb;15(2):446-55.

Madanmohan, Mahadevan SK, Balakrishnan S, Gopalakrishnan M, Prakash ES. Effect of six weeks yoga training on weight loss following step test, respiratory pressures, handgrip strength and handgrip endurance in young healthy subjects. *Indian Journal of Physiology & Pharmacology* 2008 Apr;52(2):164-70.

Malaguti M, Baldini M, Angeloni C, Biagi P, Hrelia S. High-protein-PUFA supplementation, red blood cell membranes, and plasma antioxidant activity in volleyball athletes. *International Journal of Sport Nutrition and Exercise Metabolism* 2008 Jun;18(3):301-12.

Manios Y, Moschonis G, Koutsikas K, Papoutsou S, Petraki I, Bellou E, et al. Changes in body composition following a dietary and lifestyle intervention trial: The postmenopausal health study. *Maturitas* 2009;62(1):58-65.

Masley SC, Weaver W, Peri G, Phillips SE. Efficacy of lifestyle changes in modifying practical markers of wellness and aging. *Altern Ther Health Med* 2008 Mar;14(2):24-9.

McCrary MA, Nommsen-Rivers LA, Mole PA, Lonnerdal B, Dewey KG. Randomized trial of the short-term effects of dieting compared with dieting plus aerobic exercise on lactation performance. *Am J Clin Nutr* 1999;69(5):959-67.

Mier N, Tanguma J, Millard AV, Villarreal EK, Alen M, Ory MG. A pilot walking program for Mexican-American women living in colonias at the border. *Am J Health Promot* 2011 Jan;25(3):172-5.

Moriguti JC, Das SK, Saltzman E, Corrales A, McCrary MA, Greenberg AS, et al. Effects of a 6-week hypocaloric diet on changes in body composition, hunger, and subsequent weight regain in healthy young and older adults. *Journals of Gerontology Series A-Biological Sciences & Medical Sciences* 2000 Dec;55(12):B580-B587.

Mougios V, Kazaki M, Christoulas K, Ziogas G, Petridou A. Does the intensity of an exercise programme modulate body composition changes? *INT J SPORTS MED* 2006 Mar;27(3):178-81.

Nagao T, Meguro S, Hase T, Otsuka K, Komikado M, Tokimitsu I, et al. A catechin-rich beverage improves obesity and blood glucose control in patients with type 2 diabetes. *Obesity* 2009 Feb;17(2):310-7.

Navas-Carretero S, Abete I, Zulet MA, Martinez JA. Chronologically scheduled snacking with high-protein products within the habitual diet in type-2 diabetes patients leads to a fat mass loss: a longitudinal study. *Nutr J* 2011;10:74.

Navas-Carretero S, Cuervo M, Abete I, Zulet MA, Martinez JA. Frequent consumption of selenium-enriched chicken meat by adults causes weight loss and maintains their antioxidant status. *Biol Trace Elem Res* 2011;143(1):8-19.

Nuttall FQ, Schweim K, Hoover H, Gannon MC. Effect of the LoBAG30 diet on blood glucose control in people with type 2 diabetes. *Br J Nutr* 2008 Mar;99(3):511-9.

Nybo L, Sundstrup E, Jakobsen MD, Mohr M, Hornstrup T, Simonsen L, et al. High-intensity training versus traditional exercise interventions for promoting health. *Med Sci Sports Exerc* 2010 Oct;42(10):1951-8.

Obara K, Mizutani M, Hitomi Y, Yajima H, Kondo K. Isohumulones, the bitter component of beer, improve hyperglycemia and decrease body fat in Japanese subjects with prediabetes. *Clinical nutrition (Edinburgh, Scotland)* 2009;28(3):278-84.

Oldroyd JC UNW. Randomised controlled trial evaluating lifestyle interventions in people with impaired glucose tolerance. *Diabetes Res Clin Pract* 2006;72(2):117.

Osterdahl M, Kocturk T, Koochek A, Wandell PE. Effects of a short-term intervention with a paleolithic diet in healthy volunteers. *Eur J Clin Nutr* 2008;62(5):682-5.

Panunzio MF, Caporizzi R, Antoniciello A, Cela EP, Ferguson LR, D'Ambrosio P. Randomized, controlled nutrition education trial promotes a Mediterranean diet and improves anthropometric, dietary, and metabolic parameters in adults. *Annali di Igiene* 2011 Jan;23(1):13-25.

Philippou E, Bovill-Taylor C, Rajkumar C, Vampa ML, Ntatsaki E, Brynes AE, et al. Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* 2009 Dec;58(12):1703-8.

Pimentel GD, Portero-McLellan KC, Oliveira EP, Spada AP, Oshiiwa M, Zemdegs JC, et al. Long-term nutrition education reduces several risk factors for type 2 diabetes mellitus in Brazilians with impaired glucose tolerance. *Nutr Res* 2010 Mar;30(3):186-90.

Pratt C, Wilson W, Leklem J, Kingsley L. Peer support and nutrition education for older adults with diabetes. *Journal of Nutrition for the Elderly* 1987;6(4):31-43.

Prewitt TE, Schmeisser D, Bowen PE, Aye P, Dolecek TA, Langenberg P, et al. Changes in body weight, body composition, and energy intake in women fed high- and low-fat diets. *Am J Clin Nutr* 1991 Aug;54(2):304-10.

Ryan AS, Pratley RE, Elahi D, Goldberg AP. Resistive training increases fat-free mass and maintains RMR despite weight loss in postmenopausal women. *J Appl Physiol* 1995 Sep;79(3):818-23.

Sato K, Kawamura T, Yamagiwa S. The "Senobi" breathing exercise is recommended as first line treatment for obesity. *Biomedical Research* 2010;31(4):259-62.

Schaefer EJ, Lichtenstein AH, Lamou-Fava S, McNamara JR, Schaefer MM, Rasmussen H, et al. Body weight and low-density lipoprotein cholesterol changes after consumption of a low-fat ad libitum diet. *JAMA* 1995 Nov 8;274(18):1450-5.

Scheid JL, De Souza MJ, Leidy HJ, Williams NI. Ghrelin but not peptide YY is related to change in body weight and energy availability. *Med Sci Sports Exerc* 2011 Nov;43(11):2063-71.

Shaw I, Shaw BS. Consequence of resistance training on body composition and coronary artery disease risk. *Cardiovascular Journal of South Africa* 2006;17(3):111-6.

Silbernagel G, Machann J, Unmuth S, Schick F, Stefan N, Haring HU, et al. Effects of 4-week very-high-fructose/glucose diets on insulin sensitivity, visceral fat and intrahepatic lipids: An exploratory trial. *Br J Nutr* 2011;106(1):79-86.

Soenen S, Westerterp-Plantenga MS. Changes in body fat percentage during body weight stable conditions of increased daily protein intake vs. control. *Physiol Behav* 2010 Dec 2;101(5):635-8.

Soenen S, Plasqui G, Smeets AJ, Westerterp-Plantenga MS. Protein intake induced an increase in exercise stimulated fat oxidation during stable body weight. *Physiol Behav* 2010 Dec 2;101(5):770-4.

Stote KS, Baer DJ, Spears K, Paul DR, Harris GK, Rumpler WV, et al. A controlled trial of reduced meal frequency without caloric restriction in healthy, normal-weight, middle-aged adults. *Am J Clin Nutr* 2007 Apr;85(4):981-8.

Strychar I, Cohn JS, Renier G, Rivard M, Aris-Jilwan N, Beauregard H, et al. Effects of a diet higher in carbohydrate/lower in fat versus lower in carbohydrate/higher in monounsaturated fat on postmeal triglyceride concentrations and other cardiovascular risk factors in type 1 diabetes. *Diabetes Care* 2009 Sep;32(9):1597-9.



Takada A, Nakamura R, Furukawa M, Takahashi Y, Nishimura S, Kosugi S. The relationship between weight loss and time and risk preference parameters: a randomized controlled trial.[Erratum appears in J Biosoc Sci. 2011 Jul;43(4):505-8]. J Biosoc Sci 2011 Jul;43(4):481-503.

Thomazella MC GMACD. Effects of high adherence to mediterranean or low-fat diets in medicated secondary prevention patients. Am J Cardiol 2011;108(11):1523.

Tonstad S, Alm CS, Sandvik E. Effect of nurse counselling on metabolic risk factors in patients with mild hypertension: a randomised controlled trial. EUR J CARDIOVASC NURS 2007;6(2):160-4.

Trapp EG, Chisholm DJ, Freund J, Boutcher SH. The effects of high-intensity intermittent exercise training on fat loss and fasting insulin levels of young women. Int J Obes 2008 Apr;32(4):684-91.

Veverka DV, Anderson J, Auld GW, Coulter GR, Kennedy C, Chapman PL. Use of the Stages of Change Model in Improving Nutrition and Exercise Habits in Enlisted Air Force Men. [References]. Mil Med 2003 May;168(5):373-9.

Villareal DT, Holloszy JO. Effect of DHEA on abdominal fat and insulin action in elderly women and men: a randomized controlled trial. JAMA : the journal of the American Medical Association 2004;292(18):2243-8.

Viskaal-van DM, Kok FJ, de GC. Effects of snack consumption for 8 weeks on energy intake and body weight. Int J Obes 2010 Feb;34(2):319-26.

Weiss EP, Villareal DT, Racette SB, Steger-May K, Premachandra BN, Klein S, et al. Caloric restriction but not exercise-induced reductions in fat mass decrease plasma triiodothyronine concentrations: a randomized controlled trial. Rejuvenation Research 2008 Jun;11(3):605-9.

Wieczorek-Baranowska A, Nowak A, Pilaczynska-Szczesniak L. Osteocalcin and glucose metabolism in postmenopausal women subjected to aerobic training program for 8 weeks. Metabolism 2012;61(4):542-5.

Yamashiro T, Nishikawa T, Isami S, Wei C-N, Fukumoto K, Matsuo H, et al. The effect of group-based lifestyle interventions on risk factors and insulin resistance in subjects at risk for metabolic syndrome: The Tabaruzaka Study 1. Diabestes Obes Metab 2010;12(9):790-7.

Zachariah S, Sheldon B, Shojaee-Moradie F, Jackson NC, Backhouse K, Johnsen S, et al. Insulin detemir reduces weight gain as a result of reduced food intake in patients with type 1 diabetes. Diabetes Care 2011 Jul;34(7):1487-91.

Zhang L, Feng Y, List J, Kasichayanula S, Pfister M. Dapagliflozin treatment in patients with different stages of type 2 diabetes mellitus: Effects on glycaemic control and body weight. Diabestes Obes Metab 2010;12(6):510-6.

## <12 month outcomes

Adamsson V, Reumark A, Fredriksson IB, Hammarstrom E, Vessby B, Johansson G, et al. Effects of a healthy Nordic diet on cardiovascular risk factors in hypercholesterolaemic subjects: a randomized controlled trial (NORDIET). *J Intern Med* (GBR) 2011 Feb;269(2):150-9.

Anderson-Loftin W, Barnett S, Bunn P, Sullivan P, Hussey J, Tavakoli A. Soul food light: culturally competent diabetes education. *DIABETES EDUC* 2005 Jul;31(4):555-63.

Angeles-Agdeppa I, Capanzana MV, Li-Yu J, Schollum LM, Kruger MC. High-calcium milk prevents overweight and obesity among postmenopausal women. *Food & Nutrition Bulletin* 2010 Sep;31(3):381-90.

Aoun S, Osseiran-Moisson R, Collins F, Newton R, Newton M. A self-management concept for men at the community level: The 'waist' disposal challenge. *J HEALTH PSYCHOL* 2009 Jul;14(5):663-74.

Arikawa AY, Thomas W, Schmitz KH, Kurzer MS. Sixteen weeks of exercise reduces C-reactive protein levels in young women. *Med Sci Sports Exerc* 2011 Jun;43(6):1002-9.

Barone BB, Wang NY, Bacher AC, Stewart KJ. Decreased exercise blood pressure in older adults after exercise training: contributions of increased fitness and decreased fatness. *Br J Sports Med* 2009 Jan;43(1):52-6.

Becker DJ, Gordon RY, Morris PB, Yorko J, Gordon YJ, Li M, et al. Simvastatin vs therapeutic lifestyle changes and supplements: randomized primary prevention trial. *Mayo Clin Proc* 2008 Jul;83(7):758-64.

Bedard A, Goulet J, Riverin M, Lamarche B, Lemieux S. Effects of a dietary intervention promoting the adoption of a Mediterranean food pattern on fast-food consumption among healthy French-Canadian women. *Br J Nutr* 2010 Dec;104(11):1662-5.

Boschi V, Siervo M, Nasti G, Trapanese E, D'Orsi P, Augelli E, et al. Interdisciplinary treatment of a female outpatient population. Organizational model and preliminary results. *Eating Weight Disord* 2002 Dec;7(4):268-75.

Bull FC, Jamrozik K, Blanksby BA. Tailored advice on exercise - Does it make a difference. *Am J Prev Med* 1999;16(3):230-9.

Carter-Edwards L, Bastian LA, Schultz M, Amamoo MA, Ostbye T. An internet-based weight loss intervention initiated by a newspaper. *Prev Chronic Dis* 2009 Jul;6(3):A101.

Chan CB, Ryan DAJ, Tudor-Locke C. Health benefits of a pedometer-based physical activity intervention in sedentary workers. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 2004 Dec;39(6):1215-22.

Charles C, Yuskavage J, Carlson O, John M, Tagalicud AS, Maggio M, et al. Effects of high-dose isoflavones on metabolic and inflammatory markers in healthy postmenopausal women. *Menopause (New York, NY)* 2009;16(2):395-400.

Cholewa S, Irwin JD. Project IMPACT: Brief report on a pilot programme promoting physical activity among university students. *J HEALTH PSYCHOL* 2008;13(8):1207-12.

Ciampolini M, Lovell-Smith D, Sifone M. Sustained self-regulation of energy intake. Loss of weight in overweight subjects. Maintenance of weight in normal-weight subjects. *Nutr Metab* 2010;7, 2010. Article Number: 4. Date of Publication: 2010.

Connolly S, Holden A, Turner E, Fiumicelli G, Stevenson J, Hunjan M, et al. MyAction: An innovative approach to the prevention of cardiovascular disease in the community. *Br J Cardiol* 2011;18(4):171-6.

de VE, Oenema A, Sheeran P, Brug J. Should implementation intentions interventions be implemented in obesity prevention: The impact of if-then plans on daily physical activity in Dutch adults. *International Journal of Behavioral Nutrition and Physical Activity* 2009;6:11.

Di Mauro S, Spallina G, Leotta C, Giardina M, Di Fazio I, Malaguarnera M, et al. The effects of caloric restriction and controlled physical exercise on hypertension in the elderly. *Arch Gerontol Geriatr* 1998 Jul;27(1):1-8.

Donnelly JE, Sullivan DK, Smith BK, Jacobsen DJ, Washburn RA, Johnson SL, et al. Alteration of dietary fat intake to prevent weight gain: Jayhawk Observed Eating Trial. *Obesity* 2008 Jan;16(1):107-12.

Estabrooks PA, Smith-Ray RL. Piloting a behavioral intervention delivered through interactive voice response telephone messages to promote weight loss in a pre-diabetic population. *Patient Education and Counseling* 2008;72(1):34-41.

Finucane FM, Horton J, Purslow LR, Savage DB, Brage S, Bess H, et al. A trial of aerobic exercise to reduce metabolic risk in older people: The Hertfordshire birth Cohort physical activity. *Diabetes* 2009;Vol58, ppAbstract no: 1067-P, 2009:Abstract-P.

Finucane FM, Sharp SJ, Purslow LR, Horton K, Horton J, Savage DB, et al. The effects of aerobic exercise on metabolic risk, insulin sensitivity and intrahepatic lipid in healthy older people from the Hertfordshire Cohort Study: A randomised controlled trial. *Diabetologia* 2010;53(4):624-31.

Fujii H, Watanabe Y, Ueki A, Ohno A, Kato M, Kondo K, et al. An increased dose of insulin detemir improves glycaemic control and reduces body weight of Japanese patients with diabetes. *INT J CLIN PRACT* 2010 Oct;64(11):1512-9.

Gellert KS, Aubert RE, Mikami JS. Ke 'Ano Ola: Moloka'i's community-based healthy lifestyle modification program. *Am J Public Health* 2010 May;100(5):779-83.

Gieck DJ, Olsen S. Holistic wellness as a means to developing a lifestyle approach to health behavior among college students. [References]. *J AM COLL HEALTH* 2007 Jul;56(1):29-35.

Gotsis M, Wang H, Spruijt-Metz D, Jordan-Marsh M, Valente TW. Wellness partners: design and evaluation of a web-based physical activity diary with social gaming features for adults. *JMIR Res Protoc* 2013;2(1):e10.

Goulet J, Lamarche B, Nadeau G, Lemieux S. Effect of a nutritional intervention promoting the Mediterranean food pattern on plasma lipids, lipoproteins and body weight in healthy French-Canadian women. *Atherosclerosis* 2003 Sep;170(1):115-24.

Goulet J, Lapointe A, Lamarche B, Lemieux S. Effect of a nutritional intervention promoting the Mediterranean food pattern on anthropometric profile in healthy women from the Quebec city metropolitan area. *Eur J Clin Nutr* 2007 Nov;61(11):1293-300.

Grey A, Gamble G, Ames R, Horne A, Mason B, Reid IR. Calcium supplementation does not affect CRP levels in postmenopausal women - A randomized controlled trial. *Osteoporosis Int* 2006;17(8):1141-5.

Hurling R, Catt M, De BM, Fairley BW, Hurst T, Murray P, et al. Using internet and mobile phone technology to deliver an automated physical activity program: Randomized controlled trial. *J Med Internet Res* 2007;9(2):e7.

Jacobs N, De B, I, Thijs H, Dendale P, Claes N. Effect of a cardiovascular prevention program on health behavior and BMI in highly educated adults: a randomized controlled trial. *PATIENT EDUC COUNS* 2011 Oct;85(1):122-6.

Jebb SA, Lovegrove JA, Griffin BA, Frost GS, Moore CS, Chatfield MD, et al. Effect of changing the amount and type of fat and carbohydrate on insulin sensitivity and cardiovascular risk: The RISCK (Reading, Imperial, Surrey, Cambridge, and Kings) trial. *Am J Clin Nutr* 2010;92(4):748-58.

Joo NS, Kim BT. Mobile phone short message service messaging for behaviour modification in a community-based weight control programme in Korea. [References]. *J Telemed Telecare* 2007;13(8):416-20.

Kajimoto O, Kajimoto Y, Yabune M, Nakamura T, Kotani K, Suzuki Y, et al. Tea catechins with a galloyl moiety reduce body weight and fat. *Journal of Health Science* 2005;51(2):161-71.

Katoh S, Hata S, Matsushima M, Ikemoto S, Inoue Y, Yokoyama J, et al. Troglitazone prevents the rise in visceral adiposity and improves fatty liver associated with sulfonylurea therapy - A randomized controlled trial. *Metabolism* 2001;50(4):414-7.

Kelders SM, van Gemert-Pijnen JE, Werkman A, Seydel ER. Usage and effect of a web-based intervention for the prevention of overweight; a RCT. *Studies in Health Technology & Informatics* 2010;160(Pt:1):1-32.

Khunti K, Srinivasan BT, Shutler S, Davies MJ. Effect of insulin glargine on glycaemic control and weight in obese and non-obese people with type 2 diabetes: data from the AT.LANTUS trial. *Diabetes Obes Metab* 2010 Aug;12(8):683-8.

Kim Y, Pike J, Adams H, Cross D, Doyle C, Foreyt J. Telephone intervention promoting weight-related health behaviors. *PREV MED* 2010 Mar;50(3):112-7.

Kinnunen TI, Pasanen M, Aittasalo M, Fogelholm M, Weiderpass E, Luoto R. Reducing postpartum weight retention--a pilot trial in primary health care. *Nutr J* 2007;6:21.

Kjollesdal MKR, Hjellset VT, Bjorge B, Holmboe-Ottesen G, Wandel M. Intention to change dietary habits, and weight loss among Norwegian-Pakistani women participating in a culturally adapted intervention. [References]. *Journal of Immigrant and Minority Health* 2011 Dec;13(6):1150-8.

Lim S, Choi SH, Jeong IK, Kim JH, Moon MK, Park KS, et al. Insulin-sensitizing effects of exercise on adiponectin and retinol-binding protein-4 concentrations in young and middle-aged women. *Journal of Clinical Endocrinology & Metabolism* 2008 Jun;93(6):2263-8.

Lombard CB, Deeks AA, Ball K, Jolley D, Teede HJ. Weight, physical activity and dietary behavior change in young mothers: short term results of the HeLP-her cluster randomized controlled trial. *Nutr J* 2009;8:17.

Luszczynska A, Haynes C. Changing nutrition, physical activity and body weight among student nurses and midwives: effects of a planning intervention and self-efficacy beliefs. *J HEALTH PSYCHOL* 2009 Nov;14(8):1075-84.

Ma L, Grann K, Li M, Jiang Z. A pilot study to evaluate the effect of soy isolate protein on the serum lipid profile and other potential cardiovascular risk markers in moderately hypercholesterolemic Chinese adults. *Ecol food nutr* 2011;50(6):473-85.

Makita S, Abiko A, Naganuma Y, Moriai Y, Nakamura M. Effects of telmisartan on adiponectin levels and body weight in hypertensive patients with glucose intolerance. *Metabolism* 2008 Oct;57(10):1473-8.

Mathews G, Alexander J, Rahemtulla T, Bhopal R. Impact of a cardiovascular risk control project for South Asians (Khush Dil) on motivation, behaviour, obesity, blood pressure and lipids.[Erratum appears in J Public Health (Oxf). 2008 Mar;30(1):116]. J PUBLIC HEALTH 2007 Dec;29(4):388-97.

Matsuo T, Kim MK, Murotake Y, Numao S, Kim MJ, Ohkubo H, et al. Indirect lifestyle intervention through wives improves metabolic syndrome components in men. Int J Obes 2010;34(1):136-45.

Mauceri E, Bienkowski C, Hanson KA, Doyle JJ, Jackson J, Bramer S. A health outcome assessment of the Cardio Metabolic Mission Health Program at Novartis. Journal of Occupational & Environmental Medicine 2011 Jun;53(6):647-52.

Miller-Kovach K, Hermann M, Winick M. The psychological ramifications of weight management. Journal of Women's Health & Gender-Based Medicine 1999 May;8(4):477-82.

Minges KE, Cormick G, Unglik E, Dunstan DW. Evaluation of a resistance training program for adults with or at risk of developing diabetes: an effectiveness study in a community setting. International Journal of Behavioral Nutrition & Physical Activity 2011;8:50.

Mouttapa M, Robertson TP, McEligot AJ, Weiss JW, Hoolihan L, Ora A, et al. The Personal Nutrition Planner: a 5-week, computer-tailored intervention for women. J NUTR EDUC BEHAV 2011 May;43(3):165-72.

Muniz J, Gomez-Doblas JJ, Santiago-Perez MI, Lekuona-Goya I, Murga-Eizagaetxebarria N, de Teresa-Galvan sE, et al. The effect of post-discharge educational intervention on patients in achieving objectives in modifiable risk factors six months after discharge following an episode of acute coronary syndrome, (CAM-2 Project): a randomized controlled trial. Health Qual Life Outcomes 2010;8:137.

O'Loughlin J, Paradis G, Meshefedjian G, Kishchuk N. Evaluation of an 8-week mailed healthy-weight intervention. PREV MED 1998 Mar;27(2):288-95.

Petrofsky J, Batt J, Lee B, Collins K, Yang T-N, Lemoine M, et al. The effect of an aerobic dance and diet program on cardiovascular fitness, body composition, and weight loss in women. J APPL RES 2008;8(3):179-88.

Petrofsky J, Batt J, Berk L, Bains G, Wong J, Radabaugh S, et al. A video exercise and diet program using a nutritional meal replacement shake for weight loss. J APPL RES 2011;11(1):29-36.

Petrofsky J, Batt J, Berk L, Bains G, Wong J, Radabaugh S, et al. A combination of exercise with a mini medicine ball and diet with a meal replacement shake as a synergistic program to increase fitness and produce weight loss. J APPL RES 2011;11(1):20-8.

Reisin E. The benefit of the mediterranean-style diet in patients with newly diagnosed diabetes. Curr Hypertens Rep 2010;12(2):56-8.

Rossi MCE, Nicolucci A, Di BP, Bruttomesso D, Girelli A, Ampudia FJ, et al. Diabetes interactive diary: A new telemedicine system enabling flexible diet and insulin therapy while improving quality of life: An open-label, international, multicenter, randomized study. Diabetes Care 2010;33(1):109-15.

Schmitz KH, Jensen MD, Kugler KC, Jeffery RW, Leon AS. Strength training for obesity prevention in midlife women. International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity 2003 Mar;27(3):326-33.

Seino Y, Rasmussen MF, Zdravkovic M, Kaku K. Dose-dependent improvement in glycemia with once-daily liraglutide without hypoglycemia or weight gain: A double-blind, randomized, controlled trial in Japanese patients with type 2 diabetes. Diabetes Res Clin Pract 2008 Aug;81(2):161-8.

Shai I, Erlich D, Cohen AD, Urbach M, Yosef N, Levy O, et al. The effect of personal lifestyle intervention among health care providers on their patients and clinics; the Promoting Health by Self Experience (PHASE) randomized controlled intervention trial. *PREV MED* 2012;55(4):285-91.

Shannon B, Bagby R, Wang MQ, Trenkner LL. Self-efficacy: A contributor to the explanation of eating behavior. *HEALTH EDUC RES* 1990 Dec;5(4):395-407.

Sherman BJ, Gilliland G, Speckman JL, Freund KM. The effect of a primary care exercise intervention for rural women. *PREV MED* 2007 Mar;44(3):198-201.

Simkin-Silverman L, Wing RR, Hansen DH, Klem ML. Prevention of cardiovascular risk factor elevations in healthy premenopausal women. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 1995 Sep;24(5):509-17.

Skelly AH, Carlson J, Leeman J, Soward A, Burns D. Controlled trial of nursing interventions to improve health outcomes of older African American women with type 2 diabetes. *NURS RES* 2009 Nov;58(6):410-8.

Smith AJ, Phipps WR, Arikawa AY, O'Dougherty M, Kaufman B, Thomas W, et al. Effects of aerobic exercise on premenopausal sex hormone levels: Results of the WISER study, a randomized clinical trial in healthy, sedentary, eumenorrheic women. *Cancer Epidemiology, Biomarkers & Prevention* 2011;20(6):1098-106.

Stice E, Rohde P, Shaw H, Marti CN. Efficacy trial of a selective prevention program targeting both eating disorder symptoms and unhealthy weight gain among female college students. [References]. *J Consult Clin Psychol* 2012 Feb;80(1):164-70.

Strimas R, Dionne MM. Differential effects of self-weighing in restrained and unrestrained eaters. [References]. *Personality and Individual Differences* 2010 Dec;49(8):1011-4.

Sudhakaran C, Kishore U, Anjana RM, Unnikrishnan R, Mohan V. Effectiveness of sitagliptin in Asian Indian patients with type 2 diabetes - An Indian tertiary diabetes care center experience. *Diabetes Technol Ther* 2011;13(1):27-32.

Taylor AH, Doust J, Webborn N. Randomised controlled trial to examine the effects of a GP exercise referral programme in Hailsham, East Sussex, on modifiable coronary heart disease risk factors. *J Epidemiol Community Health* 1998 Sep;52(9):595-601.

Thiesen CL, Merrill RM, Aldana SG, Diehl HA, Mahoney ML, Greenlaw RL, et al. The Coronary Health Improvement Project (CHIP) for lowering weight and improving psychosocial health. *Psychol Rep* 2011 Aug;109(1):338-52.

Tucker LA, Maxwell K. Effects of weight training on the emotional well-being and body image of females: Predictors of greatest benefit. *Am J Health Promot* 1992 May;6(5):338-44.

Tudor-Locke C, Chan CB. An Exploratory Analysis of Adherence Patterns and Program Completion of a Pedometer-Based Physical Activity Intervention. [References]. *J Phys Act Health* 2006 Apr;3(2):210-20.

Van Stralen MM, De VH, Bolman C, Mudde AN, Lechner L. Exploring the efficacy and moderators of two computer-tailored physical activity interventions for older adults: a randomized controlled trial. *Ann Behav Med* 2010 May;39(2):139-50.

Vandelanotte C, Reeves MM, Brug J, De B, I. A randomized trial of sequential and simultaneous multiple behavior change interventions for physical activity and fat intake. *PREV MED* 2008 Mar;46(3):232-7.

Ware LJ, Hurling R, Bataveljic O, Fairley BW, Hurst TL, Murray P, et al. Rates and determinants of uptake and use of an internet physical activity and weight management program in office and manufacturing work sites in England: cohort study. *J Med Internet Res* 2008;10(4):e56.

Westerterp-Plantenga MS, Wijckmans-Duijsens NE, Verboeket-van de Venne WP, de GK, van het Hof KH, Weststrate JA. Energy intake and body weight effects of six months reduced or full fat diets, as a function of dietary restraint. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 1998 Jan;22(1):14-22.

Weststrate JA, van het Hof KH, van den Berg H, Velthuis-te-Wierik EJM, de GC, Zimmermanns NJH, et al. A comparison of the effect of free access to reduced fat products or their full fat equivalents on food intake, body weight, blood lipids and fat-soluble antioxidants levels and haemostasis variables. *Eur J Clin Nutr* 1998;52(6):389-95.

Wilbur J, McDevitt JH, Wang E, Dancy BL, Miller AM, Briller J, et al. Outcomes of a home-based walking program for African-American women. *American journal of health promotion : AJHP* 2008 May;22(5):307-17.

Wilmore JH, Despres J, Stanforth PR, Mandel S, Rice T, Gagnon J, et al. Alterations in body weight and composition consequent to 20 wk of endurance training: the HERITAGE Family Study. *Am J Clin Nutr* 1999 Sep;70(3):346-52.

Winett RA, Anderson ES, Wojcik JR, Winett SG, Bowden T. Guide to health: nutrition and physical activity outcomes of a group-randomized trial of an Internet-based intervention in churches. *Ann Behav Med* 2007 Jun;33(3):251-61.

Yang W, Chen L, Ji Q, Liu X, Ma J, Tandon N, et al. Liraglutide provides similar glycaemic control as glimepiride (both in combination with metformin) and reduces body weight and systolic blood pressure in Asian population with type 2 diabetes from China, South Korea and India: a 16-week, randomized, double-blind, active control trial(\*). *Diabetes Obes Metab* 2011 Jan;13(1):81-8.

## No control group

Barnard ND, Cohen J, Jenkins DJ, Turner-McGrievy G, Gloede L, Green A, et al. A low-fat vegan diet and a conventional diabetes diet in the treatment of type 2 diabetes: a randomized, controlled, 74-wk clinical trial. *The American journal of clinical nutrition* 2009 May;89(5):1588S-96S.

Baruth M, Wilcox S, Wegley S, Buchner DM, Ory MG, Phillips A, et al. Changes in physical functioning in the Active Living Every Day program of the Active for Life Initiative. *Int J Behav Med* 2011;18(3):199-208.

Beasley JM, Schenk JM, Ludman E, Lampe JW, Reed SD, Grothaus L, et al. Brief telephone intervention increases soy intake in peri- and postmenopausal US women: the Herbal Alternatives Trial (HALT). *J AM DIET ASSOC* 2010 Aug;110(8):1189-97.

Chen L, Appel LJ, Loria C, Lin PH, Champagne CM, Elmer PJ, et al. Reduction in consumption of sugar-sweetened beverages is associated with weight loss: the PREMIER trial. *Am J Clin Nutr* 2009 May;89(5):1299-306.

Cox KL, Burke V, Beilin LJ, Puddey IB. A comparison of the effects of swimming and walking on body weight, fat distribution, lipids, glucose, and insulin in older women-the Sedentary Women Exercise Adherence Trial 2. *Metabolism* 2010;59(11):1562-73.

de Almeida-Pititto B, Griffin SJ, Sharp SJ, Hirai AT, Gimeno SGA, Ferreira SRG. A behavioral intervention in a cohort of Japanese-Brazilians at high cardiometabolic risk. *Rev Saude Publica* 2012;46(4):602-9.

Drevenhorn E, Kjellgren KI, Bengtson A. Outcomes following a programme for lifestyle changes with people with hypertension. *J CLIN NURS* 2007;16(7B):144-51.

Fujii H, Muto T, Haruyama Y, Nakade M, Kobayashi E, Ishisaki K, et al. Community-based lifestyle modification of cardiovascular disease risks in middle-aged Japanese: A 27-month update. *Tohoku J Exp Med* 2010;220(4):307-18.

Iso H. Community-based education classes for hypertension control. A 1.5-year randomized controlled trial. *Hypertension* 1996;27(4):968.

Kaku K, Rasmussen MF, Nishida T, Seino Y. Fifty-two-week, randomized, multicenter trial to compare the safety and efficacy of the novel glucagon-like peptide-1 analog liraglutide vs glibenclamide in patients with type 2 diabetes. *J Diabetes Invest* 2011;2(6):441-7.

Kawano M, Shono N, Yoshimura T, Yamaguchi M, Hirano T, Hisatomi A. Improved cardio-respiratory fitness correlates with changes in the number and size of small dense LDL: randomized controlled trial with exercise training and dietary instruction. *Intern Med* 2009;48(1):25-32.

Kemmler W, Bebenek M, Von SS, Engelke K, Kalender WA. Effect of block-periodized exercise training on bone and coronary heart disease risk factors in early post-menopausal women: A randomized controlled study. *Scand J Med Sci Sports* 2013;23(1):121-9.

Kosaka K, Noda M, Kuzuya T. Prevention of type 2 diabetes by lifestyle intervention: a Japanese trial in IGT males. *Diabetes Res Clin Pract* 2005;67(2):152-62.

Kujala UM, Jokelainen J, Oksa H, Saaristo T, Rautio N, Moilanen L, et al. Increase in physical activity and cardiometabolic risk profile change during lifestyle intervention in primary healthcare: 1-year follow-up study among individuals at high risk for type 2 diabetes. *BMJ Open* 2011;1(2):e000292.



- Liao D, Asberry PJ, Shofer JB, Callahan H, Matthys C, Boyko EJ, et al. Improvement of BMI, body composition, and body fat distribution with lifestyle modification in Japanese Americans with impaired glucose tolerance. *Diabetes Care* 2002;25(9):1504-10.
- Lindholm LH ET. The impact of health care advice given in primary care on cardiovascular risk. CELL Study Group. *BMJ (Clinical research ed )* 1995;310(6987):1105.
- Lindholm LH ET. Changes in cardiovascular risk factors by combined pharmacological and nonpharmacological strategies: the main results of the CELL Study. *J Intern Med (GBR)* 1996;240(1):13.
- Lingfors H, Lindstrom K, Persson LG, Bengtsson C, Lissner L. Lifestyle changes after a health dialogue. Results from the Live for Life health promotion programme. *SCAND J PRIM HEALTH CARE* 2003;21(4):248-52.
- Lund SS, Tarnow L, Frandsen M, Nielsen BB, Hansen BV, Pedersen O, et al. Combining insulin with metformin or an insulin secretagogue in non-obese patients with type 2 diabetes: 12 month, randomised, double blind trial. *BMJ (Clinical research ed)* 2009;Vol339, ppb4324, 2009(Clinical research ed):b4324.
- Mediano MF, Sichieri R. Insulin resistance influences weight loss in non-obese women who followed a home-based exercise program and slight caloric restriction. *DIABETES RES CLIN PRACT* 2011 Jun;92(3):361-7.
- Neve MJ, Morgan PJ, Collins CE. Behavioural factors related with successful weight loss 15 months post-enrolment in a commercial web-based weight-loss programme. *Public Health Nutr* 2012;15(7):1299-309.
- Pettee KK, Kriska AM, Conroy MB, Johnson BD, Orchard TJ, Goodpaster BH, et al. Discontinuing hormone replacement therapy: Attenuating the effect on CVD risk with lifestyle changes. *Am J Prev Med* 2007;32(6):483-9.
- Sakane N, Sato J, Tsushita K, Tsujii S, Kotani K, Tsuzaki K, et al. Prevention of type 2 diabetes in a primary healthcare setting: three-year results of lifestyle intervention in Japanese subjects with impaired glucose tolerance. *BMC Public Health* 2011;11(1):40.

## Study design

Adams JL, Nuss T, Banks C, Hartman J, Segrest W, Spears J, et al. Risk factor outcome comparison between exercise-based cardiac rehabilitation, traditional care, and an educational workshop. *J CONTIN EDUC NURS* 2007 Mar;38(2):83-8.

Almeida-Pititto B, Hirai AT, Sartorelli DS, Harima HA, Gimeno SG, Ferreira SR. Predictive factors of non-deterioration of glucose tolerance following a 2-year behavioral intervention. *Diabetol Metab Syndr* 2010;2(1).

Almeida-Pittito B, Hirai AT, Sartorelli DS, Gimeno SGA, Ferreira SRG. Impact of a 2-year intervention program on cardiometabolic profile according to the number of goals achieved. *Brazilian Journal of Medical & Biological Research* 2010;43(11):1088-94.

Anderson-Bill ES, Winett RA, Wojcik JR, Winett SG. Web-based guide to health: relationship of theoretical variables to change in physical activity, nutrition and weight at 16-months. *J Med Internet Res* 2011;13(1):e27.

Aoun S, Rosenberg M. Are rural people getting hearts smart? *Aust J Rural Health* 2004;12(2):81-8.

Bailey WA, Westman EC, Marquart ML, Guyton JR. Low glycemic diet for weight loss in hypertriglyceridemic patients attending a lipid clinic. *J clin lipidol* 2010;4(6):508-14.

Boehm G, Bracharz N, Schoberberger R. Evaluation of the sustainability of the Public Health Program "Slim without Diet (Schlank ohne Diät)". *Wien Klin Wochenschr* 2011 Jul;123(13-14):415-21.

Buysschaert M, Preumont V, Oriot PR, Paris I, Ponchon M, Scarniere D, et al. One-year metabolic outcomes in patients with type 2 diabetes treated with exenatide in routine practice. *Diabetes Metab* 2010 Nov;36(5):381-8.

Duane J, Conway W. Weight change in intensive insulin therapy for type 2 diabetes mellitus as a function of glycosylated hemoglobin (A1C) level achieved: The deep south diabetes program. *Insulin* 2008;3(4):219-31.

Fitzpatrick P, Fitz-Simon N, Lonergan M, Collins C, Daly L. Heartwatch: the effect of a primary care-delivered secondary prevention programme for cardiovascular disease on medication use and risk factor profiles. *European Journal of Cardiovascular Prevention & Rehabilitation* 2011 Feb;18(1):129-35.

Harati H, Hadaegh F, Momenan AA, Ghanei L, Bozorgmanesh MR, Ghanbarian A, et al. Reduction in incidence of type 2 diabetes by lifestyle intervention in a middle eastern community. *Am J Prev Med* 2010;38(6):628-36.

Heath GW, Wilson RH, Smith J, Leonard BE. Community-based exercise and weight control: diabetes risk reduction and glycemic control in Zuni Indians. *Am J Clin Nutr* 1991 Jun;53(6:Suppl):Suppl-1646S.

Hoffmeister H, Mensink GBM, Stolzenberg H, Hoeltz J. Reduction of coronary heart disease risk factors in the German Cardiovascular Prevention study. *Preventive Medicine: An International Journal Devoted to Practice and Theory* 1996 Mar;25(2):135-45.

Kaati G, Vester M, Sjoström M, Bygren LO. Patients attachment patterns and response to a program for alleviating cardiovascular risk factors: A 1-year follow-up of patients in a patient education program. [References]. *Patient Education and Counseling* 2001 Sep;44(3):235-41.

Linde JA, Jeffery RW, French SA, Pronk NP, Boyle RG. Self-weighing in weight gain prevention and weight loss trials. *Ann Behav Med* 2005 Dec;30(3):210-6.

May AM, Bueno-de-Mesquita HB, Boshuizen H, Spijkerman AM, Peeters PH, Verschuren WM. Effect of change in physical activity on body fatness over a 10-y period in the Doetinchem Cohort Study. *Am J Clin Nutr* 2010 Sep;92(3):491-9.

McGuire MT, Jeffery RW, French SA, Hannan PJ. The relationship between restraint and weight and weight-related behaviors among individuals in a community weight gain prevention trial. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 2001 Apr;25(4):574-80.

Sherwood NE, Jeffery RW, French SA, Hannan PJ, Murray DM. Predictors of weight gain in the Pound of Prevention study. *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity* 2000 Apr;24(4):395-403.

Watson L, Wilson BP, Alsop J, Kumar S. Weight and glycaemic control in type 2 diabetes: What is the outcome of insulin initiation? *Diabetes Obes Metab* 2011;13(9):823-31.

Whetstone LM, Kolasa KM, Dunn C, Jayaratne KS, Vodicka S, Schneider L, et al. Effects of a behavior-based weight management program delivered through a state cooperative extension and local public health department network, North Carolina, 2008-2009. *Prev Chronic Dis* 2011 Jul;8(4):A81.

Yoneda T, Shoji K, Takase H, Hibi M, Hase T, Meguro S, et al. Effectiveness and safety of 1-year ad libitum consumption of a high-catechin beverage under nutritional guidance. *Metabolic Syndrome & Related Disorders* 2009 Aug;7(4):349-56.