

Adult Obesity KQ 2 Treatment Excluded Studies List

No weight outcomes for population of interest

Bliddal H, Leeds AR, Stigsgaard L, Astrup A, and Christensen R. Weight loss as treatment for knee osteoarthritis symptoms in obese patients: 1-year results from a randomised controlled trial. *Ann Rheum Dis*. 2011;70(10):1798-803.

Camhi SM, Stefanick ML, Katzmarzyk PT, and Young DR. Metabolic syndrome and changes in body fat from a low-fat diet and/or exercise randomized controlled trial. *Obesity*. 2010;18(3):548-54.

Campbell K, Foster-Schubert K, Alfano C, Duggan C, Irwin M, and McTiernan A. Injuries in sedentary individuals enrolled in a 12-month, randomized, controlled, exercise trial. *J Phys Act Health*. 2012;9(2):198-207.

Collins CE, Morgan PJ, Jones P, Fletcher K, Martin J, Aguiar EJ, Lucas A, Neve M, McElduff P, and Callister R. Evaluation of a commercial web-based weight loss and weight loss maintenance program in overweight and obese adults: a randomized controlled trial. *BMC Public Health*. 2010;10:669.

Donnelly JE, Washburn RA, Smith BK, Sullivan DK, Gibson C, Honas JJ, and Mayo MS. A randomized, controlled, supervised, exercise trial in young overweight men and women: the Midwest Exercise Trial II (MET2). *Contemp Clin Trials*. 2012;33(4):804-10.

Felix HC, Adams B, Fausett JK, Krukowski RA, Prewitt TE, and West DS. Calculating reach of evidence-based weight loss and memory improvement interventions among older adults attending Arkansas senior centers, 2008-2011. *Prev Chronic Dis*. 2012;9:E63.

Gambineri A, Patton L, Vaccina A, Cacciari M, Morselli-Labate AM, Cavazza C, Pagotto U, and Pasquali R. Treatment with flutamide, metformin, and their combination added to a hypocaloric diet in overweight-obese women with polycystic ovary syndrome: a randomized, 12-month, placebo-controlled study. *J Clin Endocrinol Metab*. 2006;91(10):3970-80.

Harris MF, Fanaian M, Jayasinghe UW, Passey ME, McKenzie SH, Davies GP, Lyle DM, Laws RA, Schutze H, and Wan Q. A cluster randomised controlled trial of vascular risk factor management in general practice. *Med J Aust*. 2012;197(7):387-93.

Hinton PS, Rector RS, Linden MA, Warner SO, Dellsperger KC, Chockalingam A, Whaley-Connell AT, Liu Y, and Thomas TR. Weight-loss-associated changes in bone mineral density and bone turnover after partial weight regain with or without aerobic exercise in obese women. *Eur J Clin Nutr*. 2012;66(5):606-12.

Hope AA, Kumanyika SK, Shults J, and Holmes WC. Changes in Health-Related Quality of Life among African-Americans in a lifestyle weight loss program. *Quality of Life Research*. 2010;19(7):1025-33.

Kugler C, Malehsa D, Schrader E, Tegtbur U, Guetzlaff E, Haverich A, and Strueber M. A multi-modal intervention in management of left ventricular assist device outpatients: Dietary counselling, controlled exercise and psychosocial support. *Eur J Cardio-thorac Surg*. 2012;42(6):1026-32.

MacLaughlin HL, Cook SA, Kariyawasam D, Roseke M, van NM, and Macdougall IC. Nonrandomized trial of weight loss with orlistat, nutrition education, diet, and exercise in obese patients with CKD: 2-year follow-up. *Am J Kidney Dis*. 2010;55(1):69-76.

Masheb RM, Grilo CM, and Rolls BJ. A randomized controlled trial for obesity and binge eating disorder: low-energy-density dietary counseling and cognitive-behavioral therapy. *Behaviour Research & Therapy*. 2011;49(12):821-9.

Pinto AM, Kuppermann M, Nakagawa S, Vittinghoff E, Wing RR, Kusek JW, Herman WH, and Subak LL. Comparison and correlates of three preference-based health-related quality-of-life measures among overweight and obese women with urinary incontinence. *Quality of Life Research*. 2011;20(10):1655-62.

Racine EF, Lyerly J, Troyer JL, Warren-Findlow J, and McAuley WJ. The influence of home-delivered dietary approaches to stop hypertension meals on body mass index, energy intake, and percent of energy needs consumed among older adults with hypertension and/or hyperlipidemia. *J Acad Nutr Diet*. 2012;112(11):1755-62.

Radcliff TA, Bobroff LB, Lutes LD, Durning PE, Daniels MJ, Limacher MC, Janicke DM, Martin AD, and Perri MG. Comparing Costs of Telephone vs Face-to-Face Extended-Care Programs for the Management of Obesity in Rural Settings. *J Acad Nutr Diet*. 2012;112(9):1363-73.

Rickel KA, Milsom VA, Ross KM, Hoover VJ, Peterson ND, and Perri MG. Differential response of African American and Caucasian women to extended-care programs for obesity management. *Ethn Dis*. 2011;21(2):170-5.

Roffey DM, Ashdown LC, Dornan HD, Creech MJ, Dagenais S, Dent RM, and Wai EK. Pilot evaluation of a multidisciplinary, medically supervised, nonsurgical weight loss program on the severity of low back pain in obese adults. *Spine Journal: Official Journal of the North American Spine Society*. 2011;11(3):197-204.

Santonja FJ, Morales A, Villanueva RJ, and Cortes JC. Analysing the effect of public health campaigns on reducing excess weight: a modelling approach for the Spanish Autonomous Region of the Community of Valencia. *Evaluation & Program Planning*. 2012;35(1):34-9.

Schwartz AV, Johnson KC, Kahn SE, Shepherd JA, Nevitt MC, Peters AL, Walkup MP, Hodges A, Williams CC, and Bray GA. Effect of 1 year of an intentional weight loss intervention on bone mineral density in type 2 diabetes: Results from the look AHEAD randomized trial. *J Bone Miner Res*. 2012;27(3):619-27.

Shea MK, Houston DK, Nicklas BJ, Messier SP, Davis CC, Miller ME, Harris TB, Kitzman DW, Kennedy K, and Kritchevsky SB. The effect of randomization to weight loss on total mortality in older overweight and obese adults: the ADAPT Study. *The journals of gerontology*. 2010;Series A, Biological sciences and medical sciences. 65(5):519-25.

Silva MN, Markland D, Carraca EV, Vieira PN, Coutinho SR, Minderico CS, Matos MG, Sardinha LB, and Teixeira PJ. Exercise autonomous motivation predicts 3-yr weight loss in women. *Med Sci Sports Exerc*. 2011;43(4):728-37.

Tang JW, Kushner RF, Cameron KA, Hicks B, Cooper AJ, and Baker DW. Electronic tools to assist with identification and counseling for overweight patients: A randomized controlled trial. *J Gen Intern Med*. 2012;27(8):933-9.

Trolle B, Flyvbjerg A, Kesmodel U, and Lauszus FF. Efficacy of metformin in obese and non-obese women with polycystic ovary syndrome: a randomized, double-blinded, placebo-controlled cross-over trial. *Hum Reprod*. 2007;22(11):2967-73.

Trovato GM, Martines GF, Trovato FM, Pirri C, Pace P, Garozzo A, Castro A, and Catalano D. Adenovirus-36 seropositivity enhances effects of nutritional intervention on obesity, bright liver, and insulin resistance. *Digestive Diseases & Sciences*. 2012;57(2):535-44.

Vermunt PWA, Milder IEJ, Wielaard F, de Vries JHM, Baan CA, van Oers JAM, and Westert GP. A lifestyle intervention to reduce Type 2 diabetes risk in Dutch primary care: 2.5-year results of a randomized controlled trial. *Diabet Med*. 2012;29(8):e223-e231.

Wing RR, West DS, Grady D, Creasman JM, Richter HE, Myers D, Burgio KL, Franklin F, Gorin AA, Vittinghoff E, Macer J, Kusek JW, Subak LL, and Program to Reduce Incontinence by Diet and Exercise Group. Effect of weight loss on urinary incontinence in overweight and obese women: results at 12 and 18 months. *J Urol*. 2010;184(3):1005-10.

Yank V, Stafford RS, Rosas LG, and Ma J. Baseline reach and adoption characteristics in a randomized controlled trial of two weight loss interventions translated into primary care: A structured report of real-world applicability. *Contemp Clin Trials*. 2013;34(1):126-35.

Yates T, Davies MJ, Gorely T, Talbot D, Bull F, Sattar N, and Khunti K. The effect of increased ambulatory activity on markers of chronic low-grade inflammation: evidence from the PREPARE programme randomized controlled trial. *Diabet Med*. 2010;27(11):1256-63.

Intervention type does not meet criteria

Castelnuovo G, Manzoni GM, Cuzziol P, Cesa GL, Corti S, Tuzzi C, Villa V, Liuzzi A, Petroni ML, and Molinari E. TECNOB study: Ad interim results of a randomized controlled trial of a multidisciplinary telecare intervention for obese patients with type-2 diabetes. *Clin Pract Epidemiol Ment Health*. 2011;7:44-50.

Christensen JR, Faber A, Ekner D, Overgaard K, Holtermann A, and Sogaard K. Diet, physical exercise and cognitive behavioral training as a combined workplace based intervention to reduce body weight and increase physical capacity in health care workers - a randomized controlled trial. *BMC Public Health*. 2011;11:671.

Gerstel E, Pataky Z, Busnel C, Rutschmann O, Guessous I, Zumwald C, and Golay A. Impact of lifestyle intervention on body weight and the metabolic syndrome in home-care providers. *Diabetes Metab*. 2013;39(1):78-84.

Hanh T, Serog P, Fauconnier J, Batailler P, Mercier F, Roques CF, and Blin P. One-year effectiveness of a 3-week balneotherapy program for the treatment of overweight or obesity. *Evid -Based Complement Altern Med*. 2012;2012.

Jorde R, Sneve M, Torjesen P, and Figenschau Y. No improvement in cardiovascular risk factors in overweight and obese subjects after supplementation with vitamin D3 for 1 year. *J Intern Med (GBR)*. 2010;267(5):462-72.

No true control group

Ades PA, Savage PD, Lischke S, Toth MJ, Harvey-Berino J, Bunn JY, Ludlow M, and Schneider DJ. The effect of weight loss and exercise training on flow-mediated dilatation in coronary heart disease: a randomized trial. *Chest*. 2011;140(6):1420-7.

Akers JD, Cornett RA, Savla JS, Davy KP, and Davy BM. Daily self-monitoring of body weight, step count, fruit/vegetable intake, and water consumption: a feasible and effective long-term weight loss maintenance approach. *J Acad Nutr Diet*. 2012;112(5):685-92.

Andrade AM, Coutinho SR, Silva MN, Mata J, Vieira PN, Minderico CS, Melanson KJ, Baptista F, Sardinha LB, and Teixeira PJ. The effect of physical activity on weight loss is mediated by eating self-regulation. *PATIENT EDUC COUNS*. 2010;79(3):320-6.

Anton SD, LeBlanc E, Allen HR, Karabetian C, Sacks F, Bray G, and Williamson DA. Use of a computerized tracking system to monitor and provide feedback on dietary goals for calorie-restricted diets: the POUNDS LOST study. *J Diabetes Sci Technol*. 2012;6(5):1216-25.

Armamento-Villareal R, Sadler C, Napoli N, Shah K, Chode S, Sinacore DR, Qualls C, and Villareal DT. Weight loss in obese older adults increases serum sclerostin and impairs hip geometry but both are prevented by exercise training. *J Bone Miner Res*. 2012;27(5):1215-21.

Balducci S, Zanuso S, Cardelli P, Salvi L, Mazzitelli G, Bazuro A, Iacobini C, Nicolucci A, Pugliese G, and Italian Diabetes Exercise Study (IDES) Investigators. Changes in physical fitness predict improvements in modifiable cardiovascular risk factors independently of body weight loss in subjects with type 2 diabetes participating in the Italian Diabetes and Exercise Study (IDES). *Diabetes Care*. 2012;35(6):1347-54.

Barone Gibbs B, Kinzel LS, Pettee Gabriel K, Chang YF, and Kuller LH. Short- and long-term eating habit modification predicts weight change in overweight, postmenopausal women: results from the WOMAN study. *J Acad Nutr Diet*. 2012;112(9):1347-2.

Bartfield JK, Stevens VJ, Jerome GJ, Batch BC, Kennedy BM, Vollmer WM, Harsha D, Appel LJ, Desmond R, and Ard JD. Behavioral transitions and weight change patterns within the PREMIER trial. *Obesity*. 2011;19(8):1609-15.

Belalcazar LM, Reboussin DM, Haffner SM, Hoogeveen RC, Kriska AM, Schwenke DC, Tracy RP, Pi-Sunyer FX, Ballantyne CM, and Look AHEAD Research Group. A 1-year lifestyle intervention for weight loss in individuals with type 2 diabetes reduces high C-reactive protein levels and identifies metabolic predictors of change: from the Look AHEAD (Action for Health in Diabetes) study. *Diabetes Care*. 2010;33(11):2297-303.

Belalcazar LM, Ballantyne CM, Lang W, Haffner SM, Rushing J, Schwenke DC, Pi-Sunyer FX, Tracy RP, and Look Action for Health in Diabetes Research Group. Metabolic factors, adipose tissue, and plasminogen activator inhibitor-1 levels in type 2 diabetes: findings from the look AHEAD study. *Arteriosclerosis, Thrombosis & Vascular Biology*. 2011;31(7):1689-95.

Belalcazar LM, Lang W, Haffner SM, Hoogeveen RC, Pi-Sunyer FX, Schwenke DC, Balasubramanyam A, Tracy RP, Kriska AP, and Ballantyne CM. Adiponectin and the mediation of HDL-cholesterol change with improved lifestyle: The look AHEAD study. *J Lipid Res*. 2012;53(12):2726-33.

Bertoni AG, Wagenknecht LE, Kitzman DW, Marcovina SM, Rushing JT, Espeland MA, and Brain Natriuretic Peptide Subgroup of the Look AHEAD Research Group. Impact of the look AHEAD intervention on NT-pro brain natriuretic peptide in overweight and obese adults with diabetes. *Obesity (Silver Spring)*. 2012;20(7):1511-8.

Bischoff SC, Damms-Machado A, Betz C, Herpertz S, Legenbauer T, Low T, Wechsler JG, Bischoff G, Austel A, and Ellrott T. Multicenter evaluation of an interdisciplinary 52-week weight loss program for obesity with regard to body weight, comorbidities and quality of life - A prospective study. *Int J Obes.* 2012;36(4):614-24.

Borel A-L, Leblanc X, Almeras N, Tremblay A, Bergeron J, Poirier P, Despres J-P, and Series F. Sleep apnoea attenuates the effects of a lifestyle intervention programme in men with visceral obesity. *Thorax.* 2012;67(8):735-41.

Borel AL, Nazare JA, Smith J, Almeras N, Tremblay A, Bergeron J, Poirier P, and Despres JP. Improvement in insulin sensitivity following a 1-year lifestyle intervention program in viscerally obese men: contribution of abdominal adiposity. *Metabolism: Clinical & Experimental.* 2012;61(2):262-72.

Buclin-Thiebaud S, Pataky Z, Bruchez V, and Golay A. New psycho-pedagogic approach to obesity treatment: a 5-year follow-up. *PATIENT EDUC COUNS.* 2010;79(3):333-7.

Burke LE SMSSCMY. Using mHealth technology to enhance self-monitoring for weight loss: a randomized trial. *Am J Prev Med.* 2012;43(1):20-6.

Burke LE SMSSCMY. Using mHealth technology to enhance self-monitoring for weight loss: a randomized trial. *Am J Prev Med.* 2012;43(1):20-6.

Buscemi S, Batsis JA, Verga S, Carciola T, Mattina A, Citarda S, Re A, Arnone M, D'Orio L, Belmonte S, D'Angelo A, and Cerasola G. Long-term effects of a multidisciplinary treatment of uncomplicated obesity on carotid intima-media thickness. *Obesity.* 2011;19(6):1187-92.

Calugi S, Dalle GR, Compare A, Dall'Aglio E, Petroni ML, and Marchesini G. Weight loss and clinical characteristics of young adults patients seeking treatment at medical centers: Data from the QUOVADIS Study. *Eating Weight Disord.* 2012;17(4):e314-e319.

Campbell PT, Gross MD, Potter JD, Schmitz KH, Duggan C, McTiernan A, and Ulrich CM. Effect of exercise on oxidative stress: a 12-month randomized, controlled trial. *Med Sci Sports Exerc.* 2010;42(8):1448-53.

Canfi A, Gepner Y, Schwarzfuchs D, Golan R, Shahar DR, Fraser D, Witkow S, Greenberg I, Sarusi B, Vardi H, Friger M, Stampfer MJ, and Shai I. Effect of changes in the intake of weight of specific food groups on successful body weight loss during a multi-dietary strategy intervention trial. *J Am Coll Nutr.* 2011;30(6):491-501.

Chae JS, Kang R, Kwak JH, Paik JK, Kim OY, Kim M, Park JW, Jeon JY, and Lee JH. Supervised exercise program, BMI, and risk of type 2 diabetes in subjects with normal or impaired fasting glucose. *Diabetes Care.* 2012;35(8):1680-5.

Cutler L, King B, McCarthy N, Hamilton G, and Cook L. Appetite for life: an evaluation of a primary care lifestyle programme. *J Prim Health Care.* 2010;2(4):281-7.

Dalle GR, Calugi S, Petroni ML, Di DS, Marchesini G, and QUOVADIS Study Group. Weight management, psychological distress and binge eating in obesity. A reappraisal of the problem. *Appetite.* 2010;54(2):269-73.

Davis NJ, Tomuta N, Isasi CR, Leung V, and Wylie-Rosett J. Diabetes-specific quality of life after a low-carbohydrate and low-fat dietary intervention. *The Diabetes educator.* 2012;38(2):250-5.

Derosa G, Maffioli P, Ferrari I, D'Angelo A, Fogari E, Palumbo I, Randazzo S, and Cicero AF. Comparison between orlistat plus l-carnitine and orlistat alone on inflammation parameters in obese diabetic patients. *Fundamental & Clinical Pharmacology.* 2011;25(5):642-51.

Elhayany A, Lustman A, Abel R, Attal-Singer J, and Vinker S. A low carbohydrate Mediterranean diet improves cardiovascular risk factors and diabetes control among overweight patients with type 2 diabetes mellitus: a 1-year prospective randomized intervention study. *Diabetes Obes Metab*. 2010;12(3):204-9.

Faulconbridge LF, Wadden TA, Rubin RR, Wing RR, Walkup MP, Fabricatore AN, Coday M, Van DB, Mount DL, and Ewing LJ. One-year changes in symptoms of depression and weight in overweight/obese individuals with type 2 diabetes in the look AHEAD study. *Obesity*. 2012;20(4):783-93.

Ferre R, Plana N, Merino J, Aragones G, Girona J, Heras M, Coll B, Cos R, and Masana L. Effects of therapeutic lifestyle changes on peripheral artery tonometry in patients with abdominal obesity. *Nutr Metab Cardiovasc Dis*. 2012;22(2):95-102.

Fitzgibbon ML, Stolley MR, Schiffer L, Sharp LK, Singh V, and Dyer A. Obesity reduction black intervention trial (ORBIT): 18-month results. *Obesity (Silver Spring)*. 2010;18(12):2317-25.

Flood A, Mitchell N, Jaeb M, Finch EA, Laqua PS, Welsh EM, Hotop A, Langer SL, Levy RL, and Jeffery RW. Energy density and weight change in a long-term weight-loss trial. *Int j behav nutr phys act*. 2009;6:57.

Fogari R, Zoppi A, Corradi L, Preti P, Mugellini A, Lazzari P, and Derosa G. Effect of body weight loss and normalization on blood pressure in overweight non-obese patients with stage 1 hypertension. *Hypertens Res*. 2010;33(3):236-42.

Foy CG, Lewis CE, Hairston KG, Miller GD, Lang W, Jakicic JM, Rejeski WJ, Ribisl PM, Walkup MP, Wagenknecht LE, and Look AHEAD Research Group. Intensive lifestyle intervention improves physical function among obese adults with knee pain: findings from the Look AHEAD trial.[Erratum appears in *Obesity (Silver Spring)*. 2011 Jan;19(1):233]. *Obesity*. 2011;19(1):83-93.

Fuller NR, Lau NS, Denyer G, and Caterson ID. A 12-month, randomised, controlled trial to examine the efficacy of the Korean diet in an Australian overweight and obese population - A follow up analysis. *Obes Res Clin Pract*. 2012;6(4):e298-e307.

Gabriel KK, Conroy MB, Schmid KK, Storti KL, High RR, Underwood DA, Kriska AM, and Kuller LH. The impact of weight and fat mass loss and increased physical activity on physical function in overweight, postmenopausal women: results from the Women on the Move Through Activity and Nutrition study. *Menopause*. 2011;18(7):759-65.

Gagnon C, Brown C, Couture C, Kamga-Ngande CN, Hivert MF, Baillargeon JP, Carpentier AC, and Langlois MF. A cost-effective moderate-intensity interdisciplinary weight-management programme for individuals with prediabetes. *Diabetes Metab*. 2011;37(5):410-8.

Golan R, Tirosh A, Schwarzfuchs D, Harman-Boehm I, Thiery J, Fiedler GM, Bluher M, Stumvoll M, and Shai I. Dietary intervention induces flow of changes within biomarkers of lipids, inflammation, liver enzymes, and glycemic control. *Nutrition*. 2012;28(2):131-7.

Goldstein T, Kark JD, Berry EM, Adler B, Ziv E, and Raz I. The effect of a low carbohydrate energy-unrestricted diet on weight loss in obese type 2 diabetes patients - A randomized controlled trial. *e-SPEN*. 2011;6(4):e178-e186.

Goodpaster BH, DeLany JP, Otto AD, Kuller L, Vockley J, South-Paul JE, Thomas SB, Brown J, McTigue K, Hames KC, Lang W, and Jakicic JM. Effects of diet and physical activity interventions on weight loss and cardiometabolic risk factors in severely obese adults: a randomized trial. *JAMA*. 2010;304(16):1795-802.

Gregg EW, Chen H, Wagenknecht LE, Clark JM, Delahanty LM, Bantle J, Pownall HJ, Johnson KC, Safford MM, Kitabchi AE, Pi-Sunyer FX, Wing RR, Bertoni AG, and Look AHEAD Research Group. Association of an

intensive lifestyle intervention with remission of type 2 diabetes. *JAMA : the journal of the American Medical Association*. 2012;308(23):2489-96.

Gripeteg L, Torgerson J, Karlsson J, and Lindroos AK. Prolonged refeeding improves weight maintenance after weight loss with very-low-energy diets. *Br J Nutr*. 2010;103(1):141-8.

Hashemipour M, Kelishadi R, Tavalaei Zavvareh SA, and Ghatreh-Samani S. Effect of education on anthropometric indices in obese parents and children after one year of follow-up. *ARYA Atheroscler*. 2012;8(1):21-6.

Hersey JC, Khavjou O, Strange LB, Atkinson RL, Blair SN, Campbell S, Hobbs CL, Kelly B, Fitzgerald TM, Kish-Doto J, Koch MA, Munoz B, Peele E, Stockdale J, Augustine C, Mitchell G, Arday D, Kugler J, Dorn P, Ellzy J, Julian R, Grissom J, and Britt M. The efficacy and cost-effectiveness of a community weight management intervention: A randomized controlled trial of the health weight management demonstration. *PREV MED*. 2012;54(1):42-9.

Hill JO, Hauptman J, Anderson JW, Fujioka K, O'Neil PM, Smith DK, Zavoral JH, and Aronne LJ. Orlistat, a lipase inhibitor, for weight maintenance after conventional dieting: a 1-y study. *Am J Clin Nutr*. 1999;69(6):1108-16.

Hunter GR, Fisher G, Bryan DR, and Zuckerman PA. Weight loss and exercise training effect on oxygen uptake and heart rate response to locomotion. *J Strength Cond Res*. 2012;26(5):1366-73.

Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, Yukawa M, Aiello E, Potter JD, and McTiernan A. Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial. *JAMA*. 2003;289(3):323-30.

Jakicic JM, Otto AD, Lang W, Semler L, Winters C, Polzien K, and Mohr KI. The effect of physical activity on 18-month weight change in overweight adults. *Obesity*. 2011;19(1):100-9.

Jebb SA, Ahern AL, Olson AD, Aston LM, Holzapfel C, Stoll J, Amann-Gassner U, Simpson AE, Fuller NR, Pearson S, Lau NS, Mander AP, Hauner H, and Caterson ID. Primary care referral to a commercial provider for weight loss treatment versus standard care: a randomised controlled trial. *Lancet*. 2011;378(9801):1485-92.

Jolly K, Daley A, Adab P, Lewis A, Denley J, Beach J, and Aveyard P. A randomised controlled trial to compare a range of commercial or primary care led weight reduction programmes with a minimal intervention control for weight loss in obesity: the Lighten Up trial. *BMC Public Health*. 2010;Vol10, pp439, 2010:439.

Jolly K, Lewis A, Beach J, Denley J, Adab P, Deeks JJ, Daley A, and Aveyard P. Comparison of range of commercial or primary care led weight reduction programmes with minimal intervention control for weight loss in obesity: lighten Up randomised controlled trial. *BMJ*. 2011;343:d6500.

Katula JA, Vitolins MZ, Rosenberger EL, Blackwell CS, Morgan TM, Lawlor MS, and Goff DC, Jr. One-year results of a community-based translation of the Diabetes Prevention Program: Healthy-Living Partnerships to Prevent Diabetes (HELP PD) Project. *Diabetes Care*. 2011;34(7):1451-7.

Khoo J, Piantadosi C, Duncan R, Worthley SG, Jenkins A, Noakes M, Worthley MI, Lange K, and Wittert GA. Comparing effects of a low-energy diet and a high-protein low-fat diet on sexual and endothelial function, urinary tract symptoms, and inflammation in obese diabetic men. *J Sex Med*. 2011;8(10):2868-75.

Kontogianni MD, Liatis S, Grammatikou S, Perrea D, Katsilambros N, and Makrilakis K. Changes in dietary habits and their association with metabolic markers after a non-intensive, community-based lifestyle

intervention to prevent type 2 diabetes, in Greece. The DEPLAN study. *Diabetes Res Clin Pract.* 2012;95(2):207-14.

Krebs JD, Elley CR, Parry-Strong A, Lunt H, Drury PL, Bell DA, Robinson E, Moyes SA, and Mann JI. The Diabetes Excess Weight Loss (DEWL) Trial: a randomised controlled trial of high-protein versus high-carbohydrate diets over 2 years in type 2 diabetes. *Diabetologia.* 2012;55(4):905-14.

Kumanyika SK, Fassbender JE, Sarwer DB, Phipps E, Allison KC, Localio R, Morales KH, Wesby L, Harralson T, Kessler R, Tan-Torres S, Han X, Tsai AG, and Wadden TA. One-year results of the Think Health! study of weight management in primary care practices. *Obesity (Silver Spring).* 2012;20(6):1249-57.

Lapointe A, Weisnagel SJ, Provencher V, Begin C, Dufour-Bouchard AA, Trudeau C, and Lemieux S. Comparison of a dietary intervention promoting high intakes of fruits and vegetables with a low-fat approach: long-term effects on dietary intakes, eating behaviours and body weight in postmenopausal women. *Br J Nutr.* 2010;104(7):1080-90.

Leibbrandt AJ, Kieft-de Jong JC, Hogenelst MH, Snoek FJ, and Weijs PJ. Effects of the PRO-active Interdisciplinary Self-MANagement (PRISMA, Dutch DESMOND) program on dietary intake in type 2 diabetes outpatients: a pilot study. *Clin Nutr.* 2010;29(2):199-205.

Leichtle AB, Helmschrodt C, Ceglarek U, Shai I, Henkin Y, Schwarzfuchs D, Golan R, Gepner Y, Stampfer MJ, Bluher M, Stumvoll M, Thiery J, and Fiedler GM. Effects of a 2-y dietary weight-loss intervention on cholesterol metabolism in moderately obese men. *Am J Clin Nutr.* 2011;94(5):1189-95.

Levy RL, Jeffery RW, Langer SL, Graham DJ, Welsh EM, Flood AP, Jaeb MA, Laqua PS, Finch EA, Hotop AM, and Yatsuya H. Maintenance-tailored therapy vs. standard behavior therapy for 30-month maintenance of weight loss. *PREV MED.* 2010;51(6):457-9.

Look AHEAD Research Group and Wing RR. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med.* 2010;170(17):1566-75.

Malkina-Pykh IG. Effectiveness of rhythmic movement therapy for disordered eating behaviors and obesity. *Span J Psychol.* 2012;15(3):1371-87.

Milsom VA, Middleton KM, and Perri MG. Successful long-term weight loss maintenance in a rural population. *Clin Interv Aging.* 2011;6:303-9.

Morgan PJ, Lubans DR, Collins CE, Warren JM, and Callister R. 12-month outcomes and process evaluation of the SHED-IT RCT: an internet-based weight loss program targeting men. *Obesity.* 2011;19(1):142-51.

Murphy JC, McDaniel JL, Mora K, Villareal DT, Fontana L, and Weiss EP. Preferential reductions in intermuscular and visceral adipose tissue with exercise-induced weight loss compared with calorie restriction. *J Appl Physiol.* 2012;112(1):79-85.

Nackers LM, Ross KM, and Perri MG. The association between rate of initial weight loss and long-term success in obesity treatment: does slow and steady win the race? *Int J Behav Med.* 2010;17(3):161-7.

Neiberg RH, Wing RR, Bray GA, Reboussin DM, Rickman AD, Johnson KC, Kitabchi AE, Faulconbridge LF, Kitzman DW, Espeland MA, and Look AHEAD Research Group. Patterns of weight change associated with long-term weight change and cardiovascular disease risk factors in the Look AHEAD Study. *Obesity (Silver Spring).* 2012;20(10):2048-56.

Pelletier-Beaumont E, Arsenault BJ, Almeras N, Bergeron J, Tremblay A, Poirier P, and Despres J-P. Normalization of visceral adiposity is required to normalize plasma apolipoprotein B levels in response

to a healthy eating/physical activity lifestyle modification program in viscerally obese men. *Atherosclerosis*. 2012;221(2):577-82.

Phelan S, Kanaya AM, Subak LL, Hogan PE, Espeland MA, Wing RR, Burgio KL, DiLillo V, Gorin AA, West DS, Brown JS, and Look AHEAD Research Group. Weight loss prevents urinary incontinence in women with type 2 diabetes: results from the Look AHEAD trial. *J Urol*. 2012;187(3):939-44.

Rejeski WJ, Brubaker PH, Goff DC, Jr., Bearon LB, McClelland JW, Perri MG, and Ambrosius WT. Translating weight loss and physical activity programs into the community to preserve mobility in older, obese adults in poor cardiovascular health. *Arch Intern Med*. 2011;171(10):880-6.

Rejeski WJ, Ip EH, Bertoni AG, Bray GA, Evans G, Gregg EW, Zhang Q, and Look AHEAD Research Group. Lifestyle change and mobility in obese adults with type 2 diabetes. *New Engl J Med*. 2012;366(13):1209-17.

Richelsen B, Tonstad S, Rossner S, Toubro S, Niskanen L, Madsbad S, Mustajoki P, and Rissanen A. Effect of orlistat on weight regain and cardiovascular risk factors following a very-low-energy diet in abdominally obese patients: a 3-year randomized, placebo-controlled study. *Diabetes Care*. 2007;30(1):27-32.

Rock CL, Flatt SW, Sherwood NE, Karanja N, Pakiz B, and Thomson CA. Effect of a free prepared meal and incentivized weight loss program on weight loss and weight loss maintenance in obese and overweight women: a randomized controlled trial. *JAMA*. 2010;304(16):1803-10.

Rock CL, Emond JA, Flatt SW, Heath DD, Karanja N, Pakiz B, Sherwood NE, and Thomson CA. Weight loss is associated with increased serum 25-hydroxyvitamin D in overweight or obese women. *Obesity*. 2012;20(11):2296-301.

Rossen LM, Milsom VA, Middleton KR, Daniels MJ, and Perri MG. Benefits and risks of weight-loss treatment for older, obese women. *Clin Interv Aging*. 2013;8:157-66.

Saidlitz P. Effect of exercise intensity on abdominal fat loss during calorie restriction in overweight and obese postmenopausal women: A randomized, controlled trial^{1 - 4}. [French]. *Cahiers de l'Annee Gerontologique*. 2010;2(3):234-7.

Saito T, Watanabe M, Nishida J, Izumi T, Omura M, Takagi T, Fukunaga R, Bandai Y, Tajima N, Nakamura Y, Ito M, and Zensharen Study for Prevention of Lifestyle Diseases Group. Lifestyle modification and prevention of type 2 diabetes in overweight Japanese with impaired fasting glucose levels: a randomized controlled trial. *Arch Intern Med*. 2011;171(15):1352-60.

Siervo M, Nasti G, Stephan BC, Papa A, Muscariello E, Wells JC, Prado CM, and Colantuoni A. Effects of intentional weight loss on physical and cognitive function in middle-aged and older obese participants: A pilot study. *J Am Coll Nutr*. 2012;31(2):79-86.

Silva MN, Vieira PN, Coutinho SR, Minderico CS, Matos MG, Sardinha LB, and Teixeira PJ. Using self-determination theory to promote physical activity and weight control: a randomized controlled trial in women. *J Behav Med*. 2010;33(2):110-22.

Snel M, van Diepen JA, Stijnen T, Pijl H, Romijn JA, Meinders AE, Voshol P, and Jazet IM. Immediate and long-term effects of addition of exercise to a 16-week very low calorie diet on low-grade inflammation in obese, insulin-dependent type 2 diabetic patients. *Food & Chemical Toxicology*. 2011;49(12):3104-11.

Snel M, Sleddering MA, Vd Peijl ID, Romijn JA, Pijl H, Edo MA, and Jazet IM. Quality of life in type 2 diabetes mellitus after a very low calorie diet and exercise. *EUR J INTERN MED*. 2012;23(2):143-9.

Snel M, Jonker JT, Hammer S, Kerpershoek G, Lamb HJ, Meinders AE, Pijl H, de Roos A, Romijn JA, Smit JWA, and Jazet IM. Long-term beneficial effect of a 16-week very low calorie diet on pericardial fat in obese type 2 diabetes mellitus patients. *Obesity (Silver Spring)*. 2012;20(8):1572-6.

Soenen S, Bonomi AG, Lemmens SGT, Scholte J, Thijssen MAMA, Van BF, and Westerterp-Plantenga MS. Relatively high-protein or 'low-carb' energy-restricted diets for body weight loss and body weight maintenance? *Physiol Behav*. 2012;107(3):374-80.

Sorensen LB, Greve T, Kreutzer M, Pedersen U, Nielsen CM, Toubro S, and Astrup A. Weight maintenance through behaviour modification with a cooking course or neurolinguistic programming. *Can J Diet Pract Res*. 2011;72(4):181-5.

Spring B, Duncan JM, Janke EA, Kozak AT, Mcfadden HG, DeMott A, Pictor A, Epstein LH, Siddique J, Pellegrini CA, Buscemi J, and Hedeker D. Integrating technology into standard weight loss treatment: a randomized controlled trial. *JAMA Intern Med*. 2013;173(2):105-11.

Stahre L, Tarnell B, Hakanson C-E, and Hallstrom T. A randomized controlled trial of two weight-reducing short-term group treatment programs for obesity with an 18-month follow-up - duplicate of Ref ID 24302. *Int J Behav Med*. 2007;14(1):48-55.

Stewart TM, Bachand AR, Han H, Ryan DH, Bray GA, and Williamson DA. Body image changes associated with participation in an intensive lifestyle weight loss intervention. *Obesity*. 2011;19(6):1290-5.

Sukumar D, Ambia-Sobhan H, Zurfluh R, Schlussek Y, Stahl TJ, Gordon CL, and Shapses SA. Areal and volumetric bone mineral density and geometry at two levels of protein intake during caloric restriction: a randomized, controlled trial. *Journal of Bone & Mineral Research*. 2011;26(6):1339-48.

Swencionis C, Wylie-Rosett J, Lent MR, Ginsberg M, Cimino C, Wassertheil-Smoller S, Caban A, and Segal-Isaacson CJ. Weight change, psychological well-being, and vitality in adults participating in a cognitive-behavioral weight loss program. *Health Psychol*. 2013;32(4):439-46.

ter Bogt NC, Bemelmans WJ, Beltman FW, Broer J, Smit AJ, and van der Meer K. Preventing weight gain: one-year results of a randomized lifestyle intervention. *Am J Prev Med*. 2009;37(4):270-7.

Tuomilehto H, Gylling H, Peltonen M, Martikainen T, Sahlman J, Kokkarinen J, Randell J, Tukiainen H, Vanninen E, Partinen M, Tuomilehto J, Uusitupa M, Seppa J, and Kuopio Sleep Apnea Group. Sustained improvement in mild obstructive sleep apnea after a diet- and physical activity-based lifestyle intervention: postinterventional follow-up. *Am J Clin Nutr*. 2010;92(4):688-96.

Unick JL, Beavers D, Jakicic JM, Kitabchi AE, Knowler WC, Wadden TA, Wing RR, and Look AHEAD Research Group. Effectiveness of lifestyle interventions for individuals with severe obesity and type 2 diabetes: results from the Look AHEAD trial. *Diabetes Care*. 2011;34(10):2152-7.

Unick JL, Beavers D, Bond DS, Clark JM, Jakicic JM, Kitabchi AE, Knowler WC, Wadden TA, Wagenknecht LE, Wing RR, and Look AHEAD Research Group. The long-term effectiveness of a lifestyle intervention in severely obese individuals. *Am J Med*. 2013;126(3):236-2.

Venn BJ, Perry T, Green TJ, Skeaff CM, Aitken W, Moore NJ, Mann JI, Wallace AJ, Monro J, Bradshaw A, Brown RC, Skidmore PM, Doel K, O'Brien K, Frampton C, and Williams S. The effect of increasing consumption of pulses and wholegrains in obese people: a randomized controlled trial. *J Am Coll Nutr*. 2010;29(4):365-72.

Vermunt PWA, Milder IEJ, Wielaard F, de Vries JHM, van Oers HAM, and Westert GP. Lifestyle counseling for type 2 diabetes risk reduction in Dutch primary care: results of the APHRODITE study after 0.5 and 1.5 years. *Diabetes Care*. 2011;34(9):1919-25.

Villareal DT, Chode S, Parimi N, Sinacore DR, Hilton T, Armamento-Villareal R, Napoli N, Qualls C, and Shah K. Weight loss, exercise, or both and physical function in obese older adults. *New Engl J Med*. 2011;364(13):1218-29.

Wadden TA, Neiberg RH, Wing RR, Clark JM, Delahanty LM, Hill JO, Krakoff J, Otto A, Ryan DH, Vitolins MZ, and Look AHEAD Research Group. Four-year weight losses in the Look AHEAD study: factors associated with long-term success. *Obesity*. 2011;19(10):1987-98.

Wang X, Hsu FC, Isom S, Walkup MP, Kritchevsky SB, Goodpaster BH, Church TS, Pahor M, Stafford RS, and Nicklas BJ. Effects of a 12-month physical activity intervention on prevalence of metabolic syndrome in elderly men and women. *J Gerontol A Biol Sci Med Sci*. 2012;67(4):417-24.

Wesche-Thobaben JA. The development and description of the comparison group in the Look AHEAD trial. *Clinical Trials*. 2011;8(3):320-9.

West DS, Gorin AA, Subak LL, Foster G, Bragg C, Hecht J, Schembri M, Wing RR, and Program to Reduce Incontinence by Diet and Exercise (PRIDE) Research Group. A motivation-focused weight loss maintenance program is an effective alternative to a skill-based approach. *Int J Obes*. 2011;35(2):259-69.

Winett RA, Anderson ES, Wojcik JR, Winett SG, Moore S, and Blake C. Guide to health: A randomized controlled trial of the effects of a completely web-based intervention on physical activity, fruit and vegetable consumption, and body weight. *Transl Behav Med*. 2011;1(1):165-74.

Wing RR, Lang W, Wadden TA, Safford M, Knowler WC, Bertoni AG, Hill JO, Brancati FL, Peters A, Wagenknecht L, and Look AHEAD Research Group. Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care*. 2011;34(7):1481-6.

Wycherley TP, Brinkworth GD, Keogh JB, Noakes M, Buckley JD, and Clifton PM. Long-term effects of weight loss with a very low carbohydrate and low fat diet on vascular function in overweight and obese patients: Original Article. *J Intern Med (GBR)*. 2010;267(5):452-61.

Citations from the USPSTF Review considered for Adult Obesity Review but did not meet criteria

Mensink M, Blaak EE, Corpeleijn E, Saris WH, de Bruin TW, and Feskens EJ. Lifestyle intervention according to general recommendations improves glucose tolerance. *Obes Res.* 2003;11(12):1588-96.

Mitsui T, Shimaoka K, Tsuzuku S, Kajioka T, and Sakakibara H. Gentle exercise of 40 minutes with dietary counseling is effective in treating metabolic syndrome. *Tohoku J Exp Med.* 2008;215(4):355-61.

Simkin-Silverman LR, Wing RR, Boraz MA, and Kuller LH. Lifestyle intervention can prevent weight gain during menopause: results from a 5-year randomized clinical trial. *Ann Behav Med.* 2003;26(3):212-20.

Werkman A, Hulshof PJ, Stafleu A, Kremers SP, Kok FJ, Schouten EG, and Schuit AJ. Effect of an individually tailored one-year energy balance programme on body weight, body composition and lifestyle in recent retirees: a cluster randomised controlled trial. *BMC Public Health.* 2010;10:110.