

Reference List

Excluded by Population

Abdel Razek AA, Abu Zeid MM, Bilal M, Abdel Wahab NM. Virtual CT colonoscopy versus conventional colonoscopy: a prospective study. *Hepatogastroenterology* 2005;52(66):1698-702.

Abdullah M, Rani AA, Simadibrata M, Fauzi A, Syam AF. The value of fecal tumor M2 pyruvate kinase as a diagnostic tool for colorectal cancer screening. *Acta Med Indones* 2012;44(2):94-9.

Achiam MP, Andersen LPH, Klein M, Logager V, Chabanova E, Thomsen HS, et al. Differentiation between benign and malignant colon tumors using fast dynamic gadolinium-enhanced MR colonography; a feasibility study. *Eur J Radiol* 2010;74(3):e45-e50.

Arora A, Singh P. Colonoscopy in patients 80 years of age and older is safe, with high success rate and diagnostic yield. *Gastrointest Endosc* 2004;60(3):408-13.

Arora G, Mannalithara A, Singh G, Gerson LB, Triadafilopoulos G. Risk of perforation from a colonoscopy in adults: a large population-based study. *Gastrointest Endosc* 2009;69(3 Pt 2):654-64.

Bafandeh Y, Khoshbaten M, Eftekhari Sadat AT, Farhang S. Clinical predictors of colorectal polyps and carcinoma in a low prevalence region: results of a colonoscopy based study. *World J Gastroenterol* 2008;14(10):1534-8.

Baxter NN, Warren JL, Barrett MJ, Stukel TA, Doria-Rose VP. Association between colonoscopy and colorectal cancer mortality in a US cohort according to site of cancer and colonoscopist specialty. *J Clin Oncol* 2012;30(21):2664-9.

Beitler AL, Wortmann GW, RenomDeLaBaume H, Hofmann LJ, Goff JM. Endoscopy in a deployed combat support hospital: maintaining military end-strength. *Mil Med* 2006;171(6):530-3.

Bersani G, Rossi A, Ricci G, Pollino V, DeFabritiis G, Suzzi A, et al. Do ASGE guidelines for the appropriate use of colonoscopy enhance the probability of finding relevant pathologies in an open access service? *Dig Liver Dis* 2005;37(8):609-14.

Bini EJ, Park J, Francois F. Use of flexible sigmoidoscopy to screen for colorectal cancer in HIV-infected patients 50 years of age and older. *Arch Intern Med* 2006;166(15):1626-31.

Bini EJ, Green B, Poles MA. Screening colonoscopy for the detection of neoplastic lesions in asymptomatic HIV-infected subjects. *Gut* 2009;58(8):1129-34.

Bini EJ, Rajapaksa RC, Weinshel EH. Positive predictive value of fecal occult blood testing in persons taking warfarin. *Am J Gastroenterol* 2005;100(7):1586-92.

Bini EJ, Kinkhabwala A, Goldfarb DS. Predictive value of a positive fecal occult blood test increases as the severity of CKD worsens. *Am J Kidney Dis* 2006;48(4):580-6.

Bjerregaard NC, Tottrup A, Sorensen HT, Laurberg S. Diagnostic value of self-reported symptoms in Danish outpatients referred with symptoms consistent with colorectal cancer. *Colorectal Dis* 2007;9(5):443-51.

Blumenstein I, Tacke W, Bock H, Filmann N, Lieber E, Zeuzem S, et al. Prevalence of colorectal cancer and its precursor lesions in symptomatic and asymptomatic patients undergoing total colonoscopy: Results of a large prospective, multicenter, controlled endoscopy study. *Eur J Gastroenterol Hepatol* 2013;25(5):556-61.

Borda F, Jimenez FJ, Borda A, Urman J, Goni S, Ostiz M, et al. Endoscopic localization of colorectal cancer: Study of its accuracy and possible error factors. *Rev Esp Enferm Dig* 2012;104(10):512-7.

Brenner H, Chang-Claude J, Seiler CM, Sturmer T, Hoffmeister M. Potential for colorectal cancer prevention of sigmoidoscopy versus colonoscopy: Population-based case control study. *Cancer Epidemiol Biomarkers Prev* 2007;16(3):494-9.

Burling D, Halligan S, Slater A, Noakes MJ, Taylor SA. Potentially serious adverse events at CT colonography in symptomatic patients: national survey of the United Kingdom. *Radiology* 2006;239(2):464-71.

Caldeira A, Pereira E, Baldaque-Silva F, Pereira B, Sousa R, Tristan J, et al. Role of hydrocolonic sonography in the detection of colonic neoplastic lesions. *J Diagn Med Sonogr* 2011;27(4):160-6.

Calistri D, Rengucci C, Molinari C, Ricci E, Cavargini E, Scarpi E, et al. Quantitative fluorescence determination of long-fragment DNA in stool as a marker for the early detection of colorectal cancer. *Cell Oncol* 2009;31(1):11-7.

Cappell MS. Safety and efficacy of colonoscopy after myocardial infarction: An analysis of 100 study patients and 100 control patients at two tertiary cardiac referral hospitals. *Gastrointest Endosc* 2004;60(6):901-9.

Castro G, Azrak MF, Seeff LC, Royalty J. Outpatient colonoscopy complications in the CDC's Colorectal Cancer Screening Demonstration Program: A prospective analysis. *Cancer* 2013;119(SUPPL. 15):2849-54.

Chang E, Park DI, Kim YJ, Kim BK, Park JH, Kim HJ, et al. Detection of colorectal neoplasm using promoter methylation of ITGA4, SFRP2, and p16 in stool samples: A preliminary report in Korean patients. *Hepatology* 2010;57(101):720-7.

Cheung HY, Chung CC, Kwok SY, Tsang WW, Li MK. Improvement in colonoscopy performance with adjunctive magnetic endoscope imaging: a randomized controlled trial. *Endoscopy* 2006;38(3):214-7.

Cobb WS, Heniford BT, Sigmon LB, Hasan R, Simms C, Kercher KW, et al. Colonoscopic perforations: incidence, management, and outcomes. *Am Surg* 2004;70(9):750-8.

Connolly DJA, Traill ZC, Reid HS, Copley SJ, Nolan DJ. The double contrast barium enema: a retrospective single centre audit of the detection of colorectal carcinomas. *Clin Radiol* 2002;57(1):29-32.

Courtney ED, Chong D, Tighe R, Easterbrook JR, Stebbings WSL, Herson J. Screen-detected colorectal cancers show improved cancer-specific survival when compared with cancers diagnosed via the 2-week suspected colorectal cancer referral guidelines. *Colorectal Dis* 2013;15(2):177-82.

Davies RJ, Collins CD, Vickery CJ, Eyre-Brook I, Welbourn R. Reduction in the proportion of patients with colorectal cancer presenting as an emergency following the introduction of fast-track flexible sigmoidoscopy: A three-year prospective observational study. *Colorectal Dis* 2004;6(4):265-7.

De CL, Rodriguez-Pineiro AM, Rodriguez-Berrocal FJ, Cordero OJ, Martinez-Ares D, Paez dIC. Serum CD26 is related to histopathological polyp traits and behaves as a marker for colorectal cancer and advanced adenomas. *BMC cancer* 2010;10.

Duff SE, Murray D, Rate AJ, Richards DM, Kumar NAM. Computed tomographic colonography (CTC) performance: one-year clinical follow-up. *Clin Radiol* 2006;61(11):932-6.

Durkalski VL, Palesch YY, Pineau BC, Vining DJ, Cotton PB. The virtual colonoscopy study: a large multicenter clinical trial designed to compare two diagnostic screening procedures. *Control Clin Trials* 2002;23(5):570-83.

East JE, Stavrindis M, Thomas-Gibson S, Guenther T, Tekkis PP, Saunders BP. A comparative study of standard vs. high definition colonoscopy for adenoma and hyperplastic polyp detection with optimized withdrawal technique. *Aliment Pharmacol Ther* 2008;28(6):768-76.

Edwards JK, Norris TE. Colonoscopy in rural communities: can family physicians perform the procedure with safe and efficacious results? *J Am Board Fam Pract* 2004;17(5):353-8.

Ee HC, Semmens JB, Hoffman NE, Perth Teaching Hospitals Endoscopy Group. Complete colonoscopy rarely misses cancer. *Gastrointest Endosc* 2002;55(2):167-71.

Erichsen R, Baron JA, Stoffel EM, Laurberg S, Sandler RS, Sorensen HT. Characteristics and survival of interval and sporadic colorectal cancer patients: a nationwide population-based cohort study. *Am J Gastroenterol* 2013;108(8):1332-40.

Fazio L, Cotterchio M, Manno M, McLaughlin J, Gallinger S. Association between colonic screening, subject characteristics, and stage of colorectal cancer. *Am J Gastroenterol* 2005;100(11):2531-9.

Ferrante JM, McCarthy EP, Gonzalez EC, Lee JH, Chen R, Love-Jackson K, et al. Primary care utilization and colorectal cancer outcomes among Medicare beneficiaries. *Arch Intern Med* 2011;171(19):1747-57.

Fisher DA, Jeffreys A, Grambow SC, Provenzale D. Mortality and follow-up colonoscopy after colorectal cancer. *Am J Gastroenterol* 2003;98(4):901-6.

Fletcher JG, Booya F, Summers RM, Roy D, Guendel L, Schmidt B, et al. Comparative performance of two polyp detection systems on CT colonography. *Am J Roentgenol* 2007;189(2):277-82.

Frenette CT, Strum WB. Relative rates of missed diagnosis for colonoscopy, barium enema, and flexible sigmoidoscopy in 379 patients with colorectal cancer. *J Gastrointest Cancer* 2007;38(2-4):148-53.

Gatto NM, Frucht H, Sundararajan V, Jacobson JS, Grann VR, Neugut AI. Risk of perforation after colonoscopy and sigmoidoscopy: a population-based study. *J Natl Cancer Inst* 2003;95(3):230-6.

Ginnerup Pedersen B, Christiansen TEM, Bjerregaard NC, Ljungmann K, Laurberg S. Colonoscopy and multidetector-array computed-tomographic colonography: detection rates and feasibility. *Endoscopy* 2003;35(9):736-42.

Gluecker T, Meuwly JY, Pescatore P, Schnyder P, DeLarive J, Jornod P, et al. Effect of investigator experience in CT colonography. *Eur Radiol* 2002;12(6):1405-9.

Goshen E, Davidson T, Aderka D, Zwas ST. PET/CT detects abdominal wall and port site metastases of colorectal carcinoma. *Br J Radiol* 2006;79(943):572-7.

Gralnek IM, Segol O, Suissa A, Siersema PD, Carr-Locke DL, Halpern Z, et al. A prospective cohort study evaluating a novel colonoscopy platform featuring full-spectrum endoscopy. *Endoscopy* 2013;45(9):697-702.

Greenberg PD, Bertario L, Gnauck R, Kronborg O, Hardcastle JD, Epstein MS, et al. A prospective multicenter evaluation of new fecal occult blood tests in patients undergoing colonoscopy. *Am J Gastroenterol* 2000;95(5):1331-8.

Gupta S, Saunders BP, Fraser C, Kennedy RH, Ignjatovic A, Sala S, et al. The first 3 years of national bowel cancer screening at a single UK tertiary centre. *Colorectal Dis* 2012;14(2):166-73.

Haji A, Ryan S, Bjarnason I, Donaldson N, Papagrigoriadis S. Colonoscopic high frequency mini-probe ultrasound is more accurate than conventional computed tomography in the local staging of colonic cancer. *Colorectal Dis* 2012;14(8):953-9.

Halligan S, Atkin WS. CT colonography for diagnosis of symptomatic colorectal cancer: The SIGGAR trials and their implication for service delivery. *Clin Radiol* 2013;68(7):643-5.

Hancer VS, Buyukdogan M, Turkmen I, Bassullu N, Altug T, Diz-Kucukkaya R, et al. Comparison of KRAS mutation tests in colorectal cancer patients. *Gent Test and Mol Biomarkers* 2011;15(11):831-4.

Hartmann D, Bassler B, Schilling D, Adamek HE, Jakobs R, Pfeifer B, et al. Colorectal polyps: detection with dark-lumen MR colonography versus conventional colonoscopy. *Radiology* 2006;238(1):143-9.

Haykir R, Karakose S, Karabacakoglu A, Kayacetin E, Sahin M. Detection of colonic masses with MR colonography. *Turk J Gastroenterol* 2006;17(3):191-7.

Haykir R, Karakose S, Karabacakoglu A, Sahin M, Kayacetin E. Three-dimensional MR and axial CT colonography versus conventional colonoscopy for detection of colon pathologies. *World J Gastroenterol* 2006;12(15):2345-50.

Hlavaty T, Huorka M, Koller T, Zita P, Kresanova E, Rychly B, et al. Colorectal cancer screening in patients with ulcerative and Crohn's colitis with use of colonoscopy, chromoendoscopy and confocal endomicroscopy. *Eur J Gastroenterol Hepatol* 2011;23(8):680-9.

Ho JM-W, Gruneir A, Fischer HD, Fu L, Lipscombe LL, Bell CM, et al. Serious events in older Ontario residents receiving bowel preparations for outpatient colonoscopy with various comorbidity profiles: a descriptive, population-based study. *Can J Gastroenterol* 2012;26(7):436-40.

Hoepffner N, Shastri YM, Hanisch E, Rosch W, Mossner J, Caspary WF, et al. Comparative evaluation of a new bedside faecal occult blood test in a prospective multicentre study. *Aliment Pharmacol Ther* 2006;23(1):145-54.

Hoppe H, Netzer P, Spreng A, Quattropani C, Mattich J, Dinkel HP. Prospective comparison of contrast enhanced CT colonography and conventional colonoscopy for detection of colorectal neoplasms in a single institutional study using second-look colonoscopy with discrepant results. *Am J Gastroenterol* 2004;99(10):1924-35.

Horiuchi A, Nakayama Y. Improved colorectal adenoma detection with a transparent retractable extension device. *Am J Gastroenterol* 2008;103(2):341-5.

Hurlstone DP, Cross SS, Adam I, Shorhouse AJ, Brown S, Sanders DS, et al. Endoscopic morphological anticipation of submucosal invasion in flat and depressed colorectal lesions: clinical implications and subtype analysis of the kudo type V pit pattern using high-magnification-chromoscopic colonoscopy. *Colorectal Dis* 2004;6(5):369-75.

Hurlstone DP, Cross SS, Drew K, Adam I, Shorhouse AJ, Brown S, et al. An evaluation of colorectal endoscopic mucosal resection using high-magnification chromoscopic colonoscopy: a prospective study of 1000 colonoscopies. *Endoscopy* 2004;36(6):491-8.

Iannaccone R, Laghi A, Catalano C, Brink JA, Mangiapane F, Trenna S, et al. Detection of colorectal lesions: lower-dose multi-detector row helical CT colonography compared with conventional colonoscopy. *Radiology* 2003;229(3):775-81.

Iannaccone R, Catalano C, Mangiapane F, Murakami T, Lamazza A, Fiori E, et al. Colorectal polyps: detection with low-dose multi-detector row helical CT colonography versus two sequential colonoscopies. *Radiology* 2005;237(3):927-37.

Iinuma G, Moriyama N, Satake M, Miyakawa K, Tateishi U, Uchiyama N, et al. Vascular virtual endoluminal visualization of invasive colorectal cancer on MDCT colonography. *Am J Roentgenol* 2005;184(4):1194-8.

Imai N, Takeda K, Kuzuya T, Utsunomiya S, Takahashi H, Kasuga H, et al. High incidence of colonic perforation during colonoscopy in hemodialysis patients with end-stage renal disease. *Clin Gastroenterol Hepatol* 2010;8(1):55-9.

Iqbal CW, Chun YS, Farley DR. Colonoscopic perforations: a retrospective review. *J Gastrointest Surg* 2005;9(9):1229-36.

Iussich G, Correale L, Senore C, Segnan N, Laghi A, Iafrate F, et al. CT colonography: preliminary assessment of a double-read paradigm that uses computer-aided detection as the first reader. *Radiology* 2013;268(3):743-51.

Jones RM, Mongin SJ, Lazovich D, Church TR, Yeazel MW. Validity of four self-reported colorectal cancer screening modalities in a general population: differences over time and by intervention assignment. *Cancer Epidemiol Biomarkers Prev* 2008;17(4):777-84.

Kahi CJ, Azzouz F, Juliar BE, Imperiale TF. Survival of elderly persons undergoing colonoscopy: implications for colorectal cancer screening and surveillance. *Gastrointest Endosc* 2007;66(3):544-50.

Kalimutho M, Del Vecchio BG, Cretella M, Mannisi E, Sileri P, Formosa A, et al. A simplified, non-invasive fecal-based DNA integrity assay and iFOBT for colorectal cancer detection. *Int J Colorectal Dis* 2011;26(5):583-92.

Kalra N, Suri S, Bhasin DK, Sinha SK, Saravanan N, Kour T, et al. Comparison of multidetector computed tomographic colonography and conventional colonoscopy for detection of colorectal polyps and cancer. *Indian J Gastroenterol* 2006;25(5):229-32.

Karajeh MA, Sanders DS, Hurlstone DP. Colonoscopy in elderly people is a safe procedure with a high diagnostic yield: A prospective comparative study of 2000 patients. *Endoscopy* 2006;38(3):226-30.

Kawamura YJ, Okada S, Sasaki J, Tajima N, Tanaka O, Konishi F. Diagnostic accuracy of CT colonography and optical colonoscopy evaluated using surgically resected specimens. *Abdom Imaging* 2010;35(5):584-8.

Kim DH, Pickhardt PJ, Taylor AJ, Leung WK, Winter TC, Hinshaw JL, et al. CT colonography versus colonoscopy for the detection of advanced neoplasia. *New Engl J Med* 2007;357(14):1403-12.

Ko CW, Riffle S, Shapiro JA, Saunders MD, Lee SD, Tung BY, et al. Incidence of minor complications and time lost from normal activities after screening or surveillance colonoscopy. *Gastrointest Endosc* 2007;65(4):648-56.

Koch M, Kienle P, Sauer P, Willeke F, Buhl K, Benner A, et al. Hematogenous tumor cell dissemination during colonoscopy for colorectal cancer. *Surg Endosc* 2004;18(4):587-91.

Kolber M, Szafran O, Suwal J, Diaz M. Outcomes of 1949 endoscopic procedures: performed by a Canadian rural family physician. *Can Fam Physician* 2009;55(2):170-5.

Korman LY, Overholt BF, Box T, Winker CK. Perforation during colonoscopy in endoscopic ambulatory surgical centers. *Gastrointest Endosc* 2003;58(4):554-7.

Kutzner N, Hoffmann I, Linke C, Thienel T, Grzegorzczak M, Urfer W, et al. Non-invasive detection of colorectal tumours by the combined application of molecular diagnosis and the faecal occult blood test. *Cancer Lett* 2005;229(1):33-41.

La Torre M, Velluti F, Giuliani G, Di Giulio E, Ziparo V, La Torre F. Promptness of diagnosis is the main prognostic factor after colonoscopic perforation. *Colorectal Dis* 2012;14(1):e23-e26.

Laghi A, Iannaccone R, Carbone I, Catalano C, Panebianco V, Di Giulio E, et al. Computed tomographic colonography (virtual colonoscopy): blinded prospective comparison with conventional colonoscopy for the detection of colorectal neoplasia. *Endoscopy* 2002;34(6):441-6.

Leaper M, Johnston MJ, Barclay M, Dobbs BR, Frizelle FA. Reasons for failure to diagnose colorectal carcinoma at colonoscopy. *Endoscopy* 2004;36(6):499-503.

Lee SH, Chung IK, Kim SJ, Kim JO, Ko BM, Kim WH, et al. Comparison of postpolypectomy bleeding between epinephrine and saline submucosal injection for large colon polyps by conventional polypectomy: a prospective randomized, multicenter study. *World J Gastroenterol* 2007;13(21):2973-7.

Leung WK, Lam WWM, Wu JCY, So NMC, Fung SSL, Chan FKL, et al. Magnetic resonance colonography in the detection of colonic neoplasm in high-risk and average-risk individuals. *Am J Gastroenterol* 2004;99(1):102-8.

Levi Z, Hazazi R, Rozen P, Vilkin A, Waked A, Niv Y. A quantitative immunochemical faecal occult blood test is more efficient for detecting significant colorectal neoplasia than a sensitive guaiac test. *Aliment Pharmacol Ther* 2006;23(9):1359-64.

Levi Z, Rozen P, Hazazi R, Vilkin A, Waked A, Maoz E, et al. Sensitivity, but not specificity, of a quantitative immunochemical fecal occult blood test for neoplasia is slightly increased by the use of low-dose aspirin, NSAIDs, and anticoagulants. *Am J Gastroenterol* 2009;104(4):933-8.

Li BS, Wang XY, Xu AG, Ma FL, Ma QY, Li Z, et al. High-resolution melting assay (HRMA) is a simple and sensitive stool-based DNA Test for the detection of mutations in colorectal neoplasms. *Clin Colorectal Cancer* 2012;11(4):280-90.

Lidgard GP, Domanico MJ, Bruinsma JJ, Light J, Gagrath ZD, Oldham-Haltom RL, et al. Clinical performance of an automated stool DNA assay for detection of colorectal neoplasia. *Clin Gastroenterol Hepatol* 2013;11(10):1313-8.

Loffeld RJLF. Are many colorectal cancers due to missed adenomas? *Eur J Intern Med* 2009;20(1):20-3.

Lohsiriwat V, Sujarittanakarn S, Akaraviputh T, Lertakyamanee N, Lohsiriwat D, Kachinthorn U. Colonoscopic perforation: A report from World Gastroenterology Organization endoscopy training center in Thailand. *World J Gastroenterol* 2008;14(43):6722-5.

Lohsiriwat V, Sujarittanakarn S, Akaraviputh T, Lertakyamanee N, Lohsiriwat D, Kachinthorn U. What are the risk factors of colonoscopic perforation? *BMC Gastroenterol* 2009;9:71.

Lorenzo-Zuniga V, Moreno de Vega V, Domenech E, Manosa M, Planas R, Boix J. Endoscopist experience as a risk factor for colonoscopic complications. *Colorectal Dis* 2010;12(10 Online):e273-e277.

Luo MY, Shan H, Yao LQ, Zhou KR, Liang WW. Postprocessing techniques of CT colonography in detection of colorectal carcinoma. *World J Gastroenterol* 2004;10(11):1574-7.

Mai CM, Wen CC, Wen SH, Hsu KF, Wu CC, Jao SW, et al. Iatrogenic colonic perforation by colonoscopy: a fatal complication for patients with a high anesthetic risk. *Int J Colorectal Dis* 2010;25(4):449-54.

Mandelli G, Radaelli F, Paggi S, Terreni N, Gola G, Gramegna M, et al. Anticoagulant or aspirin treatment does not affect the positive predictive value of an immunological fecal occult blood test in patients undergoing colorectal cancer screening: results from a nested in a cohort case-control study. *Eur J Gastroenterol Hepatol* 2011;23(4):323-6.

Matharoo GS, Goldfarb MA. Treatment and outcomes of iatrogenic colon perforations at a community teaching hospital. *Am Surg* 2012;78(9):975-8.

Mayinger B, Neumann F, Kastner C, Haider T, Schwab D. Hexaminolevulinate-induced fluorescence colonoscopy versus white light endoscopy for diagnosis of neoplastic lesions in the colon. *Endoscopy* 2010;42(1):28-33.

McConnell RA, Kim S, Ahmad NA, Falk GW, Forde KA, Ginsberg GG, et al. Poor discriminatory function for endoscopic skills on a computer-based simulator. *Gastrointest Endosc* 2012;76(5):993-1002.

Minordi LM, Vecchioli A, Mirk P, Filigrana E, Poloni G, Bonomo L. Multidetector CT in small-bowel neoplasms. *Radiol Med* 2007;112(7):1013-25.

Moore H, Dodd N. Computed tomographic colonography (CTC); Colorectal cancer diagnosis with CTC in an Auckland population. *J Med Imaging Radiat Oncol* 2013;57(5):572-5.

Morrin MM, Farrell RJ, Raptopoulos V, McGee JB, Bleday R, Kruskal JB. Role of virtual computed tomographic colonography in patients with colorectal cancers and obstructing colorectal lesions. *Dis Colon Rectum* 2000;43(3):303-11.

Munikrishnan V, Gillams AR, Lees WR, Vaizey CJ, Boulos PB. Prospective study comparing multislice CT colonography with colonoscopy in the detection of colorectal cancer and polyps. *Dis Colon Rectum* 2003;46(10):1384-90.

Navaneethan U, Parasa S, Venkatesh PGK, Shen B. Prevalence and risk factors for colonic perforation during colonoscopy in hospitalized end-stage renal disease patients on hemodialysis. *Int J Colorectal Dis* 2012;27(6):811-6.

Newman RJ, Nichols DB, Cummings DM. Outpatient colonoscopy by rural family physicians. *Ann Fam Med* 2005;3(2):122-5.

Nishida H, Urano S. Effectiveness of repeated screening using the fecal occult blood test and its impact on reducing false-negative cancer cases. *Eur J Cancer Prev* 2011;20(3):184-9.

Niv Y, Gershtansky Y, Kenett RS, Tal Y, Birkenfeld S. Complications in colonoscopy: analysis of 7-year physician-reported adverse events. *Eur J Gastroenterol Hepatol* 2011;23(6):492-8.

Niv Y, Bogolavski I, Ilani S, Avni I, Gal E, Vilkin A, et al. Impact of colonoscopy on quality of life. *Eur J Gastroenterol Hepatol* 2012;24(7):781-6.

Oort FA, Terhaar sive Droste JS, van der Hulst RWM, Van Heukelem HA, Loffeld RJLF, Wesdorp ICE, et al. Colonoscopy-controlled intra-individual comparisons to screen relevant neoplasia: Faecal immunochemical test vs. guaiac-based faecal occult blood test. *Aliment Pharmacol Ther* 2010;31(3):432-9.

Ou CH, Kuo FC, Hsu WH, Lu CY, Yu FJ, Kuo CH, et al. Comparison of the performance of guaiac-based and two immunochemical fecal occult blood tests for identifying advanced colorectal neoplasia in Taiwan. *J Dig Dis* 2013;14(9):474-83.

Parente F, Marino B, Ilardo A, Fracasso P, Zullo A, Hassan C, et al. A combination of faecal tests for the detection of colon cancer: a new strategy for an appropriate selection of referrals to colonoscopy? A prospective multicentre Italian study. *Eur J Gastroenterol Hepatol* 2012;24(10):1145-52.

Park SH, Ha HK, Kim MJ, Kim KW, Kim AY, Yang DH, et al. False-negative results at multi-detector row CT colonography: multivariate analysis of causes for missed lesions. *Radiology* 2005;235(2):495-502.

Pedersen JW, Gentry-Maharaj A, Fourkala EO, Dawney A, Burnell M, Zaikin A, et al. Early detection of cancer in the general population: a blinded case-control study of p53 autoantibodies in colorectal cancer. *Br J Cancer* 2013;108(1):107-14.

Pellise M, Fernandez-Esparrach G, Cardenas A, Sendino O, Ricart E, Vaquero E, et al. Impact of wide-angle, high-definition endoscopy in the diagnosis of colorectal neoplasia: a randomized controlled trial. *Gastroenterology* 2008;135(4):1062-8.

Peng J, He Y, Xu J, Sheng J, Cai S, Zhang Z. Detection of incidental colorectal tumours with 18F-labelled 2-fluoro-2-deoxyglucose positron emission tomography/computed tomography scans: results of a prospective study. *Colorectal Dis* 2011;13(11):e374-e378.

Rasmussen M, Kronborg O. Upper gastrointestinal cancer in a population-based screening program with fecal occult blood test for colorectal cancer. *Scand J Gastroenterol* 2002;37(1):95-8.

Regge D, Laudi C, Galatola G, Della Monica P, Bonelli L, Angelelli G, et al. Diagnostic accuracy of computed tomographic colonography for the detection of advanced neoplasia in individuals at increased risk of colorectal cancer. *JAMA : the journal of the American Medical Association* 2009;301(23):2453-61.

Regge D, Della Monica P, Galatola G, Laudi C, Zambon A, Correale L, et al. Efficacy of computer-aided detection as a second reader for 6-9-mm lesions at CT colonography: multicenter prospective trial. *Radiology* 2013;266(1):168-76.

Repici A, Hassan C, Vitetta E, Ferrara E, Manes G, Gullotti G, et al. Safety of cold polypectomy for <10mm polyps at colonoscopy: a prospective multicenter study. *Endoscopy* 2012;44(1):27-31.

Reuterskiold MH, Lasson A, Svensson E, Kilander A, Stotzer PO, Hellstrom M. Diagnostic performance of computed tomography colonography in symptomatic patients and in patients with increased risk for colorectal disease. *Acta Radiol* 2006;47(9):888-98.

Roach SC, Hulse PA, Moulding FJ, Wilson R, Carrington BM. Magnetic resonance imaging of anal cancer. *Clin Radiol* 2005;60(10):1111-9.

Roberts-Thomson IC, Tucker GR, Hewett PJ, Cheung P, Sebben RA, Khoo EEW, et al. Single-center study comparing computed tomography colonography with conventional colonoscopy. *World J Gastroenterol* 2008;14(3):469-73.

Robinson C, Halligan S, Iinuma G, Topping W, Punwani S, Honeyfield L, et al. CT colonography: computer-assisted detection of colorectal cancer. *Br J Radiol* 2011;84(1001):435-40.

Rozen P, Knaani J, Samuel Z. Comparative screening with a sensitive guaiac and specific immunochemical occult blood test in an endoscopic study. *Cancer* 2000;89(1):46-52.

Rozen P, Comaneshter D, Levi Z, Hazazi R, Vilkin A, Maoz E, et al. Cumulative evaluation of a quantitative immunochemical fecal occult blood test to determine its optimal clinical use. *Cancer* 2010;116(9):2115-25.

Rozen P, Liphshitz I, Barchana M. Follow-up of patients undergoing both semiquantitated immunochemical fecal occult blood and colonoscopy examinations. *Eur J Cancer Prev* 2012;21(3):247-53.

Ruffin MT, Normolle DP, Eveleigh MJ, Baron JA, Bresalier RS, Marcon NE, et al. Rectal mucosal quantitative galactose oxidase-Schiff reaction as an early detection biomarker for colorectal cancer: Comparison to fecal occult stool blood test. *Cancer Biomarkers* 2011;8(2):109-12.

Sabharwal R, Vladica P, Chou R, Law WP. Helical CT in the diagnosis of acute lower gastrointestinal haemorrhage. *Eur J Radiol* 2006;58(2):273-9.

Sacher-Huvelin S, Coron E, Gaudric M, Planche L, Benamouzig R, Maunoury V, et al. Colon capsule endoscopy vs. colonoscopy in patients at average or increased risk of colorectal cancer. *Aliment Pharmacol Ther* 2010;32(9):1145-53.

Sakamoto T, Saito Y, Nakajima T, Matsuda T. Comparison of magnifying chromoendoscopy and narrow-band imaging in estimation of early colorectal cancer invasion depth: a pilot study. *Digestive endoscopy : official journal of the Japan Gastroenterological Endoscopy Society* 2011;23(2):118-23.

Selinger CP, Mullender J, Choudhury S, Jones PE, Sukumar S, Ramesh J. Flexible sigmoidoscopy does not significantly increase polyp and cancer detection yield when used to supplement CT colonography. *Digestion* 2012;85(1):55-60.

Sewitch MJ, Jiang M, Joseph L, Barkun AN, Bitton A. Rate of serious complications of colonoscopy in Quebec. *Can J Gastroenterol* 2012;26(9):611-3.

Shastri YM, Loitsch S, Hoepffner N, Povse N, Hanisch E, Rosch W, et al. Comparison of an established simple office-based immunological FOBT with fecal tumor pyruvate kinase type M2 (M2-PK) for colorectal cancer screening: prospective multicenter study. *Am J Gastroenterol* 2008;103(6):1496-504.

Shia J, Klimstra DS, Nafa K, Offit K, Guillem JG, Markowitz AJ, et al. Value of immunohistochemical detection of DNA mismatch repair proteins in predicting germline mutation in hereditary colorectal neoplasms. *Am J Surg Pathol* 2005;29(1):96-104.

Siddique I, Mohan K, Hasan F, Memon A, Patty I, Al-Nakib B. Appropriateness of indication and diagnostic yield of colonoscopy: first report based on the 2000 guidelines of the American Society for Gastrointestinal Endoscopy. *World J Gastroenterol* 2005;11(44):7007-13.

Singhal S, Verma A, Anand K. Colonoscopy for colorectal cancer screening above age 75: outcomes in symptomatic african american and Hispanic adults. *J Gastrointest Cancer* 2011;42(4):212-6.

Smiljanic S, Gill S. Patterns of diagnosis for colorectal cancer: Screening detected vs. symptomatic presentation. *Dis Colon Rectum* 2008;51(5):573-7.

Sofic A, Beslic S, Kocijancic I, Sehovic N. CT colonography in detection of colorectal carcinoma. *Radiol Oncol* 2010;44(1):19-23.

Sosna J, Blachar A, Amitai M, Barmeir E, Peled N, Goldberg SN, et al. Colonic perforation at CT colonography: assessment of risk in a multicenter large cohort. *Radiology* 2006;239(2):457-63.

Spada C, Hassan C, Munoz-Navas M, Neuhaus H, Deviere J, Fockens P, et al. Second-generation colon capsule endoscopy compared with colonoscopy. *Gastrointest Endosc* 2011;74(3):581-9.

Spinzi G, Belloni G, Martegani A, Sangiovanni A, Del Favero C, Minoli G. Computed tomographic colonography and conventional colonoscopy for colon diseases: a prospective, blinded study. *Am J Gastroenterol* 2001;96(2):394-400.

Stermer E, Lavy A, Rainis T, Goldstein O, Keren D, Zeina AR. Incidental colorectal computed tomography abnormalities: would you send every patient for a colonoscopy? *Can J Gastroenterol* 2008;22(9):758-60.

Strul H, Arber N. Fecal occult blood test for colorectal cancer screening. *Ann Oncol* 2002;13(1):51-6.

Suissa A, Bentur OS, Lachter J, Yassin K, Chermesh I, Gralnek I, et al. Outcome and complications of colonoscopy: A prospective multicenter study in Northern Israel. *Diagn Ther Endosc* 2012.

Terhaar sive Droste JS, van Turenhout ST, Oort FA, van der Hulst RWM, Steeman VA, Coblijn U, et al. Faecal immunochemical test accuracy in patients referred for surveillance colonoscopy: a multi-centre cohort study. *BMC Gastroenterol* 2012;12:94.

Thakur CS, Sood RG, Makhaik S, Sharma B, Chauhan A, Thakur S. Evaluation of the efficacy of Virtual Colonoscopy in assessment of colorectal lesions using conventional Optical Colonoscopy as the gold standard. *Egypt J Radiol Nucl Med* 2013;44(3):433-40.

Thomas S, Atchley J, Higginson A. Audit of the introduction of CT colonography for detection of colorectal carcinoma in a non-academic environment and its implications for the national bowel cancer screening programme. *Clin Radiol* 2009;64(2):142-7.

Vajdic CM, Anderson JS, Hillman RJ, Medley G, Grulich AE. Blind sampling is superior to anoscope guided sampling for screening for anal intraepithelial neoplasia. *Sex Transm Infect* 2005;81(5):415-8.

van der Sluis FJ, Loffeld RJ, Engel AF. Outcome of surgery for colonoscopic perforation. *Colorectal Dis* 2012;14(4):e187-e190.

van Gelder RE, Nio CY, Florie J, Bartelsman JF, Snel P, De Jager SW, et al. Computed tomographic colonography compared with colonoscopy in patients at increased risk for colorectal cancer. *Gastroenterology* 2004;127(1):41-8.

van Rossum LGM, van Rijn AF, Van Munster IP, Jansen JBMJ, Fockens P, Laheij RJF, et al. Earlier stages of colorectal cancer detected with immunochemical faecal occult blood tests. *Neth J Med* 2009;67(5):182-6.

Vella M, MacKenzie S, Young IE, Molloy RG, O'Dwyer PJ. Impact of video colonoscopy on stage and outcome of patients with symptomatic colorectal cancer. *Surg Endosc Interv Tech* 2004;18(8):1268-71.

Vellacott KD, Clarke D, Beech C, Joshi H. Flexible sigmoidoscopy in general practice. *Colorectal Dis* 2002;4(2):123-6.

Vironen J, Kellokumpu S, Andersson LC, Kellokumpu I. Comparison of a peanut agglutinin test and an immunochemical faecal occult blood test in detecting colorectal neoplasia in symptomatic patients. *Scand J Clin Lab Invest* 2004;64(2):140-5.

Wang D-R, Tang D. Hypermethylated SFRP2 gene in fecal DNA is a high potential biomarker for colorectal cancer noninvasive screening. *World J Gastroenterol* 2008;14(4):524-31.

Ward RL, Turner J, Williams R, Pekarsky B, Packham D, Velickovic M, et al. Routine testing for mismatch repair deficiency in sporadic colorectal cancer is justified. *J Pathol* 2005;207(4):377-84.

Ward SM, Mobsby CJ, Wilkinson S. Capsule endoscopy: A surgical perspective. *ANZ J Surg* 2008;78(1-2):28-33.

Weissfeld JL, Schoen RE, Pinsky PF, Bresalier RS, Church T, Yurgalevitch S, et al. Flexible sigmoidoscopy in the PLCO cancer screening trial: results from the baseline screening examination of a randomized trial. *J Natl Cancer Inst* 2005;97(13):989-97.

White TJ, Avery GR, Kennan N, Syed AM, Hartley JE, Monson JRT. Virtual colonoscopy vs conventional colonoscopy in patients at high risk of colorectal cancer--a prospective trial of 150 patients. *Colorectal Dis* 2009;11(2):138-45.

Wilson S, Wakelam MJO, Hobbs RFD, Ryan AV, Dunn JA, Redman VD, et al. Evaluation of the accuracy of serum MMP-9 as a test for colorectal cancer in a primary care population. *BMC cancer* 2006;6:258.

Winkleman BJ, Matthews DE, Wiebke EA. Colorectal cancer screening at a Veterans Affairs hospital. *Am J Surg* 2003;186(5):468-71.

Witt DM, Delate T, McCool KH, Dowd MB, Clark NP, Crowther MA, et al. Incidence and predictors of bleeding or thrombosis after polypectomy in patients receiving and not receiving anticoagulation therapy. *J Thromb Haemost* 2009;7(12):1982-9.

Wong BCY, Wong WM, Cheung KL, Tong TSM, Rozen P, Young GP, et al. A sensitive guaiac faecal occult blood test is less useful than an immunochemical test for colorectal cancer screening in a Chinese population. *Aliment Pharmacol Ther* 2003;18(9):941-6.

Wong CKW, Fedorak RN, Prosser CI, Stewart ME, Van Zanten SV, Sadowski DC. The sensitivity and specificity of guaiac and immunochemical fecal occult blood tests for the detection of advanced colonic adenomas and cancer. *Int J Colorectal Dis* 2012;27(12):1657-64.

Wong WM, Lam SK, Cheung KL, Tong TSM, Rozen P, Young GP, et al. Evaluation of an automated immunochemical fecal occult blood test for colorectal neoplasia detection in a Chinese population. *Cancer* 2003;97(10):2420-4.

Yajima S, Ishii M, Matsushita H, Aoyagi K, Yoshimatsu K, Kaneko H, et al. Expression profiling of fecal colonocytes for RNA-based screening of colorectal cancer. *Int J Oncol* 2007;31(5):1029-37.

Yang DH, Byeon JS, Lee KH, Yoon SM, Kim KJ, Ye BD, et al. Is endoscopic closure with clips effective for both diagnostic and therapeutic colonoscopy-associated bowel perforation? *Surg Endosc* 2010;24(5):1177-85.

Yao MD, von Rosenvinge EC, Groden C, Mannon PJ. Multiple endoscopic biopsies in research subjects: safety results from a National Institutes of Health series. *Gastrointest Endosc* 2009;69(4):906-10.

Yoong KKY, Heymann T. It is not worthwhile to perform ileoscopy on all patients. *Surg Endosc* 2006;20(5):809-11.

You Y-T, Chang CCR, Wang J-Y, Ng K-K, Chen J-S, Tang R, et al. Evaluation of contrast-enhanced computed tomographic colonography in detection of local recurrent colorectal cancer. *World J Gastroenterol* 2006;12(1):123-6.

Yousaf KR, Atiq S, Nisar MS. Detection and appearances of malignant colorectal masses through transabdominal sonography. *Pak J Med Health Sci* 2012;6(1):205-7.

Yucel C, Lev-Toaff AS, Moussa N, Durrani H. CT colonography for incomplete or contraindicated optical colonoscopy in older patients. *Am J Roentgenol* 2008;190(1):145-50.

Zaher T, Bahgat M, Ibrahim A, Ahmady M, Esmat S, Gouda H, et al. Colorectal polyps in Sharykia, Egypt: Clinico-pathological study of our experience with endoscopy. *J Med Sci (Pakistan)* 2008;8(2):196-200.

Excluded No Comparison Group of Interest

Adler A, Roll S, Marowski B, Drossel R, Rehs H-U, Willich SN, et al. Appropriateness of colonoscopy in the era of colorectal cancer screening: A prospective, multicenter study in a private-practice setting (Berlin Colonoscopy Project 1, BECOP 1). *Dis Colon Rectum* 2007;50(10):1628-38.

Adler A, Aschenbeck J, Yenerim T, Mayr M, Ainalai A, Drossel R, et al. Narrow-band versus white-light high definition television endoscopic imaging for screening colonoscopy: a prospective randomized trial. *Gastroenterology* 2009;136(2):410-715.

Adler A, Ainalai A, Aschenbeck J, Drossel R, Mayr M, Scheel M, et al. Latest generation, wide-angle, high-definition colonoscopes increase adenoma detection rate. *Clin Gastroenterol Hepatol* 2012;10(2):155-9.

Ahmed FE, Vos P. Molecular markers for human colon cancer in stool and blood identified by RT-PCR. *Anticancer Res* 2004;24(6):4127-34.

Akedo I, Ishikawa H, Ioka T, Kaji I, Narahara H, Ishiguro S, et al. Evaluation of epithelial cell proliferation rate in normal-appearing colonic mucosa as a high-risk marker for colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 2001;10(9):925-30.

Arumugam PJ, Rao GN, West J, Foster ME, Haray PN. The impact of open access flexible sigmoidoscopy: a comparison of two services. *J R Coll Surg Edinb* 2000;45(6):366-8.

Bair D, Pham J, Seaton MB, Arya N, Pryce M, Seaton TL. The quality of screening colonoscopies in an office-based endoscopy clinic. *Can J Gastroenterol* 2009;23(1):41-7.

Bardot L, Gobert JG, Kapel N. [Faecal occult blood testing: comparison of a latex agglutination test (Hemolex) and an immunoturbidimetric test (QuikRead FOB)]. *Recherche de sang occulte dans les selles: analyse comparative d'un test d'agglutination de particules de latex sensibilisees (Hemolex) et d'un test immunoturbidimetrique (QuikRead FOB)* 2004;62(3):339-43.

Baudet JS, Diaz-Bethencourt D, Aviles J, Aguirre-Jaime A. Minor adverse events of colonoscopy on ambulatory patients: the impact of moderate sedation. *Eur J Gastroenterol Hepatol* 2009;21(6):656-61.

Benhaberou-Brun D. [Colorectal cancer screening. Where are we?]. *Perspect Infirm* 2013;10(3):52-5.

Berhane C, Denning D. Incidental finding of colorectal cancer in screening colonoscopy and its cost effectiveness. *Am Surg* 2009;75(8):699-4.

Betes M, Munoz-Navas MA, Duque JM, Angos R, Macias E, Subtil JC, et al. Use of Colonoscopy as a Primary Screening Test for Colorectal Cancer in Average Risk People. *Am J Gastroenterol* 2003;98(12):2648-54.

Bezabeh T, Somorjai R, Dolenko B, Bryskina N, Levin B, Bernstein CN, et al. Detecting colorectal cancer by 1H magnetic resonance spectroscopy of fecal extracts. *NMR Biomed* 2009;22(6):593-600.

Bini EJ, Reinhold J-P, Weinshel EH, Generoso R, Salman L, Dahr G, et al. Prospective evaluation of the use and outcome of admission stool guaiac testing: The digital rectal examination on admission to the medical service (DREAMS) study. *J Clin Gastroenterol* 2006;40(9):821-7.

Bittler RD. Splenic injury due to colonoscopy: nursing considerations. *Gastroenterol Nurs* 2011;34(5):357-64.

Bleiberg H, Autier P, Huet F, Schrauwen AM, Staquet E, Delaunoit T, et al. Colorectal cancer (CRC) screening using sigmoidoscopy followed by colonoscopy: A feasibility and efficacy study on a cancer institute based population. *Ann Oncol* 2006;17(8):1328-32.

Blom J, Liden A, Jeppsson B, Holmberg L, Pahlman L. Compliance and findings in a Swedish population screened for colorectal cancer with sigmoidoscopy. *Eur J Surg Oncol* 2002;28(8):827-31.

Boellaard TN, van der Paardt MP, Hollmann MW, Eberl S, Peringa J, Schouten LJ, et al. A multi-centre randomised double-blind placebo-controlled trial to evaluate the value of a single bolus intravenous alfentanil in CT colonography. *BMC Gastroenterol* 2013;13(1).

Bonelli L, Sciallero S, Senore C, Zappa M, Aste H, Andreoni B, et al. History of negative colorectal endoscopy and risk of rectosigmoid neoplasms at screening flexible sigmoidoscopy. *Int J Colorectal Dis* 2006;21(2):105-13.

Brasso K, Ladelund S, Frederiksen BL, Jorgensen T. Psychological distress following fecal occult blood test in colorectal cancer screening--a population-based study. *Scand J Gastroenterol* 2010;45(10):1211-6.

Breen N, Wagener DK, Brown ML, Davis WW, Ballard-Barbash R. Progress in cancer screening over a decade: Results of cancer screening from the 1987, 1992, and 1998 National Health Interview Surveys. *J Natl Cancer Inst* 2001;93(22):1704-13.

Brenner H, Arndt V, Sturmer T, Stegmaier C, Ziegler H, Dhom G. Long-lasting reduction of risk of colorectal cancer following screening endoscopy. *Br J Cancer* 2001;85(7):972-6.

Brenner H, Arndt V, Stegmaier C, Ziegler H, Sturmer T. Reduction of clinically manifest colorectal cancer by endoscopic screening: Empirical evaluation and comparison of screening at various ages. *Eur J Cancer Prev* 2005;14(3):231-7.

Brenner H, Chang-Claude J, Seiler CM, Hoffmeister M. Interval cancers after negative colonoscopy: population-based case-control study. *Gut* 2012;61(11):1576-82.

Brossek D, Oosthuizen J, Pinchbeck M. Initial experience with a general population colorectal cancer screening clinic. *Am J Surg* 2006;191(5):669-72.

Cardin F, Andreotti A, Martella B, Terranova C, Militello C. Current practice in colonoscopy in the elderly. *Aging Clin Exp Res* 2012;24(SUPPL.3):9-13.

Chen JG, Cai J, Wu HL, Xu H, Zhang YX, Chen C, et al. Colorectal cancer screening: comparison of transferrin and immuno fecal occult blood test. *World J Gastroenterol* 2012;18(21):2682-8.

Chen LS, Yen AM-F, Chiu SY-H, Liao CS, Chen HH. Baseline faecal occult blood concentration as a predictor of incident colorectal neoplasia: longitudinal follow-up of a Taiwanese population-based colorectal cancer screening cohort. *Lancet Oncol* 2011;12(6):551-8.

Chew MH, Suzanah N, Ho KS, Lim JF, Ooi BS, Tang CL, et al. Colorectal cancer mass screening event utilising quantitative faecal occult blood test. *Singapore Med J* 2009;50(4):348-53.

Chiu HM, Lin JT, Chen THH, Lee YC, Chiu YH, Liang JT, et al. Elevation of C-reactive protein level is associated with synchronous and advanced colorectal neoplasm in men. *Am J Gastroenterol* 2008;103(9):2317-25.

Chung SJ, Kim YS, Yang SY, Song JH, Kim D, Park MJ, et al. Five-year risk for advanced colorectal neoplasia after initial colonoscopy according to the baseline risk stratification: a prospective study in 2452 asymptomatic Koreans. *Gut* 2011;60(11):1537-43.

Ciatto S, Martinelli F, Castiglione G, Mantellini P, Rubeca T, Grazzini G, et al. Association of FOBt-assessed faecal Hb content with colonic lesions detected in the Florence screening programme. *Br J Cancer* 2007;96(2):218-21.

Coe SG, Crook JE, Diehl NN, Wallace MB. An endoscopic quality improvement program improves detection of colorectal adenomas. *Am J Gastroenterol* 2013;108(2):219-27.

Cole SR, Tucker GR, Osborne JM, Byrne SE, Bampton PA, Fraser RJL, et al. Shift to earlier stage at diagnosis as a consequence of the National Bowel Cancer Screening Program. *Med J Aust* 2013;198(6):327-30.

Copel L, Sosna J, Kruskal JB, Raptopoulos V, Farrell RJ, Morrin MM. CT colonography in 546 patients with incomplete colonoscopy. *Radiology* 2007;244(2):471-8.

Corbett M, Chambers SL, Shadbolt B, Hillman LC, Taupin D. Colonoscopy screening for colorectal cancer: the outcomes of two recruitment methods. *Med J Aust* 2004;181(8):423-7.

Costantini AS, Martini A, Puliti D, Ciatto S, Castiglione G, Grazzini G, et al. Colorectal cancer mortality in two areas of Tuscany with different screening exposures. *J Natl Cancer Inst* 2008;100(24):1818-21.

Courtney RJ, Paul CL, Carey ML, Sanson-Fisher RW, Macrae FA, D'Este C, et al. A population-based cross-sectional study of colorectal cancer screening practices of first-degree relatives of colorectal cancer patients. *BMC cancer* 2013;13:13.

Dae HK, Seok YL, Kui SC, Ho JL, Su CP, Kim J, et al. The usefulness of colonoscopy as a screening test for detecting colorectal polyps. *Hepatogastroenterology* 2007;54(80):2240-2.

De Palma GD, Rega M, Masone S, Persico M, Siciliano S, Addeo P, et al. Conventional colonoscopy and magnified chromoendoscopy for the endoscopic histological prediction of diminutive colorectal polyps: a single operator study. *World J Gastroenterol* 2006;12(15):2402-5.

Denis B, Sauleau EA, Gendre I, Piette C, Bretagne JF, Perrin P. Measurement of adenoma detection and discrimination during colonoscopy in routine practice: an exploratory study. *Gastrointest Endosc* 2011;74(6):1325-36.

Denters MJ, Deutekom M, Bossuyt PM, Fockens P, Dekker E. Patient burden of colonoscopy after positive fecal immunochemical testing for colorectal cancer screening. *Endoscopy* 2013;45(5):342-9.

Denters MJ, Deutekom M, Fockens P, Bossuyt PM, Dekker E. Implementation of population screening for colorectal cancer by repeated fecal occult blood test in the Netherlands. *BMC Gastroenterol* 2009;9:28.

Dowling DJ, St.John DJB, Macrae FA, Hopper JL. Yield from colonoscopic screening in people with a strong family history of common colorectal cancer. *J Gastroenterol Hepatol* 2000;15(8):939-44.

Duncan JE, Sweeney WB, Trudel JL, Madoff RD, Mellgren AF. Colonoscopy in the elderly: low risk, low yield in asymptomatic patients. *Dis Colon Rectum* 2006;49(5):646-51.

Faccioli N, Foti G, Barillari M, Zaccarella A, Camera L, Biasiutti C, et al. A simplified approach to virtual colonoscopy using different intestinal preparations: preliminary experience with regard to quality, accuracy and patient acceptability. *Radiol Med* 2011;116(5):749-58.

Farrar WD, Sawhney MS, Nelson DB, Lederle FA, Bond JH. Colorectal Cancers Found After a Complete Colonoscopy. *Clin Gastroenterol Hepatol* 2006;4(10):1259-64.

Fletcher JG, Silva AC, Fidler JL, Cernigliaro JG, Manduca A, Limburg PJ, et al. Noncathartic CT colonography: Image quality assessment and performance and in a screening cohort. *AJR Am J Roentgenol* 2013;201(4):787-94.

Fludger S, Turner AM, Harvey RF, Haslam N. Controlled prospective study of faecal occult blood screening for colorectal cancer in Bury, black pudding capital of the world. *BMJ* 2002;325(7378):1444-5.

Francois F, Park J, Bini EJ. Colon pathology detected after a positive screening flexible sigmoidoscopy: A prospective study in an ethnically diverse cohort. *Am J Gastroenterol* 2006;101(4):823-30.

Fu W-P, Kam M-H, Ling W-M, Ong S-F, Suzannah N, Eu K-W. Screening for colorectal cancer using a quantitative immunochemical faecal occult blood test: A feasibility study in an Asian population. *Tech Coloproctol* 2009;13(3):225-30.

Gellad ZF, Stechuchak KM, Fisher DA, Olsen MK, McDuffie JR, Ostbye T, et al. Longitudinal adherence to fecal occult blood testing impacts colorectal cancer screening quality. *Am J Gastroenterol* 2011;106(6):1125-34.

Gilbert JM, Vaizey CJ, Cassell PG, Holden J. Feasibility study of colonoscopy as the primary screening investigation in relatives of patients with colorectal cancer. *Ann R Coll Surg Engl* 2001;83(6):415-9.

Gimeno-Garcia AZ, Quintero E, Nicolas-Perez D, Hernandez-Guerra M, Parra-Blanco A, Jimenez-Sosa A. Screening for familial colorectal cancer with a sensitive immunochemical fecal occult blood test: A pilot study. *Eur J Gastroenterol Hepatol* 2009;21(9):1062-7.

Goehde SC, Hunold P, Vogt FM, Ajaj W, Goyen M, Herborn CU, et al. Full-body cardiovascular and tumor MRI for early detection of disease: Feasibility and initial experience in 298 subjects. *Am J Roentgenol* 2005;184(2):598-611.

Goetzl MA, Outman JE, Griebing TL, Holzbeierlein JM, Weigel JW, Thrasher JB. Digital fecal occult blood testing in the ambulatory urology clinic. *J Am Coll Surg* 2008;206(1):144-7.

Gonvers JJ, Harris JK, Wietlisbach V, Burnand B, Vader JP, Froehlich F, et al. A European view of diagnostic yield and appropriateness of colonoscopy. *Hepatogastroenterology* 2007;54(75):729-35.

Gorard DA, McIntyre AS. Completion rate to caecum as a quality measure of colonoscopy in a district general hospital. *Colorectal Dis* 2004;6(4):243-9.

Goulard H, Boussac-Zarebska M, Ancelle-Park R, Bloch J. French colorectal cancer screening pilot programme: Results of the first round. *J Med Screen* 2008;15(3):143-8.

Grasset D, Morfoisse JJ, Seigneuric C. [Conditions of practice and results of colonoscopy in non-university hospitals. Results of a cross sectional, multicenter ANGH study (1)]. Conditions de realisation et resultats des coloscopies realisees dans les hopitaux non universitaires Resultats d'une enquete transversale multicentrique de l'ANGH (1) 2000;24(3):273-8.

Grassini M, Verna C, Niola P, Navino M, Battaglia E, Bassotti G. Appropriateness of colonoscopy: diagnostic yield and safety in guidelines. *World J Gastroenterol* 2007;13(12):1816-9.

Greenspan M, Rajan KB, Baig A, Beck T, Mobarhan S, Melson J. Advanced adenoma detection rate is independent of nonadvanced adenoma detection rate. *Am J Gastroenterol* 2013;108(8):1286-92.

Gupta AK, Melton LJ, Petersen GM, Timmons LJ, Vege SS, Harmsen WS, et al. Changing trends in the incidence, stage, survival, and screen-detection of colorectal cancer: a population-based study. *Clin Gastroenterol Hepatol* 2005;3(2):150-8.

Gupta S, Tong L, Anderson P, Rose B, Carter E, Koch M, et al. Measurement of colorectal cancer test use with medical claims data in a safety-net health system. *Am J Med Sci* 2013;345(2):99-103.

Habermann JK, Roblick UJ, Luke BT, Prieto DA, Finlay WJJ, Podust VN, et al. Increased Serum Levels of Complement C3a Anaphylatoxin Indicate the Presence of Colorectal Tumors. *Gastroenterology* 2006;131(4):1020-9.

Half EE, Mlynarsky L, Naftali T, Benjaminov F, Konikoff FM. False negative fecal occult blood test may be associated with increased mortality from colorectal cancer. *Dig Dis Sci* 2013;58(9):2639-45.

Harewood GC, Lawlor GO, Larson MV. Incident rates of colonic neoplasia in older patients: When should we stop screening? *J Gastroenterol Hepatol* 2006;21(6):1021-5.

Hart AR, Glover N, Howick-Baker J, Mayberry JF. An industry based approach to colorectal cancer screening in an asymptomatic population. *Postgrad Med J* 2003;79(937):646-9.

Hassan C, Pooler BD, Kim DH, Rinaldi A, Repici A, Pickhardt PJ. Computed tomographic colonography for colorectal cancer screening: Risk factors for the detection of advanced neoplasia. *Cancer* 2013;119(14):2549-54.

Helm JF, Russo MW, Biddle AK, Simpson KN, Ransohoff DF, Sandler RS. Effectiveness and economic impact of screening for colorectal cancer by mass fecal occult blood testing. *Am J Gastroenterol* 2000;95(11):3250-8.

Heuss LT, Sughanda SP, Degen LP. Endoscopy teams' judgment of discomfort among patients undergoing colonoscopy: "How bad was it really?". *Swiss Med Wkly* 2012;142:w13726.

Hewett DG, Kaltenbach T, Sano Y, Tanaka S, Saunders BP, Ponchon T, et al. Validation of a simple classification system for endoscopic diagnosis of small colorectal polyps using narrow-band imaging. *Gastroenterology* 2012;143(3):599-607.

Hill J, Walsh S, Evans DG. Screening of patients at high risk of colorectal cancer. *Colorectal Dis* 2001;3(5):308-11.

Hirakawa T, Kato J, Okumura Y, Hori K, Takahashi S, Suzuki H, et al. Detectability of colorectal neoplasia with fluorine-18-2-fluoro-2-deoxy-D-glucose positron emission tomography and computed tomography (FDG-PET/CT). *J Gastroenterol* 2012;47(2):127-35.

Hock D, Ouhadi R, Materne R, Aouchria A-S, Mancini I, Broussaud T, et al. Virtual dissection CT colonography: Evaluation of learning curves and reading times with and without computer-aided detection. *Radiology* 2008;248(3):860-8.

Hoff G, Moritz V, Bretthauer M, Aabakken L, Berset IP, Glomsaker T, et al. Incontinence after colonoscopy--an unrecognized and preventable problem. A cross-sectional study from the Gastronet quality assurance program. *Endoscopy* 2012;44(4):349-53.

Hoffman A, Sar F, Goetz M, Tresch A, Mudter J, Biesterfeld S, et al. High definition colonoscopy combined with i-Scan is superior in the detection of colorectal neoplasias compared with standard video colonoscopy: a prospective randomized controlled trial. *Endoscopy* 2010;42(10):827-33.

Hubbard RA, Johnson E, Hsia R, Rutter CM. The cumulative risk of false-positive fecal occult blood test after 10 years of colorectal cancer screening. *Cancer Epidemiol Biomarkers Prev* 2013;22(9):1612-9.

Ikematsu H, Saito Y, Yamano H. Comparative evaluation of endoscopic factors from conventional colonoscopy and narrow-band imaging of colorectal lesions. *Digestive endoscopy : official journal of the Japan Gastroenterological Endoscopy Society* 2011;23 Suppl 1:95-100.

Ikematsu H, Saito Y, Tanaka S, Uraoka T, Sano Y, Horimatsu T, et al. The impact of narrow band imaging for colon polyp detection: a multicenter randomized controlled trial by tandem colonoscopy. *J Gastroenterol* 2012;47(10):1099-107.

Imler TD, Morea J, Kahi C, Imperiale TF. Natural language processing accurately categorizes findings from colonoscopy and pathology reports. *Clin Gastroenterol Hepatol* 2013;11(6):689-94.

Imperiale TF, Glowinski EA, Lin-Cooper C, Larkin GN, Rogge JD, Ransohoff DF. Five-year risk of colorectal neoplasia after negative screening colonoscopy. *N Engl J Med* 2008;359(12):1218-24.

Jin XF, Chai TH, Shi JW, Yang XC, Sun QY. Meta-analysis for evaluating the accuracy of endoscopy with narrow band imaging in detecting colorectal adenomas. *J Gastroenterol Hepatol* 2012;27(5):882-7.

Joukar F, Majd SK, Fani A, Nazari N, Mansour-Ghanaei F. Colonoscopy outcome in North of Iran (Guilan): 2006-2009. *Int J Clin Exp Med* 2012;5(4):321-5.

Kaltenbach T, McGill SK, Kalidindi V, Friedland S, Soetikno R. Proficiency in the diagnosis of nonpolypoid colorectal neoplasm yields high adenoma detection rates. *Dig Dis Sci* 2012;57(3):764-70.

Keren D, Rainis T, Goldstein O, Stermer E, Lavy A. Significant colonic neoplasia prevalence and ASGE recommendations: Is it time for a change? *J Clin Gastroenterol* 2008;42(8):886-91.

Khankari K, Eder M, Osborn CY, Makoul G, Clayman M, Skripkauskas S, et al. Improving colorectal cancer screening among the medically underserved: A pilot study within a federally qualified health center. *J Gen Intern Med* 2007;22(10):1410-4.

Khatami S, Xuan L, Roman R, Zhang S, McConnel C, Halm EA, et al. Modestly Increased Use of Colonoscopy When Copayments Are Waived. *Clin Gastroenterol Hepatol* 2012;10(7):761-6.

Kim I-J, Kim K, Kang HC, Jang S-G, Park J-G. DHPLC analysis of adenomatous polyposis coli (APC) mutations using ready-to-use APC plates: Simple detection of multiple base pair deletion mutations. *Genet Test* 2008;12(2):295-8.

Kimberly JR, Phillips KC, Santago P, Perumpillichira J, Bechtold R, Pineau B, et al. Extracolonic findings at virtual colonoscopy: An important consideration in asymptomatic colorectal cancer screening. *J Gen Intern Med* 2009;24(1):69-73.

Ko HJ, Youn CH. Determination of the beginning age for colonoscopic screening among colonoscopy-naive individuals. *Clin Res Hepatol Gastroenterol* 2012;36(4):384-90.

Koga Y, Yasunaga M, Moriya Y, Akasu T, Fujita S, Yamamoto S, et al. Detection of colorectal cancer cells from feces using quantitative real-time RT-PCR for colorectal cancer diagnosis. *Cancer Sci* 2008;99(10):1977-83.

Koga Y, Yasunaga M, Moriya Y, Akasu T, Fujita S, Yamamoto S, et al. Detection of the DNA point mutation of colorectal cancer cells isolated from feces stored under different conditions. *Jpn J Clin Oncol* 2009;39(1):62-9.

Kremers SPJ, Mesters I, Pladdet IE, Van den Borne B, Stockbrugger RW. Participation in a sigmoidoscopic colorectal cancer screening program: A pilot study. *Cancer Epidemiol Biomarkers Prev* 2000;9(10):1127-30.

Laiyemo AO, Doubeni C, Pinsky PF, Doria-Rose VP, Sanderson AK, Bresalier R, et al. Factors associated with inadequate colorectal cancer screening with flexible sigmoidoscopy. *Cancer Epidemiol* 2012;36(4):395-9.

Lathroum L, Ramos-Mercado F, Hernandez-Marrero J, Villafana M, Cruz-Correa M. Ethnic and Sex Disparities in Colorectal Neoplasia Among Hispanic Patients Undergoing Screening Colonoscopy. *Clin Gastroenterol Hepatol* 2012;10(9):997-1001.

Lau A, Gregor JC. Resource implications for a population-based colorectal cancer screening program in Canada: a study of the impact on colonoscopy capacity and costs in London, Ontario. *Can J Gastroenterol* 2007;21(6):371-7.

Lee CK, Park DI, Lee SH, Hwangbo Y, Eun CS, Han DS, et al. Participation by experienced endoscopy nurses increases the detection rate of colon polyps during a screening colonoscopy: a multicenter, prospective, randomized study. *Gastrointest Endosc* 2011;74(5):1094-102.

Lee JG, Lum D, Urayama S, Mann S, Saavedra S, Vigil H, et al. Extended flexible sigmoidoscopy performed by colonoscopists for colorectal cancer screening: a pilot study. *Aliment Pharmacol Ther* 2006;23(7):945-51.

Lee YC, Wang HP, Chiu HM, Lin CP, Huang SP, Lai YP, et al. Factors determining post-colonoscopy abdominal pain: prospective study of screening colonoscopy in 1000 subjects. *J Gastroenterol Hepatol* 2006;21(10):1575-80.

Lee YC, Chiu HM, Chiang TH, Yen AM-F, Chiu SY-H, Chen SL-S, et al. Accuracy of faecal occult blood test and Helicobacter pylori stool antigen test for detection of upper gastrointestinal lesions. *BMJ Open* 2013;3(10):e003989.

Lei WY, Yi CH, Liu TT, Chen CL. Colonoscopic performance in a sedated health screening population: influence of colonoscope type. *Hepatogastroenterology* 2012;59(114):436-9.

Leufkens AM, DeMarco DC, Rastogi A, Akerman PA, Azzouzi K, Rothstein RI, et al. Effect of a retrograde-viewing device on adenoma detection rate during colonoscopy: the TERRACE study. *Gastrointest Endosc* 2011;73(3):480-9.

Leung FW, Cheung R, Fan RS, Fischer LS, Friedland S, Ho SB, et al. The water exchange method for colonoscopy-effect of coaching. *J Intervent Gastroenterol* 2012;2(3):122-5.

Leung K, Pinsky P, Laiyemo AO, Lanza E, Schatzkin A, Schoen RE. Ongoing colorectal cancer risk despite surveillance colonoscopy: the Polyp Prevention Trial Continued Follow-up Study. *Gastrointest Endosc* 2010;71(1):111-7.

Levin TR, Zhao W, Conell C, Seeff LC, Manninen DL, Shapiro JA, et al. Complications of colonoscopy in an integrated health care delivery system. *Ann Intern Med* 2006;145(12):880-6.

Li S, Wang J, Lu Y, Fan D. Screening and early diagnosis of colorectal cancer in China: a 12 year retrospect (1994-2006). *J Cancer Res Clin Oncol* 2007;133(10):679-86.

Lieberman DA, Weiss DG, Veterans Affairs Cooperative Study Group. One-time screening for colorectal cancer with combined fecal occult-blood testing and examination of the distal colon. *N Engl J Med* 2001;345(8):555-60.

Lin OS, Schembre DB, McCormick SE, Gluck M, Patterson DJ, Jiranek GC, et al. Risk of proximal colorectal neoplasia among asymptomatic patients with distal hyperplastic polyps. *Am J Med* 2005;118(10):1113-9.

Lin OS, Kozarek RA, Schembre DB, Ayub K, Gluck M, Drennan F, et al. Screening colonoscopy in very elderly patients: prevalence of neoplasia and estimated impact on life expectancy. *JAMA : the journal of the American Medical Association* 2006;295(20):2357-65.

Lin OS, Kozarek RA, Arai A, Gluck M, Jiranek GC, Kowdley KV, et al. The effect of periodic monitoring and feedback on screening colonoscopy withdrawal times, polyp detection rates, and patient satisfaction scores. *Gastrointest Endosc* 2010;71(7):1253-9.

Link A, Balaguer F, Shen Y, Nagasaka T, Lozano JJ, Boland CR, et al. Fecal microRNAs as novel biomarkers for colon cancer screening. *Cancer Epidemiol Biomarkers Prev* 2010;19(7):1766-74.

Liu J, Yao J, Summers RM. Scale-based scatter correction for computer-aided polyp detection in CT colonography. *Med Phys* 2008;35(12):5664-71.

Lucendo AJ, Guagnozzi D, Angueira T, Gonzalez-Castillo S, Fernandez-Fuente M, Frigal-Ruiz AB, et al. The relationship between proximal and distal colonic adenomas: Is screening sigmoidoscopy enough in the presence of a changing epidemiology? *Eur J Gastroenterol Hepatol* 2013;25(8):973-80.

Lui YW, MacAri M, Israel G, Bini EJ, Wang H, Babb J. CT Colonography Data Interpretation: Effect of Different Section Thicknesses-Preliminary Observations. *Radiology* 2003;229(3):791-7.

Luo D, Cambon AC, Wu D. Evaluating the long-term effect of FOBT in colorectal cancer screening. *Cancer Epidemiol* 2012;36(1):e54-e60.

MacK LA, Cook LS, Temple WJ, Carlson LE, Hilsden RJ, Paolucci EO. Colorectal cancer screening among first-degree relatives of colorectal cancer patients: Benefits and barriers. *Ann Surg Oncol* 2009;16(8):2092-100.

Maheshwari S, Patel T, Patel P. Screening for colorectal cancer in elderly persons: Who should we screen and when can we stop? *J Aging Health* 2008;20(1):126-39.

Matsuda T, Fujii T, Sano Y, Kudo SE, Oda Y, Igarashi M, et al. Five-year incidence of advanced neoplasia after initial colonoscopy in Japan: a multicenter retrospective cohort study. *Jpn J Clin Oncol* 2009;39(7):435-42.

McDonald R, Tomlins A, Smith S, Harmston C. Outcomes of faecal occult blood tests requested outside the UK National Bowel Cancer Screening Programme. *J Clin Pathol* 2013;66(4):330-4.

McDonald S, Lyall P, Israel L, Coates R, Frizelle FA. Why barium enemas fail to identify colorectal cancers. *ANZ J Surg* 2001;71(11):631-3.

McFarland EG, Brink JA, Pilgram TK, Heiken JP, Balfe DM, Hirselj DA, et al. Spiral CT colonography: Reader agreement and diagnostic performance with two- and three-dimensional image-display techniques. *Radiology* 2001;218(2):375-83.

Menges M, Gartner B, Georg T, Fischinger J, Zeitz M. Cost-benefit analysis of screening colonoscopy in 40- to 50-year-old first-degree relatives of patients with colorectal cancer. *Int J Colorectal Dis* 2006;21(6):596-601.

Menges M, Fischinger J, Gartner B, Georg T, Woerdehoff D, Maier M, et al. Screening colonoscopy in 40- to 50-year-old first-degree relatives of patients with colorectal cancer is efficient: A controlled multicentre study. *Int J Colorectal Dis* 2006;21(4):301-7.

Mesihovic R, Vanis N, Gribajcevic M. Test for obscure bleeding vs colonoscopy in the prevention of colorectal cancer. *Med Arh* 2008;62(3):153-5.

Miles A, Wardle J, Atkin W. Receiving a screen-detected diagnosis of cancer: the experience of participants in the UK flexible sigmoidoscopy trial. *Psychooncology* 2003;12(8):784-802.

Mizukami T, Ogata H, Hibi T. "Passive-bending colonoscope" significantly improves cecal intubation in difficult cases. *World J Gastroenterol* 2012;18(32):4454-6.

Morfoisse JJ, Grasset D, Seigneuric C. [Patient opinions after colonoscopy. Results of a cross sectional, multicenter ANGH study (2)]. *Opinion des malades apres coloscopie Resultats d'une enquetetransversale multicentrique de l'ANGH (2)* 2000;24(3):279-83.

Mudawi HMY, Mohammed Ali SEA, Dabora AA, El Tahir MA, Suliman SH, Salim OEFH, et al. American Society for Gastrointestinal Endoscopy guidelines for appropriate use of colonoscopy: are they suitable for African patients? *Trop Doct* 2012;42(3):165-7.

Muinuddin A, Aslahi R, Hopman WM, Paterson WG. Relationship between the number of positive fecal occult blood tests and the diagnostic yield of colonoscopy. *Can J Gastroenterol* 2013;27(2):90-4.

Murff HJ, Peterson NB, Fowke JH, Hargreaves M, Signorello LB, Dittus RS, et al. Colonoscopy screening in African Americans and Whites with affected first-degree relatives. *Arch Intern Med* 2008;168(6):625-31.

Myers RE, Fishbein G, Hyslop T, Hauck WW, Kutch M, Grana JR, et al. Measuring complete diagnostic evaluation in colorectal cancer screening. *Cancer Detect Prev* 2001;25(2):174-82.

Nagorni A, Bjelakovic G, Petrovic B. Narrow band imaging versus conventional white light colonoscopy for the detection of colorectal polyps. *Cochrane Database Syst Rev* 2012;1:CD008361.

Nakama H, Fattah AS, Zhang B, Kamijo N. Digital rectal examination sampling of stool is less predictive of significant colorectal pathology than stool passed spontaneously. *Eur J Gastroenterol Hepatol* 2000;12(11):1235-8.

Nakama H, Zhang B, Fattah ASMA. A cost-effective analysis of the optimum number of stool specimens collected for immunochemical occult blood screening for colorectal cancer. *Eur J Cancer* 2000;36(5):647-50.

Nakao SK, Fassler S, Sucandy I, Kim S, Zebley DM. Colorectal cancer following negative colonoscopy: is 5-year screening the correct interval to recommend? *Surg Endosc* 2013;27(3):768-73.

Nakazato M, Yamano H-O, Matsushita H-O, Sato K, Fujita K, Yamanaka Y, et al. Immunologic fecal occult blood test for colorectal cancer screening. *Japan Med Assoc J* 2006;49(5-6):203-7.

Nappi JJ, Frimmel H, Dachman AH, Yoshida H. Computerized detection of colorectal masses in CT colonography based on fuzzy merging and wall-thickening analysis. *Med Phys* 2004;31(4):860-72.

Nguyen TB, Wang S, Anugu V, Rose N, McKenna M, Petrick N, et al. Distributed human intelligence for colonic polyp classification in computer-aided detection for CT colonography. *Radiology* 2012;262(3):824-33.

Nishihara R, Wu K, Lochhead P, Morikawa T, Liao X, Qian ZR, et al. Long-term colorectal-cancer incidence and mortality after lower endoscopy. *N Engl J Med* 2013;369(12):1095-105.

Olson J, Whitney DH, Durkee K, Shuber AP. DNA stabilization is critical for maximizing performance of fecal DNA-based colorectal cancer tests. *Diagn Mol Pathol* 2005;14(3):183-91.

Oort FA, van Turenhout ST, Coupe VMH, van der Hulst RWM, Wesdorp EIC, Terhaar sive Droste JS, et al. Double sampling of a faecal immunochemical test is not superior to single sampling for detection of colorectal neoplasia: a colonoscopy controlled prospective cohort study. *BMC cancer* 2011;11:434.

Othman MO, Bradley AG, Choudhary A, Hoffman RM, Roy PK. Variable stiffness colonoscope versus regular adult colonoscope: meta-analysis of randomized controlled trials. *Endoscopy* 2009;41(1):17-24.

Ozel B, Pickhardt PJ, Kim DH, Schumacher C, Bhargava N, Winter TC. Accuracy of routine nontargeted CT without colonography technique for the detection of large colorectal polyps and cancer. *Dis Colon Rectum* 2010;53(6):911-8.

Pabby A, Suneja A, Heeren T, Farraye FA. Flexible sigmoidoscopy for colorectal cancer screening in the elderly. *Dig Dis Sci* 2005;50(11):2147-52.

Parente F, Marino B, DeVecchi N, Moretti R. Faecal occult blood test-based screening programme with high compliance for colonoscopy has a strong clinical impact on colorectal cancer. *Br J Surg* 2009;96(5):533-40.

Park SK, Park DI, Lee S-Y, Lee SK, Kim Y-H, Lee SJ, et al. Extracolonic findings of computed tomographic colonography in Koreans. *World J Gastroenterol* 2009;15(12):1487-92.

Parlak E, Yildirim M, Cekin AH, Yildiz M, Dogan AN, Dilli UD. Role of ultrasonography in diagnosis of cancers of the gastrointestinal tract. *J Clin Anal Med* 2013;4(5):377-80.

Peacock O, Watts ES, Hanna N, Kerr K, Goddard AF, Lund JN. Inappropriate use of the faecal occult blood test outside of the National Health Service colorectal cancer screening programme. *Eur J Gastroenterol Hepatol* 2012;24(11):1270-5.

Pezzoli A, Matarese V, Rubini M, Simoni M, Caravelli GC, Stockbrugger R, et al. Colorectal cancer screening: Results of a 5-year program in asymptomatic subjects at increased risk. *Dig Liver Dis* 2007;39(1):33-9.

Phillips KL, Smith ML, Ahn S, Ory MG, Hochhalter AK. Correlates of initiating colorectal cancer screening beginning at age 50. *J Community Health* 2013;38(1):23-30.

Pickhardt PJ, Taylor AJ, Gopal DV. Surface visualization at 3D endoluminal CT colonography: degree of coverage and implications for polyp detection. *Gastroenterology* 2006;130(6):1582-7.

Puente Gutierrez JJ, Marin Moreno MA, Dominguez Jimenez JL, Bernal BE, Diaz Iglesias JM. Effectiveness of a colonoscopic screening programme in first-degree relatives of patients with colorectal cancer. *Colorectal Dis* 2011;13(6):e145-e153.

Rabeneck L, Paszat LF, Saskin R, Stukel TA. Association between colonoscopy rates and colorectal cancer mortality. *Am J Gastroenterol* 2010;105(7):1627-32.

Radaelli F, Paggi S, Minoli G. Variation of quality of colonoscopy in Italy over five years: a nation-wide observational study. *Dig Liver Dis* 2013;45(1):28-32.

Rastogi A, Bansal A, Wani S, Callahan P, McGregor DH, Cherian R, et al. Narrow-band imaging colonoscopy--a pilot feasibility study for the detection of polyps and correlation of surface patterns with polyp histologic diagnosis. *Gastrointest Endosc* 2008;67(2):280-6.

Ravizza D, Bartolomei M, Santoro L, Tamayo D, Fiori G, Trovato C, et al. Positron emission tomography for the detection of colorectal adenomas. *Dig Liver Dis* 2010;42(3):185-90.

Regula J, Rupinski M, Kraszewska E, Polkowski M, Pachlewski J, Orlowska J, et al. Colonoscopy in colorectal-cancer screening for detection of advanced neoplasia. *N Engl J Med* 2006;355(18):1863-72.

Rennert G, Kislitsin D, Brenner DE, Rennert HS, Lev Z. Detecting K-ras mutations in stool from fecal occult blood test cards in multiphasic screening for colorectal cancer. *Cancer Lett* 2007;253(2):258-64.

Ross JJ, Cooley JS, Lee DL. Septic thrombophlebitis of the hepatic veins after colonoscopy: a case report. *Gastrointest Endosc* 2003;58(1):155-6.

Rundle AG, Lebowitz B, Vogel R, Levine S, Neugut AI. Colonoscopic Screening in Average-Risk Individuals Ages 40 to 49 vs 50 to 59 Years. *Gastroenterology* 2008;134(5):1311-5.

Sakamoto J, Fujiya M, Okamoto K, Nata T, Inaba Y, Moriichi K, et al. Immunoprecipitation of nucleosomal DNA is a novel procedure to improve the sensitivity of serum screening for the p16 hypermethylation associated with colon cancer. *Cancer Epidemiol* 2010;34(2):194-9.

Sakamoto T, Mitsuzaki K, Utsunomiya D, Matsuda K, Yamamura S, Urata J, et al. Detection of flat colorectal polyps at screening CT colonography in comparison with conventional polypoid lesions. *Acta Radiol* 2012;53(7):714-9.

Sali L, Falchini M, Della Monica P, Regge D, Bonanomi AG, Castiglione G, et al. CT colonography before colonoscopy in subjects with positive faecal occult blood test. Preliminary experience. *Radiol Med* 2010;115(8):1267-78.

Salonen A, Nikkila J, Jalanka-Tuovinen J, Immonen O, Rajilic-Stojanovic M, Kekkonen RA, et al. Comparative analysis of fecal DNA extraction methods with phylogenetic microarray: Effective recovery of bacterial and archaeal DNA using mechanical cell lysis. *J Microbiol Methods* 2010;81(2):127-34.

Santavirta J. Screening, sigmoidoscopy in persons aged 60 years. *Colorectal Dis* 2002;4(3):184-8.

Sato R, Fujiya M, Watari J, Ueno N, Moriichi K, Kashima S, et al. The diagnostic accuracy of high-resolution endoscopy, autofluorescence imaging and narrow-band imaging for differentially diagnosing colon adenoma. *Endoscopy* 2011;43(10):862-8.

Schoen RE, Pinsky PF, Weissfeld JL, Bresalier RS, Church T, Prorok P, et al. Results of repeat sigmoidoscopy 3 years after a negative examination. *JAMA : the journal of the American Medical Association* 2003;290(1):41-8.

Schoenfeld P, Cash B, Flood A, Dobhan R, Eastone J, Coyle W, et al. Colonoscopic screening of average-risk women for colorectal neoplasia. *New Engl J Med* 2005;352(20):2061-8.

Schoenfeld P, Shad J, Ormseth E, Coyle W, Cash B, Butler J, et al. Predictive value of diminutive colonic adenoma trial: the PREDICT trial. *Clin Gastroenterol Hepatol* 2003;1(3):195-201.

Schonfeld V, Hinzmann S. Missed colonic adenomas in routine primary care endoscopy: a prospective tandem colonoscopy study. *Z Gastroenterol* 2010;48(10):1207-10.

Schroy PC, Coe AM, Mylvaganam SR, Ahn LB, Lydotes MA, Robinson PA, et al. The Your Disease Risk Index for colorectal cancer is an inaccurate risk stratification tool for advanced colorectal neoplasia at screening colonoscopy. *Cancer Prev Res (Phila)* 2012;5(8):1044-52.

Senore C, Segnan N, Bonelli L, Sciallero S, Pennazio M, Angioli D, et al. Predicting proximal advanced neoplasms at screening sigmoidoscopy. *Dis Colon Rectum* 2004;47(8):1331-40.

Seres KA, Kirkpatrick AC, Tierney WM. The utility of an evidence-based lecture and clinical prompt as methods to improve quality of care in colorectal cancer screening. *Am J Gastroenterol* 2009;104(2):420-5.

Shapiro TF, Hoover J, Paszat LF, Burgis E, Hsieh E, Rothwell DM, et al. Colorectal cancer screening with nurse-performed flexible sigmoidoscopy: results from a Canadian community-based program. *Gastrointest Endosc* 2007;65(4):640-5.

Sherer EA, Imler TD, Imperiale TF. The effect of colonoscopy preparation quality on adenoma detection rates. *Gastrointest Endosc* 2012;75(3):545-53.

Shuhaibar M, Walsh C, Lindsay F, Lee N, Walsh P, O'Gorman P, et al. A comparative study of faecal occult blood kits in a colorectal cancer screening program in a cohort of healthy construction workers. *Ir J Med Sci* 2011;180(1):103-8.

Siersema PD, Rastogi A, Leufkens AM, Akerman PA, Azzouzi K, Rothstein RI, et al. Retrograde-viewing device improves adenoma detection rate in colonoscopies for surveillance and diagnostic workup. *World J Gastroenterol* 2012;18(26):3400-8.

Singh M, Mehta N, Murthy UK, Kaul V, Arif A, Newman N. Postpolypectomy bleeding in patients undergoing colonoscopy on uninterrupted clopidogrel therapy. *Gastrointest Endosc* 2010;71(6):998-1005.

Smith A, Young GP, Cole SR, Bampton P. Comparison of a brush-sampling fecal immunochemical test for hemoglobin with a sensitive guaiac-based fecal occult blood test in detection of colorectal neoplasia. *Cancer* 2006;107(9):2152-9.

Sohns C, Heuser M, Sossalla S, Wolff H, Obenauer S. Current role and future potential of computed tomographic colonography for colorectal polyp detection and colon cancer screening-incidental findings. *Clin Imaging* 2008;32(4):280-6.

Souques M, Zummer K. [The Hemocult II test: results of 16 years of screening tests at the Tumor Prevention Service of the City of Paris]. *Le test Hemocult II: resultats de 16 ans de pratique dans le Service de Prevention et Depistage des Tumeurs de la Ville de Paris* 2000;29(18):983-6.

Soyer P, Sirol M, Dray X, Place V, Pautrat K, Hamzi L, et al. Detection of colorectal tumors with water enema-multidetector row computed tomography. *Abdom Imaging* 2012;37(6):1092-100.

Steele RJC, McClements P, Watling C, Libby G, Weller D, Brewster DH, et al. Interval cancers in a FOBT-based colorectal cancer population screening programme: Implications for stage, gender and tumour site. *Gut* 2012;61(4):576-81.

Stegeman I, de Wijkerslooth TR, Mallant-Hent RC, de Groot K, Stroobants AK, Fockens P, et al. Implementation of population screening for colorectal cancer by repeated Fecal Immunochemical Test (FIT): third round. *BMC Gastroenterol* 2012;12:73.

Steinmetz J, Spyckerelle Y, Gueguen R, Dupre C. Colorectal cancer screening in Health Examination Centers: Evaluation in 50-74-year-old subjects between 1998 and 2003. *Gastroenterol Clin Biol* 2006;30(6-7):832-7.

Steinmetz J, Spyckerelle Y, Giordanella JP. [Colorectal cancer: assessment of two-yearly screening of the population in health care examination centres]. *Cancer colorectal: evaluation du depistage biennal dans la population des centres d'examens de sante* 2003;32(32):1496-9.

Stock C, Ihle P, Schubert I, Brenner H. Colonoscopy and fecal occult blood test use in Germany: Results from a large insurance-based cohort. *Endoscopy* 2011;43(9):771-9.

Stock C, Hoffmeister M, Birkner B, Brenner H. Performance of additional colonoscopies and yield of neoplasms within 3 years after screening colonoscopy: A historical cohort study. *Endoscopy* 2013;45(7):537-44.

Strul H, Kariv R, Leshno M, Halak A, Jakubowicz M, Santo M, et al. The prevalence rate and anatomic location of colorectal adenoma and cancer detected by colonoscopy in average-risk individuals aged 40-80 years. *Am J Gastroenterol* 2006;101(2):255-62.

Su M-Y, Tung S-Y, Lien J-M, Chiu C-T, Chen P-C, Wu C-S. Magnifying endoscopy with indigo carmine contrast for the differential diagnosis of adenomatous and hyperplastic colonic polyps. *Gastroenterol J Taiwan* 2000;17(3):162-8.

Su MY, Ho YP, Hsu CM, Chiu CT, Chen PC, Lien JM, et al. How can colorectal neoplasms be treated during colonoscopy? *World J Gastroenterol* 2005;11(18):2806-10.

Suceveanu AI, Suceveanu A, Dumitru E, Alexandrescu L, Voinea F. The feasibility of FOBT tests in colorectal cancer screening in Dobrogea. *Rom J Gastroenterol* 2005;14(3):213-7.

Sulu B, Yildiz BD, Buyukuysal C, Demir E, Gunerhan Y. Comparison of meperidine versus hyoscine during colonoscopy in the elderly: a prospective randomized study. *J Laparoendosc Adv Surg Tech A* 2012;22(7):631-4.

Summers RM, Handwerker LR, Pickhardt PJ, Van Uitert RL, Deshpande KK, Yeshwant S, et al. Performance of a previously validated CT colonography computer-aided detection system in a new patient population. *AJR Am J Roentgenol* 2008;191(1):168-74.

Synnestvedt MB, Weiner MG. Evaluation of the use of electronic health data to classify four-year mortality risk for older adults undergoing screening colonoscopies. *AMIA Annu Symp Proc* 2011;2011:1361-5.

Szura M, Bucki K, Matyja A, Kulig J. Evaluation of magnetic scope navigation in screening endoscopic examination of colorectal cancer. *Surg Endosc* 2012;26(3):632-8.

Tawn DJ, Squire CJ, Mohammed MA, Adam EJ. National audit of the sensitivity of double-contrast barium enema for colorectal carcinoma, using control charts. *Clin Radiol* 2005;60(5):558-64.

Taylor SA, Burling D, Roddie M, Honeyfield L, McQuillan J, Bassett P, et al. Computer-aided detection for CT colonography: Incremental benefit of observer training. *Br J Radiol* 2008;81(963):180-6.

Taylor T, Williamson S, Wardle J, Borrill J, Sutton S, Atkin W. Acceptability of flexible sigmoidoscopy screening in older adults in the United Kingdom. *J Med Screen* 2000;7(1):38-45.

Testoni PA, Notaristefano C, Vailati C, Di Leo M, Viale E. High-definition colonoscopy with i-Scan: better diagnosis for small polyps and flat adenomas. *World J Gastroenterol* 2012;18(37):5231-9.

Thoma MN, Castro F, Golawala M, Chen R. Detection of colorectal neoplasia by colonoscopy in average-risk patients age 40-49 versus 50-59 years. *Dig Dis Sci* 2011;56(5):1503-8.

Tiro JA, Vernon SW, Hyslop T, Myers RE. Factorial validity and invariance of a survey measuring psychosocial correlates of colorectal cancer screening among African Americans and Caucasians. *Cancer Epidemiol Biomarkers Prev* 2005;14(12):2855-61.

Triantafillidis JK, Kosmidis PA, Papalois A, Merikas E, Cheracakis P, Govosdis V, et al. Screening programs for colorectal cancer in Greece: Results of two pilot studies conducted in March 2008 and 2009. *Ann Gastroenterol* 2010;23(1):42-7.

Van Kleek E, Liu S, Conn LM, Hoadley A, Ho SB. Improving the effectiveness of fecal occult blood testing in a primary care clinic by direct colonoscopy referral for positive tests. *J Healthc Qual* 2010;32(6):62-9.

van Rijn AF, van Rossum LGM, Deutekom M, Laheij RJF, Fockens P, Bossuyt PMM, et al. Low priority main reason not to participate in a colorectal cancer screening program with a faecal occult blood test. *J Public Health (Oxf)* 2008;30(4):461-5.

van Roon AHC, Hol L, van Vuuren AJ, Francke J, Ouwendijk M, Heijens A, et al. Are fecal immunochemical test characteristics influenced by sample return time? A population-based colorectal cancer screening trial. *Am J Gastroenterol* 2012;107(1):99-107.

van Roon AHC, Goede SL, van Ballegooijen M, van Vuuren AJ, Looman CWN, Biermann K, et al. Random comparison of repeated faecal immunochemical testing at different intervals for population-based colorectal cancer screening. *Gut* 2013;62(3):409-15.

van Turenhout ST, Oort FA, Terhaar sive Droste JS, Coupe VMH, van der Hulst RW, Loffeld RJ, et al. Hemorrhoids detected at colonoscopy: an infrequent cause of false-positive fecal immunochemical test results. *Gastrointest Endosc* 2012;76(1):136-43.

Veerappan GR, Ally MR, Choi J-H, Pak JS, Maydonovitch C, Wong RKH. Extracolonic findings on CT colonography increases yield of colorectal cancer screening. *Am J Roentgenol* 2010;195(3):677-86.

Villeneuve PJ, Coombs A. Screening for colorectal cancer using the fecal occult blood test: an actuarial assessment of the impact of a population-based screening program in Canada. *Int J Technol Assess Health Care* 2003;19(4):715-23.

Wan J, Zhang Z-Q, Zhu C, Wang M-W, Zhao D-H, Fu Y-H, et al. Colonoscopic screening and follow-up for colorectal cancer in the elderly. *World J Gastroenterol* 2002;8(2):267-9.

Wang HS, Pisegna J, Modi R, Liang LJ, Atia M, Nguyen M, et al. Adenoma detection rate is necessary but insufficient for distinguishing high versus low endoscopist performance. *Gastrointest Endosc* 2013;77(1):71-8.

Weissfeld JL, Schoen RE, Pinsky PF, Bresalier RS, Doria-Rose VP, Laiyemo AO, et al. Flexible sigmoidoscopy in the randomized prostate, lung, colorectal, and ovarian (PLCO) cancer screening trial: added yield from a second screening examination. *J Natl Cancer Inst* 2012;104(4):280-9.

Wilkes JM, DiPalma JA. Brachial blood pressure monitoring versus ankle monitoring during colonoscopy. *South Med J* 2004;97(10):939-41.

Wong G, Howard K, Chapman JR, Tong A, Bourke MJ, Hayen A, et al. Test performance of faecal occult blood testing for the detection of bowel cancer in people with chronic kidney disease (DETECT) protocol. *BMC public health* 2011;11:516.

Wu D, Erwin D, Rosner GL. Estimating key parameters in FOBT screening for colorectal cancer. *Cancer causes & control : CCC* 2009;20(1):41-6.

Xirasagar S, Hurley TG, Sros L, Hebert JR. Quality and safety of screening colonoscopies performed by primary care physicians with standby specialist support. *Med Care* 2010;48(8):703-9.

Xu JW, Suzuki K. Massive-training support vector regression and Gaussian process for false-positive reduction in computer-aided detection of polyps in CT colonography. *Med Phys* 2011;38(4):1888-902.

Yamaguchi A, Hashimoto N, Tsutae W, Seino K, Osanai H, Yamamoto N, et al. Dried feces spots as an alternative DNA source: Detection of K-ras mutations in colorectal cancer screening. *Tumor Res* 2001;36:7-13.

Yang X, Slabaugh G. A robust and efficient approach to detect 3D rectal tubes from CT colonography. *Med Phys* 2011;38(11):6238-47.

Yen AW, Leung JW, Leung FW. A new method for screening and surveillance colonoscopy: Combined water-exchange and cap-assisted colonoscopy. *J Intervent Gastroenterol* 2012;2(3):114-9.

Yun Jy, Ro HJ, Park JB, Choi JB, Chung JE, Kim YJ, et al. Diagnostic performance of CT colonography for the detection of colorectal polyps. *Korean J Radiol* 2007;8(6):484-91.

Zarchy TM, Tsai F, Ramicone E, Chan LS. A risk profile for advanced proximal neoplasms on diagnostic colonoscopy. *Dig Dis Sci* 2009;54(1):151-9.

Zauber AG, Winawer SJ, O'Brien MJ, Lansdorp-Vogelaar I, van Ballegooijen M, Hankey BF, et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med* 2012;366(8):687-96.

Zheng S, Liu XY, Ding KF, Wang LB, Qiu PL, Ding XF, et al. Reduction of the incidence and mortality of rectal cancer by polypectomy: a prospective cohort study in Haining County. *World J Gastroenterol* 2002;8(3):488-92.

Excluded No Outcomes of Interest

Achiam MP, Logager VB, Chabanova E, Eegholm B, Thomsen HS, Rosenberg J. Diagnostic accuracy of MR colonography with fecal tagging. *Abdom Imaging* 2009;34(4):483-90.

Adler S, Hassan C, Metzger Y, Sompolinsky Y, Spada C. Accuracy of automatic detection of small-bowel mucosa by second-generation colon capsule endoscopy. *Gastrointest Endosc* 2012;76(6):1170-4.

Ahlquist DA, Zou H, Domanico M, Mahoney DW, Yab TC, Taylor WR, et al. Next-generation stool DNA test accurately detects colorectal cancer and large adenomas. *Gastroenterology* 2012;142(2):248-56.

Allison JE, Fraser CG. Fecal immunochemical tests had better sensitivity for detecting colorectal neoplasia in aspirin users than in nonusers. *Ann Intern Med* 2011;154(8):JC4-10.

Andreoni B, Crosta C, Lotti M, Carloni M, Marzona L, Biffi R, et al. Flexible sigmoidoscopy as a colorectal cancer screening test in the general population: recruitment phase results of a randomized controlled trial in Lombardia, Italy. *Chir Ital* 2000;52(3):257-62.

Aragones A, Schwartz MD, Shah NR, Gany FM. A randomized controlled trial of a multilevel intervention to increase colorectal cancer screening among Latino immigrants in a primary care facility. *J Gen Intern Med* 2010;25(6):564-7.

Araujo SEA, Seid VE, Caravatto PDP, Dumarco R. Incidence and management of colonoscopic colon perforations: 10 years' experience. *Hepatology* 2009;56(96):1633-6.

Arnesen RB, Adamsen S, Svendsen LB, Raaschou HO, von Benzou E, Hansen OH. Missed lesions and false-positive findings on computed-tomographic colonography: a controlled prospective analysis. *Endoscopy* 2005;37(10):937-44.

Arnesen RB, von Benzou E, Adamsen S, Svendsen LB, Raaschou HO, Hansen OH. Diagnostic performance of computed tomography colonography and colonoscopy: a prospective and validated analysis of 231 paired examinations. *Acta Radiol* 2007;48(8):831-7.

Baier M, Calonge N, Cutter G, McClatchey M, Schoentgen S, Hines S, et al. Validity of self-reported colorectal cancer screening behavior. *Cancer Epidemiol Biomarkers Prev* 2000;9(2):229-32.

Baker SG, Kramer BS. Estimating the cumulative risk of a false-positive under a regimen involving various types of cancer screening tests. *J Med Screen* 2008;15(1):18-22.

Bhangu A, Bowley DM, Horner R, Baranowski E, Raman S, Karandikar S. Volume and accreditation, but not specialty, affect quality standards in colonoscopy. *Br J Surg* 2012;99(10):1436-44.

Braun KL, Fong M, Kaanoi ME, Kamaka ML, Gotay CC. Testing a culturally appropriate, theory-based intervention to improve colorectal cancer screening among Native Hawaiians. *Prev Med* 2005;40(6):619-27.

Brenner H, Hoffmeister M, Arndt V, Stegmaier C, Altenhofen L, Haug U. Protection from right- and left-sided colorectal neoplasms after colonoscopy: population-based study. *J Natl Cancer Inst* 2010;102(2):89-95.

Buchner AM, Shahid MW, Heckman MG, McNeil RB, Cleveland P, Gill KR, et al. High-definition colonoscopy detects colorectal polyps at a higher rate than standard white-light colonoscopy. *Clin Gastroenterol Hepatol* 2010;8(4):364-70.

Burnand B, Harris JK, Wietlisbach V, Froehlich F, Vader JP, Gonvers JJ. Use, appropriateness, and diagnostic yield of screening colonoscopy: an international observational study (EPAGE). *Gastrointest Endosc* 2006;63(7):1018-26.

Castells A, Bessa X, Quintero E, Bujanda L, Cubiella J, Salas D, et al. Risk of advanced proximal neoplasms according to distal colorectal findings: comparison of sigmoidoscopy-based strategies. *J Natl Cancer Inst* 2013;105(12):878-86.

Chatman N, Sutherland JR, van der Zwan R, Abraham N. A survey of patient understanding and expectations of sedation/anaesthesia for colonoscopy. *Anaesth Intensive Care* 2013;41(3):369-73.

Chung SJ, Kim YS, Yang SY, Song JH, Park MJ, Kim JS, et al. Prevalence and risk of colorectal adenoma in asymptomatic Koreans aged 40-49 years undergoing screening colonoscopy. *J Gastroenterol Hepatol* 2010;25(3):519-25.

Ciatto S, Castiglione G. Role of double-contrast barium enema in colorectal cancer screening based on fecal occult blood. *Tumori* 2002;88(2):95-8.

Corley DA, Jensen CD, Marks AR. Can we improve adenoma detection rates? A systematic review of intervention studies. *Gastrointest Endosc* 2011;74(3):656-65.

Cornett D, Barancin C, Roeder B, Reichelderfer M, Frick T, Gopal D, et al. Findings on optical colonoscopy after positive CT colonography exam. *Am J Gastroenterol* 2008;103(8):2068-74.

Coronado GD, Golovaty I, Longton G, Levy L, Jimenez R. Effectiveness of a clinic-based colorectal cancer screening promotion program for underserved Hispanics. *Cancer* 2011;117(8):1745-54.

Cotton PB, Durkalski VL, Pineau BC, Palesch YY, Mauldin PD, Hoffman B, et al. Computed tomographic colonography: What results should we believe? *Evid -Based Gastroenterol* 2004;5(3):84-5.

Croswell JM, Kramer BS, Kreimer AR, Prorok PC, Xu JL, Baker SG, et al. Cumulative incidence of false-positive results in repeated, multimodal cancer screening. *Ann Fam Med* 2009;7(3):212-22.

Cruz-Correa M, Schultz K, Jagannath S, Harris M, Kantsevov S, Bedine M, et al. Performance characteristics and comparison of two fecal occult blood tests in patients undergoing colonoscopy. *Dig Dis Sci* 2007;52(4):1009-13.

de Wijkerslooth TR, de Haan MC, Stoop EM, Deutekom M, Fockens P, Bossuyt PMM, et al. Study protocol: population screening for colorectal cancer by colonoscopy or CT colonography: a randomized controlled trial. *BMC Gastroenterol* 2010;10:47.

Doria-Rose VP, Levin TR, Selby JV, Newcomb PA, Richert-Boe KE, Weiss NS. The incidence of colorectal cancer following a negative screening sigmoidoscopy: Implications for screening interval. *Gastroenterology* 2004;127(3):714-22.

Doubeni CA, Weinmann S, Adams K, Kamineni A, Buist DSM, Ash AS, et al. Screening colonoscopy and risk for incident late-stage colorectal cancer diagnosis in average-risk adults: A nested case-control study. *Ann Intern Med* 2013;158(5 PART 1):312-20.

Egan CA, Smith EP, Taylor TB, Meyer LJ, Samowitz WS, Zone JJ. Fecal occult blood testing in a noncompliant inner city minority population: Increased compliance and adherence to screening procedures without loss of test sensitivity using stool obtained at the time of in-office rectal examination. *Am J Gastroenterol* 2001;96(6):1908-13.

Eickhoff A, Van Dam J, Jakobs R, Kudis V, Hartmann D, Damian U, et al. Computer-assisted colonoscopy (the NeoGuide Endoscopy System): results of the first human clinical trial ("PACE study"). *Am J Gastroenterol* 2007;102(2):261-6.

Ellul P, Fogden E, Simpson CL, Nickerson CLR, Mckaig BC, Swarbrick ET, et al. Downstaging of colorectal cancer by the National Bowel Cancer Screening programme in England: first round data from the first centre. *Colorectal Dis* 2010;12(5):420-2.

Federici A, Giorgi Rossi P, Borgia P, Bartolozzi F, Farchi S, Gausticchi G. The immunochemical faecal occult blood test leads to higher compliance than the guaiac for colorectal cancer screening programmes: a cluster randomized controlled trial. *J Med Screen* 2005;12(2):83-8.

Ferrante JM, Lee JH, McCarthy EP, Fisher KJ, Chen R, Gonzalez EC, et al. Primary care utilization and colorectal cancer incidence and mortality among medicare beneficiaries: A population-based, case-control study. *Ann Intern Med* 2013;159(7):437-46.

Gluecker T, Dorta G, Keller W, Jornod P, Meuli R, Schnyder P. Performance of multidetector computed tomography colonography compared with conventional colonoscopy. *Gut* 2002;51(2):207-11.

Goodyear SJ, Leung E, Menon A, Pedamallu S, Williams N, Wong LS. The effects of population-based faecal occult blood test screening upon emergency colorectal cancer admissions in Coventry and north Warwickshire. *Gut* 2008;57(2):218-22.

Graser A, Stieber P, Nagel D, Schafer C, Horst D, Becker CR, et al. Comparison of CT colonography, colonoscopy, sigmoidoscopy and faecal occult blood tests for the detection of advanced adenoma in an average risk population. *Gut* 2009;58(2):241-8.

Gross CP, Soulos PR, Ross JS, Cramer LD, Guerrero C, Tinetti ME, et al. Assessing the impact of screening colonoscopy on mortality in the medicare population. *J Gen Intern Med* 2011;26(12):1441-9.

Gross SA, Buchner AM, Crook JE, Cangemi JR, Picco MF, Wolfsen HC, et al. A comparison of high definition-image enhanced colonoscopy and standard white-light colonoscopy for colorectal polyp detection. *Endoscopy* 2011;43(12):1045-51.

Groth S, Krause H, Behrendt R, Hill H, Borner M, Basturk M, et al. Capsule colonoscopy increases uptake of colorectal cancer screening. *BMC Gastroenterol* 2012;12:80.

Guittet L, Guillaume E, Levillain R, Beley P, Tichet J, Lantieri O, et al. Analytical comparison of three quantitative immunochemical fecal occult blood tests for colorectal cancer screening. *Cancer Epidemiol Biomarkers Prev* 2011;20(7):1492-501.

Gupta N, Bansal A, Rao D, Early DS, Jonnalagadda S, Edmundowicz SA, et al. Accuracy of in vivo optical diagnosis of colon polyp histology by narrow-band imaging in predicting colonoscopy surveillance intervals. *Gastrointest Endosc* 2012;75(3):494-502.

Gupta S, Halm EA, Rockey DC, Hammons M, Koch M, Carter E, et al. Comparative effectiveness of fecal immunochemical test outreach, colonoscopy outreach, and usual care for boosting colorectal cancer screening among the underserved a randomized clinical trial. *JAMA Intern Med* 2013;173(18):1725-32.

Halligan S, Taylor SA, Dehmeshki J, Amin H, Ye X, Tsang J, et al. Computer-assisted detection for CT colonography: external validation. *Clin Radiol* 2006;61(9):758-63.

Haug U, Knudsen AB, Brenner H, Kuntz KM. Is fecal occult blood testing more sensitive for left- versus right-sided colorectal neoplasia? A systematic literature review. *Expert Rev Mol Diagn* 2011;11(6):605-16.

Hoffman RM, Steel S, Yee EFT, Massie L, Schrader RM, Murata GH. Colorectal cancer screening adherence is higher with fecal immunochemical tests than guaiac-based fecal occult blood tests: a randomized, controlled trial. *Prev Med* 2010;50(5-6):297-9.

Hol L, de Jonge V, Van Leerdam ME, van Ballegooijen M, Looman CWN, Van Vuuren AJ, et al. Screening for colorectal cancer: comparison of perceived test burden of guaiac-based faecal occult blood test, faecal immunochemical test and flexible sigmoidoscopy. *Eur J Cancer* 2010;46(11):2059-66.

Holme O, Hoie O, Matre J, Stallemo A, Garborg K, Hasund A, et al. Magnetic endoscopic imaging versus standard colonoscopy in a routine colonoscopy setting: a randomized, controlled trial. *Gastrointest Endosc* 2011;73(6):1215-22.

Hong SN, Choe WH, Lee JH, Kim SI, Kim JH, Lee TY, et al. Prospective, randomized, back-to-back trial evaluating the usefulness of i-SCAN in screening colonoscopy. *Gastrointest Endosc* 2012;75(5):1011-21.

Horiuchi A, Nakayama Y, Kajiyama M, Fujii H, Tanaka N. Usefulness of a small-caliber, variable-stiffness colonoscope as a backup in patients with difficult or incomplete colonoscopy. *Am J Gastroenterol* 2004;99(10):1936-40.

Horiuchi A, Nakayama Y, Kato N, Ichise Y, Kajiyama M, Tanaka N. Hood-assisted colonoscopy is more effective in detection of colorectal adenomas than narrow-band imaging. *Clin Gastroenterol Hepatol* 2010;8(4):379-83.

Hosokawa O, Hattori M, Douden K, Hayashi H, Ohta K, Kaizaki Y. Difference in accuracy between gastroscopy and colonoscopy for detection of cancer. *Hepatogastroenterology* 2007;54(74):442-4.

Huang SW, Hsu CM, Jeng WJ, Yen TC, Su MY, Chiu CT. A Comparison of Positron Emission Tomography and Colonoscopy for the Detection of Advanced Colorectal Neoplasms in Subjects Undergoing a Health Check-Up. *PLoS ONE* 2013;8(7).

Huang Y, Ge W, London V, Li Q, Cai S, Zhang S, et al. Diagnostic inconsistency of faecal immunochemical tests for haemoglobin in population screening of colorectal cancer. *Clin Chem Lab Med* 2013;51(11):2173-80.

Janda M, Hughes KL, Auster JF, Leggett BA, Newman BM. Repeat participation in colorectal cancer screening utilizing fecal occult blood testing: a community-based project in a rural setting. *J Gastroenterol Hepatol* 2010;25(10):1661-7.

Johnson CD. The national CT colonography trial: Assessment of accuracy in participants 65 years of age and older. *Radiology* 2012;263(2):401-8.

Johnson CD, Chen MH, Toledano AY, Heiken JP, Dachman A, Kuo MD, et al. Accuracy of CT colonography for detection of large adenomas and cancers. *N Engl J Med* 2008;359(12):1207-17.

Jorgensen OD, Kronborg O, Fenger C, Rasmussen M. Influence of long-term colonoscopic surveillance on incidence of colorectal cancer and death from the disease in patients with precursors (adenomas). *Acta Oncol* 2007;46(3):355-60.

Kahi CJ, Anderson JC, Waxman I, Kessler WR, Imperiale TF, Li X, et al. High-definition chromocolonoscopy vs. high-definition white light colonoscopy for average-risk colorectal cancer screening. *Am J Gastroenterol* 2010;105(6):1301-7.

Kaminski MF, Bretthauer M, Zauber AG, Kuipers EJ, Adami HO, van Ballegooijen M, et al. The NordICC Study: rationale and design of a randomized trial on colonoscopy screening for colorectal cancer. *Endoscopy* 2012;44(7):695-702.

Kang HY, Kang HW, Kim SG, Kim JS, Park KJ, Jung HC, et al. Incidence and management of colonoscopic perforations in Korea. *Digestion* 2008;78(4):218-23.

Kapidzic A, Korfage IJ, Van DL, Van Roon AHC, Reijerink JCIY, Zauber AG, et al. Quality of life in participants of a CRC screening program. *Br J Cancer* 2012;107(8):1295-301.

Karl J, Wild N, Tacke M, Andres H, Garczarek U, Rollinger W, et al. Improved Diagnosis of Colorectal Cancer Using a Combination of Fecal Occult Blood and Novel Fecal Protein Markers. *Clin Gastroenterol Hepatol* 2008;6(10):1122-8.

Ker TS, Wasserberg N, Beart RWJ. Colonoscopic perforation and bleeding of the colon can be treated safely without surgery. *Am Surg* 2004;70(10):922-4.

Kim DH, Pooler BD, Weiss JM, Pickhardt PJ. Five year colorectal cancer outcomes in a large negative CT colonography screening cohort. *Eur Radiol* 2012;22(7):1488-94.

Kim YS, Kim N, Kim SH, Park MJ, Lim SH, Yim JY, et al. The efficacy of intravenous contrast-enhanced 16-row multidetector CT colonography for detecting patients with colorectal polyps in an asymptomatic population in Korea. *J Clin Gastroenterol* 2008;42(7):791-8.

Koss K, Maxton D, Jankowski JAZ. Faecal dimeric M2 pyruvate kinase in colorectal cancer and polyps correlates with tumour staging and surgical intervention. *Colorectal Dis* 2008;10(3):244-8.

Kouklakis G, Mpoumpouaris A, Gatopoulou A, Efraimidou E, Manolas K, Lirantzopoulos N. Endoscopic resection of large pedunculated colonic polyps and risk of postpolypectomy bleeding with adrenaline injection versus endoloop and hemoclip: a prospective, randomized study. *Surg Endosc* 2009;23(12):2732-7.

Ladabaum U, Allen J, Wandell M, Ramsey S. Colorectal cancer screening with blood-based biomarkers: Cost-effectiveness of methylated septin 9 DNA versus current strategies. *Cancer Epidemiol Biomarkers Prev* 2013;22(9):1567-76.

Laghi A, Iannaccone R, Trenna S, Mangiapane F, Sinibaldi G, Piacentini F, et al. Multislice spiral CT colonography in the evaluation of colorectal neoplasms. *Radiol Med* 2002;104(5-6):394-403.

Larsen IK, Grotmol T, Almendingen K, Hoff G. Lifestyle as a predictor for colonic neoplasia in asymptomatic individuals. *BMC Gastroenterol* 2006;6:5.

Lee TJW, Rutter MD, Blanks RG, Moss SM, Goddard AF, Chilton A, et al. Colonoscopy quality measures: Experience from the NHS Bowel Cancer Screening Programme. *Gut* 2012;61(7):1050-7.

Leufkens AM, Van Oijen MGH, Vleggaar FP, Siersema PD. Factors influencing the miss rate of polyps in a back-to-back colonoscopy study. *Endoscopy* 2012;44(5):470-5.

Leung WK, Lau JY, Suen BY, Wong GL, Chow DK, Lai LH, et al. Repeat-screening colonoscopy 5 years after normal baseline-screening colonoscopy in average-risk chinese: A prospective study. *Am J Gastroenterol* 2009;104(8):2028-34.

Liedenbaum MH, Venema HW, Stoker J. Radiation dose in CT colonography--trends in time and differences between daily practice and screening protocols. *Eur Radiol* 2008;18(10):2222-30.

Liles EG, Perrin N, Rosales AG, Feldstein AC, Smith DH, Mosen DM, et al. Change to FIT increased CRC screening rates: Evaluation of a US screening outreach program. *Am J Managed Care* 2012;18(10):588-95.

Ling BS, Schoen RE, Trauth JM, Wahed AS, Eury T, Simak DM, et al. Physicians encouraging colorectal screening: a randomized controlled trial of enhanced office and patient management on compliance with colorectal cancer screening. *Arch Intern Med* 2009;169(1):47-55.

Liou J-M, Lin J-T, Huang S-P, Chiu H-M, Wang H-P, Lee Y-C, et al. Screening for colorectal cancer in average-risk Chinese population using a mixed strategy with sigmoidoscopy and colonoscopy. *Dis Colon Rectum* 2007;50(5):630-40.

Lipkus IM, Samsa GP, Dement J, Skinner CS, Green LS, Pompeii L, et al. Accuracy of self-reports of fecal occult blood tests and test results among individuals in the carpentry trade. *Prev Med* 2003;37(5):513-9.

Loktionov A, Bandaletova T, Llewelyn AH, Dion C, Lywood HGG, Lywood RCG, et al. Colorectal cancer detection by measuring DNA from exfoliated colonocytes obtained by direct contact with rectal mucosa. *Int J Oncol* 2009;34(2):301-11.

Luning TH, Keemers-Gels ME, Barendregt WB, Tan ACIT, Rosman C. Colonoscopic perforations: a review of 30,366 patients. *Surg Endosc* 2007;21(6):994-7.

Macari M, Bini EJ, Xue X, Milano A, Katz SS, Resnick D, et al. Colorectal neoplasms: prospective comparison of thin-section low-dose multi-detector row CT colonography and conventional colonoscopy for detection. *Radiology* 2002;224(2):383-92.

Macari M, Bini EJ, Jacobs SL, Naik S, Lui YW, Milano A, et al. Colorectal polyps and cancers in asymptomatic average-risk patients: evaluation with CT colonography. *Radiology* 2004;230(3):629-36.

Mang T, Kolligs FT, Schaefer C, Reiser MF, Graser A. Comparison of diagnostic accuracy and interpretation times for a standard and an advanced 3D visualisation technique in CT colonography. *Eur Radiol* 2011;21(3):653-62.

Mang T, Hermosillo G, Wolf M, Bogoni L, Salganicoff M, Raykar V, et al. Time-efficient CT colonography interpretation using an advanced image-gallery-based, computer-aided "first-reader" workflow for the detection of colorectal adenomas. *Eur Radiol* 2012;22(12):2768-79.

Mayinger B, Neumann F, Kastner C, Degitz K, Hahn EG, Schwab D. Early detection of premalignant conditions in the colon by fluorescence endoscopy using local sensitization with hexaminolevulinate. *Endoscopy* 2008;40(2):106-9.

Morikawa T, Kato J, Yamaji Y, Wada R, Mitsushima T, Sakaguchi K, et al. Sensitivity of immunochemical fecal occult blood test to small colorectal adenomas. *Am J Gastroenterol* 2007;102(10):2259-64.

Multicentre Australian Colorectal-neoplasia Screening (MACS) Group. A comparison of colorectal neoplasia screening tests: a multicentre community-based study of the impact of consumer choice. *Med J Aust* 2006;184(11):546-50.

Myers RE, Bittner-Fagan H, Daskalakis C, Sifri R, Vernon SW, Cocroft J, et al. A randomized controlled trial of a tailored navigation and a standard intervention in colorectal cancer screening. *Cancer Epidemiol Biomarkers Prev* 2013;22(1):109-17.

Nelson DB. Colonoscopy versus double-contrast barium enema. *Gastroenterology* 2000;119(5):1402-3.

Nelson RL, Schwartz A. A survey of individual preference for colorectal cancer screening technique. *BMC cancer* 2004;4.

Ng SC, Ching JYL, Chan V, Wong MCS, Suen BY, Hirai HW, et al. Diagnostic accuracy of faecal immunochemical test for screening individuals with a family history of colorectal cancer. *Aliment Pharmacol Ther* 2013;38(7):835-41.

Nietert PJ, Jenkins RG, Nemeth LS, Ornstein SM. An application of a modified constrained randomization process to a practice-based cluster randomized trial to improve colorectal cancer screening. *Contemp Clin Trials* 2009;30(2):129-32.

Niv Y, Dickman R, Figer A, Abuksis G, Fraser G. Case-control study of screening colonoscopy in relatives of patients with colorectal cancer. *Am J Gastroenterol* 2003;98(2):486-9.

Ockenga J, Stuhmann M, Ballmann M, Teich N, Keim V, Dork T, et al. A randomized, controlled trial to assess a novel colorectal cancer screening strategy: The conversion strategy: A comparison of sequential sigmoidoscopy and colonoscopy with immediate conversion from sigmoidoscopy to colonoscopy in patients with an abnormal screening sigmoidoscopy. *Am J Gastroenterol* 2000;95(8):2074-9.

Oono Y, Iriguchi Y, Doi Y, Tomino Y, Kishi D, Oda J, et al. A retrospective study of immunochemical fecal occult blood testing for colorectal cancer detection. *Clin Chim Acta* 2010;411(11-12):802-5.

Ore L, Hagoel L, Lavi I, Rennert G. Screening with faecal occult blood test (FOBT) for colorectal cancer: assessment of two methods that attempt to improve compliance. *Eur J Cancer Prev* 2001;10(3):251-6.

Pickhardt PJ, Choi JR, Hwang I, Pineau BC. Screening computed tomographic colonography in asymptomatic adults: As good as colonoscopy? *Evid -Based Gastroenterol* 2004;5(3):82-3.

Pilz JB, Portmann S, Peter S, Beglinger C, Degen L. Colon Capsule Endoscopy compared to Conventional Colonoscopy under routine screening conditions. *BMC Gastroenterol* 2010;10:66.

Pinsky PF, Schoen RE, Weissfeld JL, Kramer B, Hayes RB, Yokochi L, et al. Variability in flexible sigmoidoscopy performance among examiners in a screening trial. *Clin Gastroenterol Hepatol* 2005;3(8):792-7.

Potter MB, Namvargolian Y, Hwang J, Walsh JME. Improving colorectal cancer screening: a partnership between primary care practices and the American Cancer Society. *J Cancer Educ* 2009;24(1):22-7.

Potter MB, Phengrasamy L, Hudes ES, McPhee SJ, Walsh JME. Offering annual fecal occult blood tests at annual flu shot clinics increases colorectal cancer screening rates. *Ann Fam Med* 2009;7(1):17-23.

Potter MB, Walsh JME, Yu TM, Gildengorin G, Green LW, McPhee SJ. The effectiveness of the FLU-FOBT program in primary care a randomized trial. *Am J Prev Med* 2011;41(1):9-16.

Potter MB, Ackerson LM, Gomez V, Walsh JME, Green LW, Levin TR, et al. Effectiveness and reach of the FLU-FIT program in an integrated health care system: a multisite randomized trial. *Am J Public Health* 2013;103(6):1128-33.

Pox CP, Altenhofen L, Brenner H, Theilmeier A, Von Stillfried D, Schmiegel W. Efficacy of a nationwide screening colonoscopy program for colorectal cancer. *Gastroenterology* 2012;142(7):1460-7.

Quintero E, Castells A, Bujanda L, Cubiella J, Salas D, Lanas A, et al. Colonoscopy versus fecal immunochemical testing in colorectal-cancer screening. *N Engl J Med* 2012;366(8):697-706.

Rae LC, Gibberd RW. Survival of patients with colorectal cancer detected by a community screening program. *Med J Aust* 2000;172(1):13-5.

Rasmussen M, Fenger C, Kronborg O. Diagnostic yield in a biennial Hemocult-II screening program compared to a once-only screening with flexible sigmoidoscopy and Hemocult-II. *Scand J Gastroenterol* 2003;38(1):114-8.

Rockey DC, Paulson E, Niedzwiecki D, Davis W, Bosworth HB, Sanders L, et al. Analysis of air contrast barium enema, computed tomographic colonography, and colonoscopy: Prospective comparison. *Lancet* 2005;365(9456):305-11.

Rogal SS, Pinsky PF, Schoen RE. Relationship between detection of adenomas by flexible sigmoidoscopy and interval distal colorectal cancer. *Clin Gastroenterol Hepatol* 2013;11(1):73-8.

Rohlin A, Wernersson J, Engwall Y, Wiklund L, Bjork J, Nordling M. Parallel sequencing used in detection of mosaic mutations: Comparison with four diagnostic DNA screening techniques. *Hum Mutat* 2009;30(6):1012-20.

Rokkas T, Papaxoinis K, Triantafyllou K, Ladas SD. A meta-analysis evaluating the accuracy of colon capsule endoscopy in detecting colon polyps. *Gastrointest Endosc* 2010;71(4):792-8.

Roy HK, Turzhitsky V, Kim Y, Goldberg MJ, Watson P, Rogers JD, et al. Association between rectal optical signatures and colonic neoplasia: Potential applications for screening. *Cancer Res* 2009;69(10):4476-83.

Roy HK, Gomes AJ, Ruderman S, Bianchi LK, Goldberg MJ, Stoyneva V, et al. Optical measurement of rectal microvasculature as an adjunct to flexible sigmoidoscopy: Gender-specific implications. *Cancer Prev Res* 2010;3(7):844-51.

Rozen P, Levi Z, Hazazi R, Waked A, Vilkin A, Maoz E, et al. Identification of colorectal adenomas by a quantitative immunochemical faecal occult blood screening test depends on adenoma characteristics, development threshold used and number of tests performed. *Aliment Pharmacol Ther* 2009;29(8):906-17.

Russmann S, Lamerato L, Marfatia A, Motsko SP, Pezzullo JC, Olds G, et al. Risk of impaired renal function after colonoscopy: A cohort study in patients receiving either oral sodium phosphate or polyethylene glycol. *Am J Gastroenterol* 2007;102(12):2655-63.

Saar B, Meining A, Beer A, Settles M, Helmberger H, Frimberger E, et al. Prospective study on bright lumen magnetic resonance colonography in comparison with conventional colonoscopy. *Br J Radiol* 2007;80(952):235-41.

Sali L, Grazzini G, Carozzi F, Castiglione G, Falchini M, Mallardi B, et al. Screening for colorectal cancer with FOBT, virtual colonoscopy and optical colonoscopy: Study protocol for a randomized controlled trial in the Florence district (SAVE study). *Trials* 2013;14(1).

Sali L, Falchini M, Bonanomi AG, Castiglione G, Ciatto S, Mantellini P, et al. CT colonography after incomplete colonoscopy in subjects with positive faecal occult blood test. *World J Gastroenterol* 2008;14(28):4499-504.

Sanaka MR, Deepinder F, Thota PN, Lopez R, Burke CA. Adenomas are detected more often in morning than in afternoon colonoscopy. *Am J Gastroenterol* 2009;104(7):1659-65.

Saraya T, Ikematsu H, Fu KI, Tsunoda C, Yoda Y, Oono Y, et al. Evaluation of complications related to therapeutic colonoscopy using the bipolar snare. *Surg Endosc* 2012;26(2):533-40.

Schoofs N, Deviere J, Van Gossum A. PillCam colon capsule endoscopy compared with colonoscopy for colorectal tumor diagnosis: a prospective pilot study. *Endoscopy* 2006;38(10):971-7.

Scott RG, Edwards JT, Fritschi L, Foster NM, Mendelson RM, Forbes GM. Community-based screening by colonoscopy or computed tomographic colonography in asymptomatic average-risk subjects. *Am J Gastroenterol* 2004;99(6):1145-51.

Senore C, Ederle A, Fantin A, Andreoni B, Bisanti L, Grazzini G, et al. Acceptability and side-effects of colonoscopy and sigmoidoscopy in a screening setting. *J Med Screen* 2011;18(3):128-34.

Shastri YM, Loitsch S, Nowak R, Povse N, Stein J. Prospective comparative evaluation of an office-based rapid immunological test with a Guaiac-based fecal occult blood test for colorectal cancer screening in general population with average-risk. *Clin Lab* 2008;54(9-10):385-7.

Siddiqui AA, Sifri R, Hyslop T, Andrel J, Rosenthal M, Vernon SW, et al. Race and response to colon cancer screening interventions. *Prev Med* 2011;52(3-4):262-4.

Singer RB. Fecal occult blood testing and the incidence of colorectal cancer. *J Insur Med* 2002;34(2):89-91.

Stegeman I, De Wijkerslooth TR, Stoop EM, van Leerdam M, van Ballegooijen M, Kraaijenhagen RA, et al. Risk factors for false positive and for false negative test results in screening with fecal occult blood testing. *Int J Cancer* 2013;133(10):2408-14.

Stoop EM, de Haan MC, de Wijkerslooth TR, Bossuyt PM, van Ballegooijen M, Nio CY, et al. Participation and yield of colonoscopy versus non-cathartic CT colonography in population-based screening for colorectal cancer: a randomised controlled trial. *Lancet Oncol* 2012;13(1):55-64.

Summers RM, Yao J, Pickhardt PJ, Franaszek M, Bitter I, Brickman D, et al. Computed tomographic virtual colonoscopy computer-aided polyp detection in a screening population. *Gastroenterology* 2005;129(6):1832-44.

Suzuki K, Yoshida H, Nappi J, Armato III SG, Dachman AH. Mixture of expert 3D massive-training ANNs for reduction of multiple types of false positives in CAD for detection of polyps in CT colonography. *Med Phys* 2008;35(2):694-703.

Taupin D, Chambers SL, Corbett M, Shadbolt B. Colonoscopic screening for colorectal cancer improves quality of life measures: a population-based screening study. *Health Qual Life Outcomes* 2006;4:82.

Taylor KL, Shelby R, Gelmann E, McGuire C. Quality of life and trial adherence among participants in the prostate, lung, colorectal, and ovarian cancer screening trial. *J Natl Cancer Inst* 2004;96(14):1083-94.

Thiis-Evensen E, Hoff GS, Sauar J, Majak BM, Vatn MH. The effect of attending a flexible sigmoidoscopic screening program on the prevalence of colorectal adenomas at 13-year follow-up. *Am J Gastroenterol* 2001;96(6):1901-7.

van Rossum LGM, van Rijn AF, van Oijen MGH, Fockens P, Laheij RJF, Verbeek ALM, et al. False negative fecal occult blood tests due to delayed sample return in colorectal cancer screening. *Int J Cancer* 2009;125(4):746-50.

van Rossum LGM, van Rijn AF, Verbeek ALM, van Oijen MGH, Laheij RJF, Fockens P, et al. Colorectal cancer screening comparing no screening, immunochemical and guaiac fecal occult blood tests: a cost-effectiveness analysis. *Int J Cancer* 2011;128(8):1908-17.

van Turenhout ST, van Rossum LGM, Oort FA, Laheij RJF, van Rijn AF, Terhaar sive Droste JS, et al. Similar fecal immunochemical test results in screening and referral colorectal cancer. *World J Gastroenterol* 2012;18(38):5397-403.

Wada Y, Kudo Se, Misawa M, Ikehara N, Hamatani S. Vascular pattern classification of colorectal lesions with narrow band imaging magnifying endoscopy. *Digestive endoscopy : official journal of the Japan Gastroenterological Endoscopy Society* 2011;23 Suppl 1:106-11.

Wallace PM, Suzuki R. Regional, racial, and gender differences in colorectal cancer screening in middle-aged African-Americans and Whites. *J Cancer Educ* 2012;27(4):703-8.

Walsh JME, Salazar R, Nguyen TT, Kaplan C, Nguyen LK, Hwang J, et al. Healthy colon, healthy life: a novel colorectal cancer screening intervention. *Am J Prev Med* 2010;39(1):1-14.

Wang YR, Cangemi JR, Loftus EVJ, Picco MF. Risk of colorectal cancer after colonoscopy compared with flexible sigmoidoscopy or no lower endoscopy among older patients in the United States, 1998-2005. *Mayo Clin Proc* 2013;88(5):464-70.

Warren JL, Klabunde CN, Mariotto AB, Meekins A, Topor M, Brown ML, et al. Adverse events after outpatient colonoscopy in the Medicare population. *Ann Intern Med* 2009;150(12):849-W152.

Whyte S, Chilcott J, Halloran S. Reappraisal of the options for colorectal cancer screening in England. *Colorectal Dis* 2012;14(9):e547-e561.

Wilschut JA, Hol L, Dekker E, Jansen JB, van Leerdam ME, Lansdorp-Vogelaar I, et al. Cost-effectiveness analysis of a quantitative immunochemical test for colorectal cancer screening. *Gastroenterology* 2011;141(5):1648-55.

Wood JJ, Foy CJW, Valori R, Lucarotti ME, Fowler AL, Dowler K, et al. Does use of a colonoscopy imaging device improve performance? A cohort study. *Surg Endosc* 2012;26(12):3616-21.

Yee J, Akerkar GA, Hung RK, Steinauer-Gebauer AM, Wall SD, Mcquaid KR. Colorectal neoplasia: performance characteristics of CT colonography for detection in 300 patients. *Radiology* 2001;219(3):685-92.

Yen AW, Leung JW, Leung FW. A novel method with significant impact on adenoma detection: Combined water-exchange and cap-assisted colonoscopy. *Gastrointest Endosc* 2013;77(6):944-8.

Yoshinaga M, Watabe R, Yanagisawa J, Harada N, Motomura S, Nawata H, et al. The interval between flexible sigmoidoscopy screening examinations can be expanded beyond five years. *Hepatogastroenterology* 2001;48(38):437-9.

Young PE, Gentry AB, Cash BD. The utility of flexible sigmoidoscopy after a computerized tomographic colonography revealing only rectosigmoid lesions. *Aliment Pharmacol Ther* 2008;27(6):520-7.

Zhang GJ, Zhou T, Liu ZL, Tian HP, Xia SS. Plasma miR-200c and miR-18a as potential biomarkers for the detection of colorectal carcinoma. *Molecular and Clinical Oncology* 2013;1(2):379-84.

Zou H, Taylor WR, Harrington JJ, Hussain FTN, Cao X, Loprinzi CL, et al. High Detection Rates of Colorectal Neoplasia by Stool DNA Testing With a Novel Digital Melt Curve Assay. *Gastroenterology* 2009;136(2):459-70.

Zubarik R, Ganguly E, Benway D, Ferrentino N, Moses P, Vecchio J. Procedure-related abdominal discomfort in patients undergoing colorectal cancer screening: a comparison of colonoscopy and flexible sigmoidoscopy. *Am J Gastroenterol* 2002;97(12):3056-61.

Excluded Wrong Study Design

Azuara D, Rodriguez-Moranta F, De OJ, Soriano-Izquierdo A, Mora J, Guardiola J, et al. Novel methylation panel for the early detection of colorectal tumors in stool DNA. *Clin Colorectal Cancer* 2010;9(3):168-76.

Baxter NN, Goldwasser MA, Paszat LF, Saskin R, Urbach DR, Rabeneck L. Association of colonoscopy and death from colorectal cancer. *Ann Intern Med* 2009;150(1):1-8.

Brenner H, Haug U, Arndt V, Stegmaier C, Altenhofen L, Hoffmeister M. Low risk of colorectal cancer and advanced adenomas more than 10 years after negative colonoscopy. *Gastroenterology* 2010;138(3):870-6.

Cotterchio M, Manno M, Klar N, McLaughlin J, Gallinger S. Colorectal screening is associated with reduced colorectal cancer risk: a case-control study within the population-based Ontario Familial Colorectal Cancer Registry. *Cancer causes & control : CCC* 2005;16(7):865-75.

Faiss S. The missed colorectal cancer problem. *Dig Dis* 2011;29 Suppl 1:60-3.

Gillespie JS, Kelly BE. Double contrast barium enema and colorectal carcinoma: sensitivity and potential role in screening. *Ulster Med J* 2001;70(1):15-8.

Hamashima C, Sobue T, Muramatsu Y, Saito H, Moriyama N, Kakizoe T. Comparison of observed and expected numbers of detected cancers in the research center for cancer prevention and screening program. *Jpn J Clin Oncol* 2006;36(5):301-8.

Jacob BJ, Moineddin R, Sutradhar R, Baxter NN, Urbach DR. Effect of colonoscopy on colorectal cancer incidence and mortality: an instrumental variable analysis. *Gastrointest Endosc* 2012;76(2):355-64.

Jacob BJ, Sutradhar R, Moineddin R, Baxter NN, Urbach DR. Methodological approaches to population based research of screening procedures in the presence of selection bias and exposure measurement error: colonoscopy and colorectal cancer outcomes in Ontario. *BMC Med Res Methodol* 2013;13:59.

Jafri SM, Arora A. Silent perforation: an iatrogenic complication of colonoscopy. *Surg Laparosc Endosc Percutan Tech* 2007;17(5):452-4.

Kahi CJ, Imperiale TF, Juliar BE, Rex DK. Effect of screening colonoscopy on colorectal cancer incidence and mortality. *Clin Gastroenterol Hepatol* 2009;7(7):770-11.

Lamah M, Norris J, Caffarey SM, Broughton M, Marks CG. Effect of faecal occult blood testing on colorectal cancer mortality in the surveillance of subjects at moderate risk of colorectal neoplasia: A case-control study. *Int J Colorectal Dis* 2001;16(5):313-7.

Lee KJ, Inoue M, Otani T, Iwasaki M, Sasazuki S, Tsugane S, et al. Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: a prospective cohort study in Japan. *Cancer Detect Prev* 2007;31(1):3-11.

Lejeune C, Arveux P, Dancourt V, Bejean S, Bonithon-Kopp C, Faivre J. Cost-effectiveness analysis of fecal occult blood screening for colorectal cancer. *Int J Technol Assess Health Care* 2004;20(4):434-9.

Li S, Nie Z, Li N, Li J, Zhang P, Yang Z, et al. Colorectal cancer screening for the natural population of Beijing with sequential fecal occult blood test: A multicenter study. *Chin Med J* 2003;116(2):200-2.

Libby G, Brewster DH, McClements PL, Carey FA, Black RJ, Birrell J, et al. The impact of population-based faecal occult blood test screening on colorectal cancer mortality: a matched cohort study. *Br J Cancer* 2012;107(2):255-9.

Macafee DAL, Waller M, Whynes DK, Moss S, Scholefield JH. Population screening for colorectal cancer: the implications of an ageing population. *Br J Cancer* 2008;99(12):1991-2000.

Manser CN, Bachmann LM, Brunner J, Hunold F, Bauerfeind P, Marbet UA. Colonoscopy screening markedly reduces the occurrence of colon carcinomas and carcinoma-related death: a closed cohort study. *Gastrointest Endosc* 2012;76(1):110-7.

McClements PL, Madurasinghe V, Thomson CS, Fraser CG, Carey FA, Steele RJC, et al. Impact of the UK colorectal cancer screening pilot studies on incidence, stage distribution and mortality trends. *Cancer Epidemiol* 2012;36(4):e232-e242.

Mulder SA, Van Soest EM, Dieleman JP, van Rossum LGM, Ouwendijk RJT, Van Leerdam ME, et al. Exposure to colorectal examinations before a colorectal cancer diagnosis: A case-control study. *Eur J Gastroenterol Hepatol* 2010;22(4):437-43.

Nakajima M, Saito H, Soma Y, Sobue T, Tanaka M, Munakata A. Prevention of advanced colorectal cancer by screening using the immunochemical faecal occult blood test: A case-control study. *Br J Cancer* 2003;89(1):23-8.

Newcomb PA, Storer BE, Morimoto LM, Templeton A, Potter JD. Long-term efficacy of sigmoidoscopy in the reduction of colorectal cancer incidence. *J Natl Cancer Inst* 2003;95(8):622-5.

Niv Y, Lev-El M, Fraser G, Abuksis G, Tamir A. Protective effect of faecal occult blood test screening for colorectal cancer: worse prognosis for screening refusers. *Gut* 2002;50(1):33-7.

Petersen CR, Adamsen S, Gocht-Jensen P, Arnesen RB, Hart-Hansen O. Splenic injury after colonoscopy. *Endoscopy* 2008;40(1):76-9.

Philippart P, Takeh H, Brandelet B, de Koster E, Mendes da Costa P. Presacral hemorrhage: a rare complication of colonoscopy. *Hepatogastroenterology* 2005;52(66):1725-6.

Quallick MR, Brown WR. Rectal perforation during colonoscopic retroflexion: a large, prospective experience in an academic center. *Gastrointest Endosc* 2009;69(4):960-3.

Rabeneck L, Lewis JD, Paszat LF, Saskin R, Stukel TA. Risk of proximal and distal colorectal cancer following flexible sigmoidoscopy: a population-based cohort study. *Am J Gastroenterol* 2008;103(8):2075-82.

Rozen P, Shabtai EI, Liphshitz I, Barchana M. Risk for colorectal cancer in elderly persons and possible methodologies for their screening. *Eur J Gastroenterol Hepatol* 2011;23(5):431-7.

Sharaf RN, Ladabaum U. Comparative effectiveness and cost-effectiveness of screening colonoscopy vs. sigmoidoscopy and alternative strategies. *Am J Gastroenterol* 2013;108(1):120-32.

Singh H, Nugent Z, Demers AA, Kliewer EV, Mahmud SM, Bernstein CN. The reduction in colorectal cancer mortality after colonoscopy varies by site of the cancer. *Gastroenterology* 2010;139(4):1128-37.

Suttie SA, Shaikh IA, Amin AI, Daniel T, Yalamarthi S. The effect of a screening programme on the outcome of colorectal cancer. *Colorectal Dis* 2012;14(8):937-42.

Trecca A, Gaj F, Pasciuto A, Serafini S, Bella A. Advanced diagnostic tools: Virtual colonoscopy and magnifying chromoendoscopy. *Tech Coloproctol* 2004;8(SUPPL. 2):S279-S282.

Trombold J, Farmer RW, McCafferty M. The impact of colorectal cancer screening in a veteran hospital population. *Am Surg* 2013;79(3):296-300.

Ventura L, Zappa M, Carreras G, Ciatto S, Grazzini G. What is the best screening strategy to detect advanced colorectal adenomas? Simulation from ongoing Italian screening experiences. *Tumori* 2011;97(5):547-50.

Vijan S, Hwang EW, Hofer TP, Hayward RA. Which colon cancer screening test? A comparison of costs, effectiveness, and compliance. *Am J Med* 2001;111(8):593-601.