




Breast Cancer Screening for Women Not at Increased Risk

Updated Task Force Recommendations for Women

For women aged 60–69 years: we conditionally recommend *screening* with mammography every two to three years.

- The balance of benefits and harms is *more favourable* in this group.
- This recommendation is conditional because some women may wish to not be screened if they are concerned about potential harms.
- Women in this age group should have a discussion with their health care provider to decide if screening is best for them.

Screening 1000 women aged 60–69 over 7 years



With screening:

256 women will have a false positive test result

35 women will have an unnecessary biopsy

21 women will be diagnosed with breast cancer. Among these 21 women:

No reliable data on the number of women who were treated for a breast cancer that would have never caused a problem

1 breast cancer death will be prevented

1087 women in this age group would need to be screened to prevent one death

Who do these recommendations *not* apply to?

These recommendations don't apply to anyone at increased risk of breast cancer, such as those with a personal or family history of breast cancer.

What is screening?

Screening is done to attempt to detect potential disease or illness in people who do not have any signs or symptoms of disease.

What is a mammogram?

It is an x-ray of the breast(s) to identify potential cancer.

Why is shared decision making important?

Screening is a personal decision. It is important to weigh the benefits and harms of screening for women in your age group (as shown below) with your health care provider to decide what is best for you.

What are the harms of screening for breast cancer?

Overdiagnosis - Not all breast cancers will cause harm to a woman in her lifetime. With screening, some women will be diagnosed with a cancer that would not have caused them a problem in their lifetime; this is called 'overdiagnosis' and leads to unnecessary treatment.

False positives - A false positive test occurs in someone who tested positive (abnormal mammography) but who ultimately is shown not to have cancer. It can lead to additional testing, including biopsy, and may cause psychological and physical harm.

In general, harms of screening are greater in younger women and decrease with age.

What are the benefits of screening for breast cancer?

There is evidence that shows that screening lowers a woman's risk of dying from breast cancer. In general, the benefits of screening increase with age.

This tool is not a decision aid but is intended to be one step in the shared decision making process.